




Health benefits of xylitol

Asma Gasmi Benahmed¹ · Amin Gasmi² · Maria Arshad² · Mariia Shanaida³ · Roman Lysiuk^{4,5} · Massimiliano Peana⁶ · Irena Pshyk-Titko^{5,7} · Stepan Adamiv⁸ · Yurii Shanaida³ · Geir Bjørklund⁹ 

Received: 9 April 2020 / Revised: 23 May 2020 / Accepted: 31 May 2020 / Published online: 7 July 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Abstract

Many diseases, including caries, chronic inflammatory diseases, diabetes, and obesity, are associated with uncontrolled sugar consumption. Artificial sweeteners are commonly used in food and pharmaceutical industries as sugar substitutes for the prevention of several dental and body diseases; they also have a favorable impact on body weight as they may help to restrict simple sugar consumption. Xylitol is a sugar alcohol that is commonly used as a sweetener. It can be found naturally or artificially prepared mainly from plant materials chemically or by fermentation of hemicelluloses from agricultural biomass by yeast or bacteria strains. This polyol has a significant antiplaque effect on teeth surface and can reduce the gingival inflammation; it is being used as a preventive agent for dental caries due to decreasing the growth levels of pathogenic *Streptococcus mutans* and *Streptococcus sanguis* at the very early stages. Xylitol can bind with calcium ion leading to consequent remineralization of teeth enamel; it is also able to prevent osteoporosis. This polyol can treat respiratory tract and middle ear diseases due to its antibacterial and anti-inflammatory potential and prevent some diseases which cannot be cured through antibiotics or surgery. Xylitol can reduce constipation, diabetes, obesity, and other body syndromes or illnesses; it has also revealed its stimulating effect on digestion and immune system. However, it can produce some side effects such as irritable bowel syndrome, diarrhea, nephrolithiasis, etc., when consumed in excessive amounts. Different vehicles are used for delivering the xylitol into the human body, but chewing gums occupy a leading position. The present review is devoted to comprehensive analyses of the positive and negative effects of this polyol on human health.

Key Points

- The health benefits of xylitol are not limited to oral hygiene.
- Xylitol efficiently stimulates the immune system, digestion, lipid and bone metabolism.
- Xylitol helps in glycemic and obesity control; reduces ear and respiratory infections.
- Xylitol treats diseases that cannot be cured through antibiotics or by surgery.

Keywords Xylitol · Artificial sweetener · Oral microbiota · Caries · Metabolic disease · Health care · Preventive effect

✉ Geir Bjørklund
bjorklund@conem.org

¹ Académie Internationale de Médecine Dentaire Intégrative, Paris, France

² Société Francophone de Nutrithérapie et de Nutriginétique Appliquée, Villeurbanne, France

³ I. Horbachevsky Ternopil National Medical University, Ternopil, Ukraine

⁴ Danylo Halytsky Lviv National Medical University, Lviv, Ukraine

⁵ CONEM Ukraine Life Science Research Group, Danylo Halytsky Lviv National Medical University, Lviv, Ukraine

⁶ Department of Chemistry and Pharmacy, University of Sassari, Sassari, Italy

⁷ Dental Clinics Dr. Irena Pshyk-Titko, Lviv, Ukraine

⁸ Odessa National Medical University, Odessa, Ukraine

⁹ Council for Nutritional and Environmental Medicine (CONEM), Toften 24, 8610 Mo i Rana, Norway

Introduction

The global increase in sugar consumption causes health problems, including exacerbation of caries, gingivitis, metabolic syndrome, obesity, and diabetes mellitus (Burt 2006; Wolnerhanssen and Meyer-Gerspach 2019; Wolnerhanssen et al. 2019). This has driven consumers to demand healthier sweeteners such as sugar alcohols. These polyols are present in many natural products, such as fruits and vegetables, and are also added to foods and pharmaceuticals as artificial sweeteners. Commonly used polyols are xylitol, sorbitol, erythritol, maltitol, lactitol, mannitol, and isomalt (Rice et al. 2019). They are widely used in the manufacturing of food and pharmaceuticals because of their favorable physicochemical properties (Ortiz et al. 2013). Xylitol is a 5-carbon sugar alcohol, white crystalline carbohydrate, and commonly used as an artificial sweetener (Fig. 1). It was identified in the late nineteenth century, naturally present in numerous edible plants (including fruits and berries) and mushrooms. It is also artificially manufactured in the industry from xylan-rich plant material, such as beechwood and birch wood (El-Marakby et al. 2017).

According to the World Health Organization, dental caries is a widely spread disease linked to the prevalence of pathogenic microorganisms (mainly *Streptococcus mutans* and *Streptococcus sobrinus*) in the oral cavity and with the quantity of fermentable carbohydrates ingested (Petersen 2004). Strategies to maximize caries prevention, of course, foresee the use of sugar substitutes (Mäkinen 2011). Xylitol has a great potential to inhibit or reduce the growth of oral pathogenic bacteria. Its antibacterial potential has made it famous in diverse preventive dental medicines or products. Clinical studies revealed that xylitol, as physiologic sugar alcohol of natural origin, can be used as an effective caries-limiting sweetener (Honkala et al. 2014; Mäkinen 2011; Silva et al. 2009).




Fig. 1 The structure of xylitol ((2R,3R,4S)-pentane-1,2,3,4,5-pentol)

Throughout World War II, the Scandinavian countries faced a severe sugar shortage. Then, they used xylitol as a substitute for sugar. After that, dental professionals started using xylitol to prevent dental caries and to improve oral health.

In previous decades, researchers mainly were focused on the influences of xylitol and other artificial sweeteners on oral health care while the antihyperglycaemic effects have only been studied recently (Wolnerhanssen et al. 2019). Xylitol has many beneficial characteristics for humans (Fig. 2) (Chukwuma and Islam 2017); for example, it is involved in the release of a minimal amount of insulin. It is considered a good sugar substitute for the people who have diabetes and are on a very low carbohydrate diet to overcome the high glycaemic index in their bloodstream. Xylitol does not require insulin hormone to enter in the cells; that is why it is considered as a good energy source for people with diabetes. It is found in many food items, particularly labeled as a “low-carb” diet. It is considered a non-toxic compound in mammals with the exception of dogs.


Nowadays, xylitol has been studied worldwide and accepted as an artificial sweetener globally (Peterson 2013). Xylitol and other polyols are approved for use in a wide range of foods, and many of them have common designations, for instance: xylitol (E 967), isomalt (E 953); lactitol (E 966); maltitol (E 965); mannitol (E 421); sorbitol (E 420) (Tennant 2014).

What is xylitol?

The word “Xylitol” derived from the Greek word “Xylo” means wood, and “itol” is the suffix denotes the sugar alcohols. Xylitol is an alcoholic compound that is naturally present in plants and is produced by some bacteria and fungi (Ortiz et al. 2013). It is generally classified as sugar alcohol or polyalcohol. Its chemical formula is consisting of 5 carbon skeleton $\text{CH}_2\text{OH}(\text{CHOH})_3\text{CH}_2\text{OH}$ (Fig. 1) (Granstrom et al. 2007). Physically, it is comprised of white solid, crystalline, or granular structure that is soluble in water.

Xylitol occurs in a very low quantity in fruits, vegetables, and plant raw materials, as well as its precursor xylose (Olennikov et al. 2009; Shanaida et al. 2017; Ur-Rehman et al. 2015). It is mainly obtained commercially from birch wood or other plants that are used as raw material to manufacture food additives and medicinal products (Riley et al. 2015). Recovery of waste constituents from plants manufacturing have both economic and social importance (Fierascu et al. 2020).

Xylitol can be produced chemically or by biotechnological processes (Fig. 2) (Rice et al. 2019). Chemical production uses extensive machinery and expensive chemicals, thus increasing the basic cost. The biotechnological method utilizes

Fig. 2 Xylitol production and health benefits

agricultural wastes, which offer the possibilities of economic production of xylitol. The precursor xylose is produced from agricultural biomass by enzymatic hydrolysis and can be converted to xylitol primarily by yeast or bacteria strains (Carneiro et al. 2019; Felipe Hernandez-Perez et al. 2019; Jain and Mulay 2014; Stoklosa et al. 2019; Ur-Rehman et al. 2015). Hemicellulosic hydrolysates are the primary raw materials for the production of xylitol (Felipe Hernandez-Perez et al. 2019; Sene et al. 2011; Yuan et al. 2020). Recombinant *Escherichia coli* has been revealed the better producer of xylitol (Abd Rahman et al. 2020; Yuan et al. 2020).

Biotechnologically engineered *Zymomonas mobilis* ATCC ZW658 strain showed the ability to co-utilize glucose and xylose, exhibiting a 1.65-fold increase in the xylose utilization rate compared with the parent one (Sarkar et al. 2020). There has been constructed a strain of recombinant yeast *Saccharomyces cerevisiae* efficiently producing xylitol using wheat stalk hydrolysate as a source of xylose (Reshamwala and Lali 2020). Biosynthesis of value-added functional sugars, including xylitol using biotechnologically engineered yeast *Yarrowia lipolytica* as sustainable carbon sources, are also among promising trends in the latest investigations (Bilal et al. 2020; Xu et al. 2019).

Xylitol vehicles

Xylitol is considered as a safe and efficient tooth decay preventive substance when used regularly, although its uses have

been restricted due to the absence of formulations. Xylitol can, among others, be taken from candies, chewing gums, sweets, and syrups (Jones 2000). Substantial research proposes that 5–6 g and a minimum of three exposures per day from candies or chewing gums are required for a better clinical effect (Milgrom et al. 2009a). Most xylitol clinical trials include almost entirely lozenges or chewing gums and tooth decay assessed in school-going children's teeth (Lynch and Milgrom 2003). To get the effective results of xylitol, almost two vehicles are adequate and safe for toddlers. Many studies have been conducted to evaluate the effective results of xylitol by using topical xylitol syrups during primary teeth eruption in toddlers (Milgrom et al. 2009b).

The common modality for the delivery of xylitol is chewing gum (Soderling 2009b). It enhances the process of cleaning away acid and acts as remineralized for tooth by uptaking calcium phosphate. The prescribed time for chewing the gum is 20 min after eating. The study also showed that the regular consumption of xylitol gummy bear canapés or snacks helps reduce germs responsible for producing dental caries in school-going children. It is adequately reported that eating gummy bear snacks consisting of xylitol is responsible for reducing mutans streptococci growth levels (Ganter et al. 2020; Nayak et al. 2014).

Mouth rinses comprised of xylitol can reduce the development of biofilms. They also reduce the growth levels of *S. sanguis* and *S. mutans* at the very early stages. Among both, the *S. mutans*' colonies are highly sensitive to xylitol mouth rinses. In the USA, xylitol is used in small nonclinical

quantities as an artificial sweetener in a gimmick or in several food items and in children's supplements, including vitamins (El-Marakby et al. 2017). Meanwhile, there has also been a debate in previous literature about whether xylitol-containing toothpaste at lower doses is most effective.

However, the synergistic effect of fluoride and xylitol cannot be ruled out. Researches have been conducted on new xylitol vehicles, for example, xylitol-releasing dummy or pediatric xylitol syrups. (Milgrom et al. 2009a). Fluoride toothpaste with 10% xylitol reduced caries by 13% when compared to a fluoride-only toothpaste analyzed in 4216 children (Duane 2015). Chewing sugar-free gums reduced caries increment more effectively in comparison to non-chewing controls using alternatives such as rinses, tablets, and candies (Newton et al. 2019). Hence, this substance can be used as an active element in gums, toothpaste, and antimicrobial mouthwashes, especially in patients with candidiasis (Talattof et al. 2018).

Artificially sweetened soft drinks have appeared as an alternative sweetness without changes in caloric intake because most of the sugars are obtained from such drinks; also, sweeteners are often used to replace sugars in various yogurts, puddings, and ice cream (Sylvetsky et al. 2011). Inclusion of a xylitol drinking water additive for seven months reduced the tooth plaque in dogs on 5.1% and the calculus accumulation on 14.9% (Lowe and Anthony 2020). Natural sweeteners have a positive influence on body weight as they may help to restrict simple sugar consumption (Mooradian et al. 2017). It should be noted that xylitol present in beverages causes side effects (such as diarrhea) at lower levels than when present in solid food (Mäkinen 2016).

Antiplaque, antigingivitic, and remineralizing properties of xylitol

Biofilm formation is a natural process in the oral cavity but needs to be under the control of regular brushing to prevent the development of periodontal diseases and caries (Verkaik et al. 2011). Dental plaque is a biofilm produced by bacteria, salivary proteins, and food substances (Badet et al. 2008).

Lactobacillus, *S. mutans*, and *S. sobrinus*, and *Actinomyces viscosus* are four major components of the oral cavity microflora producing acid through the fermentation of sugars from consumed food (Thabuis et al. 2013). This acid can induce a partial dissolution of minerals in tooth enamel, followed by the formation of a dental cavity infected by oral bacteria. As bacteria of the oral cavity do not ferment polyols, their use does not increase the acid content there.

Xylitol has significant antiplaque effect on teeth; thus, it is being used as a preventive agent for dental caries. Xylitol cannot be fermented by the microorganisms of dental plaque (Takahashi et al. 2003). Studies showed evidence that oral flora cannot metabolize xylitol. The use of xylitol daily has

shown a severe reduction of colonies of *S. mutans* in dental plaque. Xylitol decreases the growth of bacteria. *S. mutans* carry the xylitol sugar inside the cell during an energy-consuming cycle, and this phenomenon is responsible for inhibition of growth and, ultimately, inhibition of plaque (Alves et al. 2013). Xylitol is further converted into the xylitol-5-phosphate by phosphoenolpyruvate. This phenomenon results in the production of intracellular vacuoles and degradation of the cell membrane. Therefore, *S. mutans* bacteria are responsible for their own death. *S. mutans* then also dephosphorylates the xylitol-5-phosphate, and this dephosphorylated sugar molecule is ejected from the cell. This ejection occurs with an energy expenditure without energy contribution from the metabolism of xylitol.

Consequently, xylitol reduces or inhibits the growth of mutans streptococci through starving the bacteria (El-Marakby et al. 2017). The antimicrobial effects of xylitol, xylitol-probiotic, and fluoride toothpaste on *S. mutans* plaque were observed in 13–15-year-old children (Arat Maden et al. 2017). Thus, regular use of toothpaste containing a combination of fluorides and other appropriate components can prevent some oral diseases. There has been a study about the individuals having xylitol regularly. Their plaque sample examination showed that there is a substantial reduction in plaque adhesiveness, and there is a significant reduction of extracellular polysaccharides produced by *S. mutans* to develop biofilms (Burt 2006). The anti-caries potential of xylitol is mainly related to its effect on dental plaque and cariogenic bacteria (El-Marakby et al. 2017).

As a consequence of the regular use of xylitol, numerous effects appear. The first effect is very selective in *S. mutans*, which become xylitol-resistant. A second effect is an increase in the concentration of ammonia and many basic amino acids when the dental plaque is exposed to xylitol. Consequently, plaque acids become neutralized. So that is how xylitol showed its antiplaque properties (Maguire and Rugg-Gunn 2003).

Candidiasis is the most common fungal infection among the human population caused mostly by *Candida albicans* (Talattof et al. 2018). As sugar intake promotes the *C. albicans* growth, xylitol can decrease the risk of candidiasis in the oral mucosa; the minimum inhibition concentration of xylitol for *C. albicans* was found to be 20×10^4 µg/mL (Talattof et al. 2018).

Xylitol is also responsible for increasing mineralization on dental tissues. Several clinical and laboratory studies reported the remineralization of dental caries by consuming xylitol gums regularly. It increases the salivary flow in case of consuming as chewing gum or large xylitol tablets. Chewing gums with xylitol demonstrated good antibiofilm and remineralized potential on damaged by caries teeth areas (Gargouri et al. 2018). Xylitol- and sorbitol-containing chewing gums are also responsible for reducing gingival

inflammation (Shyama et al. 2006). Such gums enhance the potential of remineralization. Gums containing xylitol showed some superior effects regarding remineralization and reduction of plaque formation (Steinberg et al. 1992). Although remineralization has been noticed on all such experiments carried with non-sugar sweeteners substances, the remineralization process occurs due to enhanced saliva flow that is rich in phosphate and calcium.

The efficiency of xylitol chewing gums enriched with propolis was established *in vitro* by the remineralizing effect of softly demineralized dentin (Gargouri et al. 2020).

Effect of xylitol on oral, nasopharyngeal, and gut microbiota

Biofilm control is crucial for preventing of such oral infections as caries and periodontal diseases. Xylitol can provoke decreasing oral biofilm formation by the inhibition of bacterial β -glucosidase in human saliva, which is essential for biofilm formation in the oral cavity (Teixeira Essenfelder et al. 2019). *S. mutans* can incorporate xylitol, which then can inhibit the process of glycolysis of these microorganisms by the use of a fructose phosphotransferase system. Inside the bacteria cell, the molecule of this sugar is phosphorylated, resulting in the production of xylitol-5-phosphate, which cannot be metabolized (Teixeira Essenfelder et al. 2019).

A 30-day supplementation with xylitol and maltitol chewing gums led to a significant decrease in the presence of oral pathogenic microorganisms *S. mutans*, *S. sobrinus*, and *Actinomyces viscosus*; that effect was directly linked to an increase in plaque pH by these polyols which cannot be fermented by oral bacteria and can similarly improve general oral health (Thabuis et al. 2013). The chewing gum with xylitol decreased the *S. mutans*, and *S. sobrinus* counts significantly comparatively to the sorbitol gum group (Bahador et al. 2012).

The everyday use of xylitol and isothiazolinone demonstrated the pronounced antimicrobial effect against *Pseudomonas aeruginosa* ATCC 9027 and *Staphylococcus aureus* ATCC 6538 biofilms in a dose-dependent manner and may be applied to reduce the undesirable biofilms *in vitro* (Zhou et al. 2019). High levels of antimicrobial activity against among others *S. aureus* and *C. albicans* were demonstrated by the essential oils of plants containing such predominant components as thymol, carvacrol, and other aromatic terpenoids (Kozłowska et al. 2015; Marchese et al. 2016). It would be interesting to investigate the combined cumulative effect of xylitol and essential oil components, such as menthol, in chewing gum (Santos et al. 2014). It was revealed the positive effect of chewing gums containing probiotics (*Lactobacillus reuteri*) and xylitol on *S. mutans* counts and gingival scores (Kaur et al. 2018).

Factors such as oral hygiene and fermentable carbohydrate exposure frequency should also be controlled during the studies of the anti-cariogenic effect of xylitol (Janket et al. 2019). However, the contrast between xylitol and other sweeteners, such as sorbitol or sucralose, was not significantly different (Janket et al. 2019). Xylitol can inhibit the glucosyltransferase, which blocks the utilization of glucose by *S. mutans* and adhesion this pathogenic microorganism to the tooth surface (Janket et al. 2019). The effects of xylitol, sucrose, and their combination have been studied *in vitro* on *S. mutans* and *S. sobrinus* (Salli et al. 2016). While the addition of 1% sucrose significantly increased the degree of bacterial strains, the study performed with increasing the concentration (2–5%) of xylitol showed a decrease in bacterial counts also in the presence of sucrose.

Xylitol-containing chewing gum inhibited the increase in total salivary bacteria in the case of short-term research of 70 healthy adult men (Takeuchi et al. 2018).

Xylitol reduces the *S. mutans* counts and *lactobacilli* in saliva. There are very few other species that are beneficial in the oral microbiome (Badet et al. 2008). According to a study, by consuming xylitol, the lactobacilli were unidentified in oral microbiome while the *S. mutans* were observed in a minimal amount in the oral cavity. Therefore, this proved that xylitol has excellent effects on oral microbiome communities as it has anti-caries and antiplaque potential. It inhibits the growth of all those bacterial colonies which have a role in chronic periodontitis (Rafeek et al. 2019).

The recent research outcomes of a randomized clinical trial among children concerning the comparison of effectiveness for mouthwashes comprising green tea with xylitol vs. green tea alone (two times daily during 14 days). The salivary *S. mutans* and *Lactobacillus* colony count revealed a more significant reduction with the administration of green tea with xylitol mouthwash (Hajiahmadi et al. 2019).

Xylitol has also affect on the growth of nasopharyngeal microbiota. This could be important when noticing the respiratory pathologies caused by nasopharyngeal bacteria, such as *S. pneumoniae*. These bacteria respond to xylitol in the same way as *S. mutans* respond to xylitol in the oral cavity. To evaluate this statement, a study has been conducted in which they grew some nasopharyngeal bacteria in media that was consisted of xylitol sugar. The results of this study showed that xylitol could significantly reduce the growth levels of *S. pneumoniae* and *S. mitis* during the growth phase (Sakallioğlu et al. 2014). This effect enhanced with an increase in the concentration of xylitol in growth media. There also was observed the slight inhibition of growth of β -hemolytic streptococci only by 5% xylitol concentration. However, no inhibition in the case of *Haemophilus influenzae* and *Moraxella catarrhalis* has been demonstrated (Kontiohari et al. 1995). Xylitol (5% and 10%) significantly decreased the biomass of such pathogenic bacteria like *Staphylococcus*

aureus and *S. epidermidis*, and *Pseudomonas aeruginosa* involved in the formation of biofilms in case of chronic rhinosinusitis (Jain et al. 2016).

In an experimental model of rhinitis medicamentosa in rats, the application of xylitol solution during 15 days demonstrated the effectiveness for nasal mucosas, evaluated histopathologically, as of a reference drug mometasone (Cam et al. 2019).

Gut microbiota is responsible for forming many bioactive compounds from dietary substances which can maintain the whole-body metabolism (Ruiz-Ojeda et al. 2019, 2020). For example, better glucose metabolism can be induced by the help of dietary fibers that are linked with the greater quantity of *Prevotella* (Kovatcheva-Datchary et al. 2015). Same as some food additives and derivatives can produce some effects on the metabolism of the host after association with microbiota present in the gut. The study has been conducted with xylitol on mice model to check xylitol's effect on gut microbiota. They supply the xylitol 40 mg/kg bodyweight of mice in the diet. Xylitol induced substantial changes in the gut microbiota of mice (Tamura et al. 2013). They observed the changes in genera *Barnesiella*, which was significantly reduced in gut microbiota while the *Feacalibaculum* was significantly increased. *Barnesiella* and *Feacalibaculum* both have been spotted in the mice gut microbiota as well as human gut microbiota. The study concluded that the xylitol could induce changes in gut microbiota or bacterial colonies present in the gut, but it does not affect lipid metabolism (Uebanso et al. 2017).

Xylitol in the prevention of dental caries

Dental caries is one of the most common infections that affect the oral cavity associated with high costs for families. Research shows that it is still a significant health concern in most developing countries, affecting 60–90% of school-going children and many adults (Riggs et al. 2019). The broadening of the available spectrum of prophylactic substances (specifically xylitol) in an accessible form is hugely relevant at present. It was suggested that the positive effect of polyols in caries prevention might depend on the number of OH-groups in their molecule: erythritol \geq xylitol $>$ sorbitol (Mäkinen 2010). According to some authors, the prevalence of permanent tooth decay in children aged 6–16 exceeds 70 percent with an intensity of 2,86 index SIC – $5,20 \pm 0,26$. The highest increase in permanent tooth decay is observed in the age groups of 6–7 years (by 2.91-fold) and 11–12 years (by 1.37-fold) (Chukhray 2018). Dental caries is a multifactorial disease, which depends on the use of fermentable carbohydrates by acidogenic and aciduric microorganisms forming the dental biofilms. It has been established that natural protective mechanisms of the oral cavity of children with high-

intensity decay are lowered. In particular, the level of IgA secretion is approximately 2.4 g/L, lysozyme (29.44 L/g), *Lactobacillus* concentration (41.02 percent), all who are below normal levels (Cherepyk 2017). The consumption of sucrose in large quantities at the age of 3 years can enhance the risk of increased colonies of mutans streptococci and increase the risk for dental caries. Several studies conclude that refined sugar like sucrose is a significant cause of developing caries in children and even in adults. *S. mutans* is the main producer of acids during the metabolism of sugars, mainly sucrose. The spreading of these acids at the interface between dental biofilms and enamel promotes a fast decrease of pH and a resultant stimulus to the demineralization process. Acids further diffuse into the enamel and dentin layer and dissolve the minerals there.

S. mutans forms biofilms known as dental plaque on the tooth surfaces, and it can also be responsible for the infective endocarditis (Lemos et al. 2019). *S. mutans* was firstly isolated from carious lesions by J. Clarke in 1924. However, it was considered an important etiologic agent in caries by the middle of the 1960s only (Lemos et al. 2019). Xylitol sugar has the potential to reduce the growth of *S. mutans* and, consequently, the production of dental caries.

Xylitol disturbs the bacterial metabolism and especially their energy production. It triggers the process of energy consumption cycle, and bacterial cell death occurs due to this phenomenon. Xylitol can reduce plaque formation. It decreases the demineralization of enamel and dentin (Ly et al. 2006). It also has an inhibitory effect on bacterial adherence and the production of biofilms in the oral cavity (Ammons et al. 2011). Very first dental caries prevention by xylitol was studied and researched in Finland in the 1970s, by using different animal models. The first xylitol related study in humans is referred to as “Turku Sugar Studies,” and it was demonstrated a reduction of dental plaque through xylitol consumption. In Finland, for the first time, xylitol was introduced into chewing gum to cure dental caries by improving the health of the oral cavity.

The use of xylitol as a caries-preventing agent has been endorsed by many public regulatory affairs (Kitchens 2005; Mäkinen 2011). Xylitol-containing food and oral hygiene adjuvants have been shown to reduce the growth of dental plaque by decreasing the growth of caries-associated bacteria (Mäkinen 2011). Application of xylitol wipes might be a useful complement in reducing the cariogenic bacteria, especially *S. mutans*, and thus could be considered as an adjunct oral hygiene tool for caries prevention in children (Kayalvizhi et al. 2018).

The most efficient vehicle or mode of delivery of xylitol is, however, considered the chewing gum. The caries preventive effect of 1 year using of xylitol-containing chewing gum in an adult population showed that it had a reduction of caries disease at teeth level of 23% (Cocco et al. 2017). After using a

chewing gum containing a high dose of xylitol during the half-year period, it was proved to be efficacious in caries prevention in high-risk children (Campus et al. 2013).

Other professional agents that can prevent dental caries are consisted of some toothpaste and mouthwash that contain xylitol components and fluoride as well. Toothpaste containing xylitol and fluoride may be more effective than toothpaste with fluoride only (Duane 2015). As dental caries are mostly experienced in young school-going children, so to treat dental caries in children xylitol-containing candies, gummy bears, gums, and lozenges are better and fascinating options.

Xylitol is better-tolerated polyol comparatively to d-glucitol or d-mannitol, and its highest safe doses typically range 20–70 g per day; the quantity of xylitol recommended for caries inhibition is about 10 g/day for adults; children should intake smaller quantities (Mäkinen 2016). Researchers revealed the need for well-designed clinical studies with adequate control groups after studying the prevention of caries after consuming xylitol-containing candies and lozenges (Antonio et al. 2011). A systematic review (Marghalani et al. 2017) examined the effectiveness of xylitol in reducing dental caries in children compared to placebo, no treatment, or other preventive strategies revealed a small effect size in randomized trials and a very low level of evidence that makes its preventive effect uncertain. Anyway, from all the studies and researches until now, it is concluded that the xylitol products are very beneficial and helpful to treat dental caries, to reduce the growth of mutans streptococci, to eliminate the plaque, and ultimately to improve the health of oral cavity (Janakiram et al. 2017).

Xylitol and insulin

Research showed that the administration of xylitol could induce a rapid release of insulin in plasma. The study has been done in dogs (0.4 g/kg xylitol, intravenously). The prompt rise in plasma insulin was observed as a result, though the procedure or mechanism behind the stimulation of the release of insulin is still ambiguous. Xylitol enters the pentose phosphate pathway through D-xylulose-5-phosphate. Therefore, it shares the mutual metabolic pathway with glucose. The researchers further also checked the effects of insulin by changing the doses of xylitol and its administration by the oral route.

The results showed that oral administration of xylitol incites a more substantial effect on the secretion of insulin than the intravenous route (Kuzuya et al. 1969). In another research (Kuzuya et al. 1971) performed on humans and some mammals, both xylitol and glucose were administered intravenously in equal quantity. The significant differences in insulin responses among different species have been observed. In man, glucose produced a definite increase in plasma insulin as compared to xylitol. While in cows and goats, glucose and xylitol

both increased the plasma insulin markedly to an approximate extent. In horses, the insulin release was instead a bit poor and sluggish after administration of glucose and xylitol. Rabbits also showed the same extent of insulin release in response to glucose and xylitol. In rats, glucose showed higher insulin releases in plasma as compared to xylitol (Kishore et al. 2012). However, in dogs, the results were a bit different from all other mammals: xylitol showed a definite higher insulin peak as compared to glucose. The reason behind this species variation could not be explained (Kuzuya et al. 1971).

The utilization of carbohydrates has been studied and compared the results with glucose utilization during the endogenous insulin secretion inhibition. The experiment has been carried out in 28 healthy volunteers, and they were divided into further five groups (sorbitol, fructose, glucose, xylitol, and saline). The results concluded that sorbitol, fructose, and xylitol are oxidized more as compare to glucose during the suppression of insulin secretion without any substantial increase in glycemic index (de Kalbermatten et al. 1980). Therefore, low glycemic and insulinemic influences of xylitol are beneficial (Janket et al. 2019).

Xylitol and inflammation

Xylitol has an inhibitory effect on inflammatory reactions. The important and vital process that is involved in inflammation is angiogenesis, which occurs during tumor formation and metastasis. Many research studies have been carried out on natural compounds that are responsible for reducing or inhibiting inflammatory responses. Xylitol is one of the natural compounds that can inhibit inflammatory responses. Xylitol can inhibit the expression of inflammatory cytokines that are triggered by lipopolysaccharides (LPS). It has been discovered that inflammation and angiogenesis share a mutual or common signaling pathway. Xylitol is responsible for inhibiting the invasion, migration, and tube formation of human umbilical vein endothelial cells (HUVECs). *In vivo* studies on rats showed that xylitol also inhibited angiogenesis in a mouse Matrigel plug assay. By treating the rats with xylitol, there was a decreased mRNA expression of vascular endothelial growth factor (VEGF), matrix-metalloproteinase-2 (MMP-2), and basic fibroblast growth factor (BFGF). This anti-inflammatory and anti-angiogenic potential of xylitol can be exerted via inhibition of the NF- κ B pathway and Akt activation pathway. These results showed that xylitol has a beneficial role against inflammation and angiogenesis (Yi and Kim 2013).

Porphyromonas gingivalis is the most identified bacteria responsible for causing dental pathologies. The LPS secreted by *P. gingivalis* is the main factor in the development of periodontitis. Inflammatory cytokines, induced by LPS, play a significant role in the destruction of gingival tissues. Xylitol

can inhibit the production of inflammatory cytokines and reduce the secretion of LPS by *P. gingivalis*. The pretreatment with xylitol can suppress the expression of cytokines that cause inflammation (Han et al. 2005). Xylitol also inhibits the TNF- α and IL-1 β gene expression and, ultimately, protein synthesis. Studies showed that LPS-induced activation of the NF- κ B pathway could also be inhibited by xylitol that is involved in inflammatory responses. Xylitol also displayed an inhibitory effect on the growth of *P. gingivalis*.

All these findings collectively indicate that xylitol can have an excellent clinical effect not only for dental pathologies but also to inhibit LPS-trigger inflammatory cytokine expression (Han et al. 2005). Preparations containing xylitol are known to be used effectively in the comprehensive treatment of endogenous intoxication in suppurative diseases in maxillofacial surgery (Medvid 2011).

Xylitol in the prevention of obesity and metabolic syndrome

Obesity has become a significant health concern in the whole world by being the principal risk factor for several diseases, such as cardiovascular diseases, metabolic syndrome, stroke, atherosclerosis, and even cancer. Obesity is caused by several environmental factors involving diet and nutrient composition. A diet containing high-fat components is responsible for causing metabolic syndromes, among others, obesity, dyslipidemia, and hyperglycemia. Xylitol is used in medications as low-calorie sugar. While taking xylitol, a small increase in blood glucose level can be observed, a rapid release of insulin occurs, and glucose level can overcome in diabetic and healthy individuals due to the release of insulin. Therefore, it is recommended that diabetic patients can uptake xylitol as a sugar substitute and as an energy source in place of other carbohydrates dietary products. In the liver, xylitol is first phosphorylated and then converted into xylulose-5-phosphate through a metabolic pentose phosphate pathway. This xylitol metabolite, xylulose-5-phosphate, is responsible for activating nuclear transport activities and DNA-binding activities of carbohydrates response element of a binding protein (ChREBP) via the activation of another enzyme known as protein phosphatase 2A (PP2A). ChREBP is a commonly known transcription factor that triggers or stimulates the genes associated with a lipogenic enzyme such as acetyl coenzyme A carboxylase (ACC) and fatty acid synthase (FAS).

Consequently, this increased lipogenesis in the liver results in obesity or stenosis. Therefore, in that case, excessive consumption of xylitol combined with a high-fat diet can activate lipogenesis genes, and it induces obesity in people and ultimately leading to metabolic syndrome. However, still, the long-term effects of xylitol on carbohydrate and lipid metabolism cannot be fully understood—more studies and research

are required to reveal its mechanism. Because xylitol has been used in appropriate doses for some beneficial effects such as it is useful to reduce the postprandial hyperglycemia and, ultimately, diabetes, obesity, and metabolic syndrome (Amo et al. 2011; Pearlman et al. 2017). It was observed that gastric emptying was significantly slower with xylitol intake; thus, it can prevent the hunger sensation and food intake (Janket et al. 2019). The impact of xylitol sugar is very low on the blood glycemic index as compared to other sugars, so it indicates that xylitol can be a beneficial substitute for natural sugars that, among others, can develop cardiac diseases and diabetes (Soderling 2009a).

Xylitol and prevention from other diseases

Except for the diseases already discussed, such as oral cavity diseases, obesity, and metabolic syndrome, some researches showed that xylitol could prevent some other diseases, too, for example, some respiratory diseases, among others pneumonia and middle ear infection (Lee and Park 2014).

The bacterium responsible for pneumonia is *S. pneumonia*. This bacterium is becoming resistant to beta-lactam antibiotics and penicillin day by day. Pneumonia affects mostly young children. The currently developed antibiotics and vaccines do not have the potential to overcome this disease because its pathogen has developed resistance against them. Alternative preventive treatment or cure is necessary to overcome this disease. Studies showed that the five-carbon sugar alcohol xylitol could cure this disease. The derivative of xylitol, known as xylitol phosphate that is produced by *S. pneumonia*, has the potential to inhibit the growth of them. Consequently, the growth of these resistant bacteria can be stopped by using xylitol (Palchaudhuri et al. 2015).

The most common middle ear infection in the USA among young children is known as acute otitis media. In this infection, bacteria create a sticky fluid in the middle ear right behind the eardrum. Severe pain can be experienced in that situation. Fever and ear secretion can also occur in this disease among children, although no severe complications are noted in acute otitis media (Uhari et al. 2000). The treatment of this disease is a bit costly, and the pathogen can develop antibiotic resistance. Surgery is also an inappropriate and costly option to manage middle ear infection in children. Therefore, xylitol is considered an appropriate option.

Xylitol can be found naturally in many fruits, among other strawberries, raspberries, plums, and rowan berries. Artificially, it is present in among others gums, gummy bears, candies, and pills. So, children can easily and happily consume xylitol to treat their oral cavity infections and middle ear infections.

The xylitol syrup is also available as a medicine for young children. Xylitol can be administered in gums, syrups, and

lozenges, and these products can eliminate acute otitis media among young children with no severe respiratory infections and side effects (Azarpazhooh et al. 2016). Another study reported that young individuals have a very low incidence of acute otitis media if they are consuming xylitol oral solution in a dose of 2–5 g daily. Despite all these studies, xylitol is not being used at the global level consistently (Vernacchio et al. 2014).

Xylitol has been reported to improve barrier function and inhibit the growth of potential skin pathogenic microorganisms. As a non-digestible organic compound, xylitol enters the colon where it is fermented by the colonic microbiota such as *Anaerostipes* species to produce butyrate (Salli et al. 2019). Sugar alcohols have some prebiotic properties (Mäkinen 2016). Xylitol can improve bone mineral density and modulate the immune system, which, together with its antimicrobial effect, contributes to a reduced respiratory tract infection risk. Safety and efficacy of aerosolized xylitol (15%, 5 ml) used twice a day for 2 weeks in patients hospitalized with pulmonary cystic fibrosis, has been proven (Singh et al. 2020). As a low caloric sweetener, xylitol may contribute to weight management due to its ability to increase satiety (Salli et al. 2019).

Safety and eventual adverse effects of xylitol in adults

Despite xylitol sugar possess many positive effects, its consumption is usually associated with irritable bowel syndrome (IBS) in adults. Abnormal flatulence also has been observed by consuming xylitol (Mäkinen 2016). Therefore, xylitol consumption can also affect human health negatively. It is thought to be a burden in economic terms of health care costs. Although it has been found that sugar alcohols can be useful in the treatment of severe or chronic constipation. For instance, D-glucitol has been utilized in many commercial preparations (Mäkinen 2016). No previous study has primarily focused on osmotic diarrhea linked with the overuse of sugar alcohol substances. Fewer studies are present in which difficulties related to sugar alcohols have been discussed. It is necessary to focus that osmotic diarrhea, which can be observed by consuming slowly absorbed polyols and carbohydrates, is not considered as a disease but somewhat an unpretentious physiochemical response of the gastrointestinal tract in the presence of polyols in the gut (Mäkinen 2016). When an individual consumes carbohydrate or polyol solutions, in response, water draw from the body into the lumen of the gut, this phenomenon causes osmotic diarrhea (Grillaud et al. 2005). Osmotic diarrhea can also be seen in some diseases like pancreatic diseases (Oku and Nakamura 2007). It is also necessary to focus that severe osmotic effects can also be seen when people eat a large number of grains, cereals, vegetables, and certain fruits. In healthy people, consuming the extra-

large amount of substances, for example, vitamin C, salts containing magnesium, and some antibiotics can cause acute cases of osmotic diarrhea and bowel diseases (Mäkinen 2016).

Xylitol can be tolerated well, even in large quantities. Though susceptible persons may require adapting to digest higher intake of xylitol, by adapting to tolerate the higher amounts, xylitol can increase the activity levels of enzyme polyol dehydrogenase. This enzyme is beneficial to tolerate and absorb the xylitol in the body. Xylitol sugar is slowly absorbed and metabolized by the body. It can induce minimal alterations in insulin release.

Xylitol in the body can bind with calcium ion and some other cations. This phenomenon contributes to the remineralization of the enamel of teeth. The complexes of xylitol-calcium facilitate the absorption of calcium throughout the gut wall. This indicates that xylitol can be beneficial in preventing osteoporosis (Vasilescu et al. 2011).

However, several studies reported increased the risk factors for urolithiasis in the case of long-term xylitol intake, which could be attributable to osmotic diarrhea that may cause metabolic acidosis (Janket et al. 2019).

Safety and eventual adverse effects of xylitol in children

Xylitol is a useful option to prevent dental caries and acute otitis media (AOM) in children. A study (Vernacchio et al. 2007) has described its efficiency when xylitol is given to the children five times a day to prevent AOM. Despite this advantage, xylitol is not considered as the best option worldwide to cure acute otitis media in young children because there are two significant reasons. One of them is the impractical schedule of dosing five times, which is difficult to follow. The other actual reason is that xylitol can produce some gastrointestinal side effects while treating acute otitis media in young children and especially in infants (Mäkinen 2016). The study (Vernacchio et al. 2007) was done to check the xylitol side effects in children. The researchers did the dosing of xylitol five times a day in children 6–36 months of age. The subjects tolerate the xylitol solution very well. Only three subjects out of 30 developed the gastrointestinal side effects. The subjects who developed side effects were usually infants. The side effects observed among children were comprised of the development of excessive gas, loose stool, and diarrhea. The rates of diarrhea were more significant as compared to other side effects in children reported by parents (Vernacchio et al. 2007). Therefore, this study concluded that infants have more chances to develop gastrointestinal side effects while consuming xylitol solution daily. However, still, further research is required to approve these observations.

Overuse of xylitol is responsible for the accumulation of xylitol in the gut lumen that will lead to retention of water and

consequently cause diarrhea in young children. Also, other studies showed that an excessive quantity of xylitol would lead to the development of gas and bloating (Storey et al. 2007). Xylitol solution, which cannot be absorbed, is eliminated after metabolizing into carbon dioxide. In 1985 European Union's Scientific Committee on Food published a report in which it has been stated that taking or consuming 50 g of xylitol solution can cause diarrhea. The union also confirmed that if the sweeteners containing xylitol consumed too much, it could induce some laxative side effects in children.

Conclusions

Xylitol is belonging to a carbohydrate class of biomolecules, known as a sugar alcohol, and used as a sweetener. It can be found naturally in a very low quantity. Commercially, xylitol is produced through the recovery of waste constituents from plants manufacturing chemically or by biotechnological processes.

Chewing gums occupy a leading position among the different vehicles used for delivering the xylitol into the human body. This biomolecule has excellent potential to prevent dental caries, to support the remineralization of teeth enamel, and acts as antiplaque and anti-gingivitis to improve the health of the oral cavity among young children. Xylitol also has good prospects in the area of inhibiting the growth of antibiotic-resistant bacteria. This sugar alcohol has excellent anti-inflammatory potential and can cure chronic inflammatory diseases efficiently; it can play a positive role in the treatment, among others, middle ear infections and respiratory diseases such as pneumonia. Xylitol is also responsible for releases insulin in blood plasma in mammals; it can also reduce obesity and some other metabolic syndromes. When taken in excessive amounts, xylitol can produce some side effects, such as diarrhea and loose stools. However, several studies have demonstrated its many beneficial effects on human health. Xylitol can prevent and treat various diseases, which not can be treated with antibiotics or surgery. Still, advanced research is needed to explore more the beneficial effects of xylitol.

Authors' contributions AGB, AG, and MA contributed to the acquisition of data and writing the first version of the manuscript. MS, RL, MP, IPT, SA, and YS supported the conception, and revised the manuscript for essential intellectual content. GB contributed to the acquisition of data, drafting and revising the manuscript, and coordinated the project. All authors approved the final version of the manuscript.

Compliance with ethical standards

Conflict of interest The authors declare that they have no competing interests.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Abd Rahman NH, Md Jahim J, Abdul Munaim MS, R AR, Fuzi SFZ, Md Illias R (2020) Immobilization of recombinant *Escherichia coli* on multi-walled carbon nanotubes for xylitol production. *Enzym Microb Technol* 135:109495. <https://doi.org/10.1016/j.enzmictec.2019.109495>
- Alves FRF, Neves MAS, Silva MG, Rocas IN, Siqueira JF Jr (2013) Antibiofilm and antibacterial activities of farnesol and xylitol as potential endodontic irrigants. *Braz Dent J* 24:224–229
- Ammons MCB, Ward LS, James GA (2011) Anti-biofilm efficacy of a lactoferrin/xylitol wound hydrogel used in combination with silver wound dressings. *Int Wound J* 8(3):268–273. <https://doi.org/10.1111/j.1742-481X.2011.00781.x>
- Amo K, Arai H, Uebanso T, Fukaya M, Koganei M, Sasaki H, Yamamoto H, Taketani Y, Takeda E (2011) Effects of xylitol on metabolic parameters and visceral fat accumulation. *J Clin Biochem Nutr* 49(1):1–7. <https://doi.org/10.3164/jcbs.10-111>
- Antonio AG, Pierro VS, Maia LC (2011) Caries preventive effects of xylitol-based candies and lozenges: a systematic review. *J Public Health Dent* 71(2):117–124. <https://doi.org/10.1111/j.1752-7325.2010.00208.x>
- Arat Maden E, Altun C, Acikel C (2017) The Efficacy of xylitol, xylitol-probiotic and fluoride dentifrices in plaque reduction and gingival inflammation in children: a randomised controlled clinical trial. *Oral Health Prev Dent* 15(2):117–121. <https://doi.org/10.3290/j.ohpd.a37973>
- Azarapazhooh A, Lawrence HP, Shah PS (2016) Xylitol for preventing acute otitis media in children up to 12 years of age. *Cochrane Database Syst Rev* 8:CD007095. <https://doi.org/10.1002/14651858.CD007095.pub3>
- Badet C, Furiga A, Thebaud N (2008) Effect of xylitol on an in vitro model of oral biofilm. *Oral Health Prev Dent* 6(4):337–341
- Bahador A, Lesan S, Kashi N (2012) Effect of xylitol on cariogenic and beneficial oral streptococci: a randomized, double-blind crossover trial. *Iran J Microbiol* 4(2):75–81
- Bilal M, Xu S, Iqbal HMN, Cheng H (2020) Yarrowia lipolytica as an emerging biotechnological chassis for functional sugars biosynthesis. *Crit Rev Food Sci Nutr*. <https://doi.org/10.1080/10408398.2020.1739000>
- Burt BA (2006) The use of sorbitol- and xylitol-sweetened chewing gum in caries control. *J Am Dent Assoc* 137(2):190–196. <https://doi.org/10.14219/jada.archive.2006.0144>
- Cam B, Sari M, Mide A, Gergin O (2019) Xylitol treats nasal mucosa in rhinitis medicamentosa: an experimental rat model study. *Eur Arch Otorhinolaryngol* 276(11):3123–3130. <https://doi.org/10.1007/s00405-019-05605-3>
- Campus G, Cagetti MG, Sale S, Petruzzi M, Solinas G, Strohmenger L, Lingstrom P (2013) Six months of high-dose xylitol in high-risk caries subjects—a 2-year randomised, clinical trial. *Clin Oral Investig* 17(3):785–791. <https://doi.org/10.1007/s00784-012-0774-5>
- Carneiro C, de Paula ESFC, Almeida JRM (2019) Xylitol production: identification and comparison of new producing yeasts. *Microorganisms* 7(11). <https://doi.org/10.3390/microorganisms7110484>
- Cherepyk OM (2017) Substantiation of early prevention of caries of temporary teeth in children of Precarpathian region [in Ukrainian]. PhD (Dentistry) dissertation, Ivano-Frankivsk National Medical University

- Chukhray NL (2018) Substantiation of dental caries prevention in children based on associated risk factors and their influence on the formation of enamel resistance [in Ukrainian]. DScD manuscript, University of Lviv
- Chukwuma CI, Islam MS (2017) Xylitol: one name, numerous benefits. In: Merillon J-M, Ramawat KG (eds) Sweeteners: Pharmacology, Biotechnology, and Applications. Springer International Publishing, Cham, pp 1–27
- Cocco F, Carta G, Cagetti MG, Strohmenger L, Lingström P, Campus G (2017) The caries preventive effect of 1-year use of low-dose xylitol chewing gum. A randomized placebo-controlled clinical trial in high-caries-risk adults. Clin Oral Investig 21(9):2733–2740. <https://doi.org/10.1007/s00784-017-2075-5>
- de Kalbarmatten N, Ravussin E, Maeder E, Geser C, Jequier E, Felber JP (1980) Comparison of glucose, fructose, sorbitol, and xylitol utilization in humans during insulin suppression. Metabolism 29(1):62–67. [https://doi.org/10.1016/0026-0495\(80\)90099-2](https://doi.org/10.1016/0026-0495(80)90099-2)
- Duane B (2015) Xylitol and caries prevention. Evid Based Dent 16(2):37–38. <https://doi.org/10.1038/sj.ebd.6401088>
- El-Marakby AM, Al-Sabri FA, Mohamed SG, Labib LM (2017) Anticariogenic effect of five-carbon sugar: xylitol. JDOH 3(6):081
- Felipe Hernandez-Perez A, de Arruda PV, Sene L, da Silva SS, Kumar Chandel A, de Almeida Felipe MDG (2019) Xylitol bioproduction: state-of-the-art, industrial paradigm shift, and opportunities for integrated biorefineries. Crit Rev Biotechnol 39(7):924–943. <https://doi.org/10.1080/07388551.2019.1640658>
- Fierascu RC, Fierascu I, Ortan A, Georgiev MI, Sieniawska E (2020) Innovative approaches for recovery of phytoconstituents from medicinal/aromatic plants and biotechnological production. Molecules 25(2). <https://doi.org/10.3390/molecules25020309>
- Ganter J, Hellwig E, Doerken S, Al-Ahmad A (2020) In vitro evaluation of the cariogenic potential of rebaudioside A compared to sucrose and xylitol. Clin Oral Investig 24(1):113–122. <https://doi.org/10.1007/s00784-019-02908-x>
- Gargouri W, Zmantar T, Kammoun R, Kechaou N, Ghoul-Mazgar S (2018) Coupling xylitol with remineralizing agents improves tooth protection against demineralization but reduces antibiofilm effect. Microb Pathog 123:177–182. <https://doi.org/10.1016/j.micpath.2018.06.038>
- Gargouri W, Kammoun R, Elleuche M, Tlili M, Kechaou N, Ghoul-Mazgar S (2020) Effect of xylitol chewing gum enriched with propolis on dentin remineralization in vitro. Arch Oral Biol 112:104684. <https://doi.org/10.1016/j.archoralbio.2020.104684>
- Granstrom TB, Izumori K, Leisola M (2007) A rare sugar xylitol. Part I: the biochemistry and biosynthesis of xylitol. Appl Microbiol Biotechnol 74(2):277–281. <https://doi.org/10.1007/s00253-006-0761-3>
- Grillaud M, Bandon D, Nancy J, Delbos Y, Vaysse F (2005) The polyols in pediatric dentistry: advantages of xylitol. Arch Pediatr 12(7):1180–1186. <https://doi.org/10.1016/j.arcped.2005.01.032>
- Hajiahmadi M, Yegdaneh A, Homayoni A, Parishani H, Moshkelgosha H, Salari-Moghaddam R (2019) Comparative Evaluation of Efficacy of “Green Tea” and “Green Tea with Xylitol” mouthwashes on the salivary *Streptococcus mutans* and *Lactobacillus* colony count in children: a randomized clinical trial. J Contemp Dent Pract 20(10):1190–1194
- Han SJ, Jeong SY, Nam YJ, Yang KH, Lim HS, Chung J (2005) Xylitol inhibits inflammatory cytokine expression induced by lipopolysaccharide from *Porphyromonas gingivalis*. Clin Diagn Lab Immunol 12(11):1285–1291. <https://doi.org/10.1128/CDLI.12.11.1285-1291.2005>
- Honkala S, Runnel R, Saag M, Olak J, Nommela R, Russak S, Makinen PL, Vahlberg T, Falony G, Makinen K, Honkala E (2014) Effect of erythritol and xylitol on dental caries prevention in children. Caries Res 48(5):482–490. <https://doi.org/10.1159/000358399>
- Jain H, Mulay S (2014) A review on different modes and methods for yielding a pentose sugar: xylitol. Int J Food Sci Nutr 65(2):135–143. <https://doi.org/10.3109/09637486.2013.845651>
- Jain R, Lee T, Hardcastle T, Biswas K, Radcliff F, Douglas R (2016) The in vitro effect of xylitol on chronic rhinosinusitis biofilms. Rhinology 54(4):323–328. <https://doi.org/10.4193/Rhin15.380>
- Janakiram C, Deepan Kumar CV, Joseph J (2017) Xylitol in preventing dental caries: a systematic review and meta-analyses. J Nat Sci Biol Med 8(1):16–21. <https://doi.org/10.4103/0976-9668.198344>
- Janket SJ, Benwait J, Isaac P, Ackerson LK, Meurman JH (2019) Oral and systemic effects of xylitol consumption. Caries Res 53(5):491–501. <https://doi.org/10.1159/000499194>
- Jones AH (2000) Xylitol delivery. United States Patent.
- Kaur K, Nekkanti S, Madiyal M, Choudhary P (2018) Effect of chewing gums containing probiotics and xylitol on oral health in children: a randomized controlled trial. J Int Oral Health 10(5):237–243. https://doi.org/10.4103/jioh.jioh_170_18
- Kayalvizhi G, Nivedha D, Sajeev R, Prathima GS, Suganya M, Ramesh V (2018) Evaluating the efficacy of xylitol wipes on cariogenic bacteria in 19- to 35-month-old children: a double-blind randomized controlled trial. Int J Clin Pediatr Dent 11(1):13–17. <https://doi.org/10.5005/jp-journals-10005-1476>
- Kishore P, Kehlenbrink S, Hu M, Zhang K, Gutierrez-Juarez R, Koppaka S, El-Maghrabi MR, Hawkins M (2012) Xylitol prevents NEFA-induced insulin resistance in rats. Diabetologia 55(6):1808–1812. <https://doi.org/10.1007/s00125-012-2527-z>
- Kitchens DH (2005) Xylitol in the prevention of oral diseases. Spec Care Dentist 25(3):140–144. <https://doi.org/10.1111/j.1754-4505.2005.tb01424.x>
- Kontikari T, Uhari M, Koskela M (1995) Effect of xylitol on growth of nasopharyngeal bacteria in vitro. Antimicrob Agents Chemother 39(8):1820–1823. <https://doi.org/10.1128/aac.39.8.1820>
- Kovatcheva-Datchary P, Nilsson A, Akrami R, Lee YS, De Vadder F, Arora T, Hallen A, Martens E, Björck I, Backhed F (2015) Dietary fiber-induced improvement in glucose metabolism is associated with increased abundance of prevotella. Cell Metab 22(6):971–982. <https://doi.org/10.1016/j.cmet.2015.10.001>
- Kozłowska M, Laudy AE, Przybyl J, Ziarno M, Majewska E (2015) Chemical composition and antibacterial activity of some medicinal plants from Lamiaceae family. Acta Pol Pharm 72(4):757–767
- Kuzuya T, Kanazawa Y, Kosaka K (1969) Stimulation of insulin secretion by xylitol in dogs. Endocrinology 84(2):200–207. <https://doi.org/10.1210/endo-84-2-200>
- Kuzuya T, Kanazawa Y, Hayashi M, Kikuchi M, Ide T (1971) Species difference in plasma insulin responses to intravenous xylitol in man and several mammals. Endocrinol Jpn 18(4):309–320. <https://doi.org/10.1507/endocrj1954.18.309>
- Lee BD, Park MK (2014) Effects and safety of xylitol on middle ear epithelial cells. J Int Adv Otol 10(1):19–24
- Lemos JA, Palmer SR, Zeng L, Wen ZT, Kajfasz JK, Freires IA, Abranches J, Brady LJ (2019) The biology of *Streptococcus mutans*. Microbiol Spectr 7(1). <https://doi.org/10.1128/microbiolspec.GPP3-0051-2018>
- Lowe C, Anthony J (2020) Pilot study of the effectiveness of a xylitol-based drinking water additive to reduce plaque and calculus accumulation in dogs. Can Vet J 61(1):63–68
- Ly KA, Milgrom P, Rothen M (2006) Xylitol, sweeteners, and dental caries. Pediatr Dent 28(2):154–163 discussion 192–8
- Lynch H, Milgrom P (2003) Xylitol and dental caries: an overview for clinicians. J Calif Dent Assoc 31(3):205–209
- Maguire A, Rugg-Gunn AJ (2003) Xylitol and caries prevention — is it a magic bullet? Br Dent J 194(8):429–436. <https://doi.org/10.1038/sj.bdj.4810022>
- Mäkinen KK (2010) Sugar alcohols, caries incidence, and remineralization of caries lesions: a literature review. Int J Dent 2010:981072–981023. <https://doi.org/10.1155/2010/981072>

- Mäkinen KK (2011) Sugar alcohol sweeteners as alternatives to sugar with special consideration of xylitol. *Med Princ Pract* 20(4):303–320. <https://doi.org/10.1159/000324534>
- Mäkinen KK (2016) Gastrointestinal disturbances associated with the consumption of sugar alcohols with special consideration of xylitol: scientific review and instructions for dentists and other health-care professionals. *Int J Dent* 2016:5967907–5967916. <https://doi.org/10.1155/2016/5967907>
- Marchese A, Orhan IE, Daglia M, Barbieri R, Di Lorenzo A, Nabavi SF, Gortzi O, Izadi M, Nabavi SM (2016) Antibacterial and antifungal activities of thymol: a brief review of the literature. *Food Chem* 210: 402–414. <https://doi.org/10.1016/j.foodchem.2016.04.111>
- Marghalani AA, Guinto E, Phan M, Dhar V, Tinanoff N (2017) Effectiveness of xylitol in reducing dental caries in children. *Pediatr Dent* 39(2):103–110
- Medvid YO (2011) Optimization of complex treatment and correction of endointoxication of patients with diffuse phlegmons in maxillofacial area [in Ukrainian]. PhD (Dentistry) manuscript, University of Lviv
- Milgrom P, Ly KA, Rothen M (2009a) Xylitol and its vehicles for public health needs. *Adv Dent Res* 21(1):44–47. <https://doi.org/10.1177/0895937409335623>
- Milgrom P, Ly KA, Tut OK, Mancil L, Roberts MC, Briand K, Gancio MJ (2009b) Xylitol pediatric topical oral syrup to prevent dental caries: a double-blind randomized clinical trial of efficacy. *Arch Pediatr Adolesc Med* 163(7):601–607. <https://doi.org/10.1001/archpediatrics.2009.77>
- Mooradian AD, Smith M, Tokuda M (2017) The role of artificial and natural sweeteners in reducing the consumption of table sugar: a narrative review. *Clin Nutr ESPEN* 18:1–8. <https://doi.org/10.1016/j.clnesp.2017.01.004>
- Nayak PA, Nayak UA, Khandelwal V (2014) The effect of xylitol on dental caries and oral flora. *Clin Cosmet Investig Dent* 6:89–94. <https://doi.org/10.2147/CCIDE.S55761>
- Newton JT, Awojobi O, Nasseripour M, Warburton F, Di Giorgio S, Gallagher JE, Banerjee A (2019) A systematic review and meta-analysis of the role of sugar-free chewing gum in dental caries. *JDR Clin Trans Res*. <https://doi.org/10.1177/2380084419887178>
- Oku T, Nakamura S (2007) Threshold for transitory diarrhea induced by ingestion of xylitol and lactitol in young male and female adults. *J Nutr Sci Vitaminol (Tokyo)* 53(1):13–20. <https://doi.org/10.3177/jnsv.53.13>
- Olenikov DN, Rokhin AV, Tankhaeva LM (2009) Lamiaceae carbohydrates. VI. Water-soluble polysaccharides from *Lophanthus chinensis*. *Chem Nat Compd* 45(3):300–303. <https://doi.org/10.1007/s10600-009-9358-2>
- Ortiz ME, Bleckwedel J, Raya RR, Mozzi F (2013) Biotechnological and in situ food production of polyols by lactic acid bacteria. *Appl Microbiol Biotechnol* 97(11):4713–4726. <https://doi.org/10.1007/s00253-013-4884-z>
- Palchaudhuri S, Palchaudhuri A, Chatterjee B (2015) Xylitol in prevention of dental caries and respiratory disease caused by *Diplococcus streptococcus*. *Arch Med Microbiol* 6(4):4
- Pearlman M, Obert J, Casey L (2017) The Association between artificial sweeteners and obesity. *Curr Gastroenterol Rep* 19(12):64. <https://doi.org/10.1007/s11894-017-0602-9>
- Petersen PE (2004) Challenges to improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. *Int Dent J* 54(6 Suppl 1):329–343. <https://doi.org/10.1111/j.1875-595x.2004.tb00009.x>
- Peterson ME (2013) Xylitol. *Top Companion Anim Med* 28(1):18–20. <https://doi.org/10.1053/j.tcam.2013.03.008>
- Rafeek R, Carrington CVF, Gomez A, Harkins D, Torralba M, Kuelbs C, Addae J, Moustafa A, Nelson KE (2019) Xylitol and sorbitol effects on the microbiome of saliva and plaque. *J Oral Microbiol* 11(1): 1536181. <https://doi.org/10.1080/20002297.2018.1536181>
- Reshamwala SMS, Lali AM (2020) Exploiting the NADPH pool for xylitol production using recombinant *Saccharomyces cerevisiae*. *Biotechnol Prog*. <https://doi.org/10.1002/btpr.2972>
- Rice T, Zannini E, E KA, Coffey A (2019) A review of polyols - biotechnological production, food applications, regulation, labeling and health effects. *Crit Rev Food Sci Nutr*. <https://doi.org/10.1080/10408398.2019.1625859>
- Riggs E, Kilpatrick N, Slack-Smith L, Chadwick B, Yelland J, Muthu MS, Gomersall JC (2019) Interventions with pregnant women, new mothers and other primary caregivers for preventing early childhood caries. *Cochrane Database Syst Rev* 2019(11). <https://doi.org/10.1002/14651858.CD012155.pub2>
- Riley P, Moore D, Ahmed F, Sharif MO, Worthington HV (2015) Xylitol-containing products for preventing dental caries in children and adults. *Cochrane Database Syst Rev* 3:CD010743. <https://doi.org/10.1002/14651858.CD010743.pub2>
- Ruiz-Ojeda FJ, Plaza-Diaz J, Saez-Lara MJ, Gil A (2019) Effects of sweeteners on the gut microbiota: a review of experimental studies and clinical trials. *Adv Nutr* 10(suppl_1):S31–S48. <https://doi.org/10.1093/advances/nmy037>
- Ruiz-Ojeda FJ, Plaza-Diaz J, Saez-Lara MJ, Gil A (2020) Corrigendum for Ruiz-Ojeda et al. Effects of sweeteners on the gut microbiota: a review of experimental studies and clinical trials. *Advances in Nutrition*, Volume 10, Issue suppl_1, January 2019, Pages S31–S48. *Adv Nutr* 11(2):468. <https://doi.org/10.1093/advances/nmz112>
- Sakallioglu O, Guvenc IA, Cingi C (2014) Xylitol and its usage in ENT practice. *J Laryngol Otol* 128(7):580–585. <https://doi.org/10.1017/S0022215114001340>
- Salli KM, Forssten SD, Lahtinen SJ, Ouwehand AC (2016) Influence of sucrose and xylitol on an early *Streptococcus mutans* biofilm in a dental simulator. *Arch Oral Biol* 70:39–46. <https://doi.org/10.1016/j.archoralbio.2016.05.020>
- Salli K, Lehtinen MJ, Tiitonen K, Ouwehand AC (2019) Xylitol's health benefits beyond dental health: a comprehensive review. *Nutrients* 11(8). <https://doi.org/10.3390/nu11081813>
- Santos MG, Carpinteiro DA, Thomazini M, Rocha-Selmi GA, da Cruz AG, Rodrigues CEC, Favaro-Trindade CS (2014) Coencapsulation of xylitol and menthol by double emulsion followed by complex coacervation and microcapsule application in chewing gum. *Food Res Int* 66:454–462. <https://doi.org/10.1016/j.foodres.2014.10.010>
- Sarkar P, Mukherjee M, Goswami G, Das D (2020) Adaptive laboratory evolution induced novel mutations in *Zymomonas mobilis* ATCC ZW658: a potential platform for co-utilization of glucose and xylose. *J Ind Microbiol Biotechnol* 47(3):329–341. <https://doi.org/10.1007/s10295-020-02270-y>
- Sene L, Arruda PV, Oliveira SM, Felipe MG (2011) Evaluation of sorghum straw hemicellulosic hydrolysate for biotechnological production of xylitol by *Candida guilliermondii*. *Braz J Microbiol* 42(3): 1141–1146. <https://doi.org/10.1590/S1517-838220110002000036>
- Shanaida M, Kernychna I, Shanaida Y (2017) Chromatographic Analysis of organic acids, amino acids, and sugars in *Ocimum americanum* L. *Acta Pol Pharm* 74(2):729–734
- Shyama M, Honkala E, Honkala S, Al-Mutawa SA (2006) Effect of xylitol candies on plaque and gingival indices in physically disabled school pupils. *J Clin Dent* 17(1):17–21
- Silva TC, Pereira AFF, Machado MAAM, Buzalaf MAR (2009) The use of xylitol as a strategy for prevention of dental caries. *Revista Odonto Ciencia* 24(2):205–212
- Singh S, Hornick D, Fedler J, Launsbach JL, Teresi ME, Santacroce TR, Cavanaugh JE, Horan R, Nelson G, Starnes TD, Zabner J, Durairaj L (2020) Randomized controlled study of aerosolized hypertonic xylitol versus hypertonic saline in hospitalized patients with pulmonary exacerbation of cystic fibrosis. *J Cyst Fibros* 19(1):108–113. <https://doi.org/10.1016/j.jcf.2019.06.016>
- Soderling E (2009a) Controversies around xylitol. *Eur J Dent* 3(2):81–82

- Soderling EM (2009b) Xylitol, mutans streptococci, and dental plaque. *Adv Dent Res* 21(1):74–78. <https://doi.org/10.1177/0895937409335642>
- Steinberg LM, Odusola F, Mandel ID (1992) Remineralizing potential, antiplaque and antigingivitis effects of xylitol and sorbitol sweetened chewing gum. *Clin Prev Dent* 14(5):31–34
- Stoklosa RJ, Nghiem NP, Latona RJ (2019) Xylose-enriched ethanol fermentation stillage from sweet sorghum for xylitol and astaxanthin production. *Fermentation* 5(4):84
- Storey D, Lee A, Bornet F, Brouns F (2007) Gastrointestinal tolerance of erythritol and xylitol ingested in a liquid. *Eur J Clin Nutr* 61(3):349–354. <https://doi.org/10.1038/sj.ejcn.1602532>
- Sylvetsky A, Rother KI, Brown R (2011) Artificial sweetener use among children: epidemiology, recommendations, metabolic outcomes, and future directions. *Pediatr Clin N Am* 58(6):1467–14xi. <https://doi.org/10.1016/j.pcl.2011.09.007>
- Takahashi K, Fukazawa M, Motohira H, Ochiai K, Nishikawa H, Miyata T (2003) A pilot study on antiplaque effects of mastic chewing gum in the oral cavity. *J Periodontol* 74(4):501–505. <https://doi.org/10.1902/jop.2003.74.4.501>
- Takeuchi K, Asakawa M, Hashiba T, Takeshita T, Saeki Y, Yamashita Y (2018) Effects of xylitol-containing chewing gum on the oral microbiota. *J Oral Sci* 60(4):588–594. <https://doi.org/10.2334/josnurd.17-0446>
- Talattof Z, Azad A, Zahed M, Shahradnia N (2018) Antifungal activity of xylitol against *Candida albicans*: an in vitro study. *J Contemp Dent Pract* 19(2):125–129. <https://doi.org/10.5005/jp-journals-10024-2225>
- Tamura M, Hoshi C, Hori S (2013) Xylitol affects the intestinal microbiota and metabolism of daidzein in adult male mice. *Int J Mol Sci* 14(12):23993–24007. <https://doi.org/10.3390/ijms141223993>
- Teixeira Essenfelder L, Gomes AA, Miquelutti D, da Silva GF, Magalhães MLB (2019) Effect of xylitol on salivary β -glucosidase in humans. *Eur J Oral Sci* 127(5):472–475. <https://doi.org/10.1111/eos.12649>
- Tennant DR (2014) Potential intakes of total polyols based on UK usage survey data. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 31(4):574–586. <https://doi.org/10.1080/19440049.2014.886132>
- Thabuis C, Cheng CY, Wang X, Pochat M, Han A, Miller L, Wils D, Guerin-Deremaux L (2013) Effects of maltitol and xylitol chewing-gums on parameters involved in dental caries development. *Eur J Paediatr Dent* 14(4):303–308
- Uebanso T, Kano S, Yoshimoto A, Naito C, Shimohata T, Mawatari K, Takahashi A (2017) Effects of consuming xylitol on gut microbiota and lipid metabolism in mice. *Nutrients* 9(7). <https://doi.org/10.3390/nu9070756>
- Uhari M, Tapiainen T, Kontiokari T (2000) Xylitol in preventing acute otitis media. *Vaccine* 19(Suppl 1):S144–S147. [https://doi.org/10.1016/s0264-410x\(00\)00294-2](https://doi.org/10.1016/s0264-410x(00)00294-2)
- Ur-Rehman S, Mushtaq Z, Zahoor T, Jamil A, Murtaza MA (2015) Xylitol: a review on bioproduction, application, health benefits, and related safety issues. *Crit Rev Food Sci Nutr* 55(11):1514–1528. <https://doi.org/10.1080/10408398.2012.702288>
- Vasilescu R, Ionescu A, Mihai A, Carniciu S, Ionescu-Tîrgoviște C (2011) Sweeteners and metabolic diseases: xylitol as a new player. In: *Proc Rom Acad Series B* 2:125–128
- Verkaik MJ, Busscher HJ, Jager D, Slomp AM, Abbas F, van der Mei HC (2011) Efficacy of natural antimicrobials in toothpaste formulations against oral biofilms in vitro. *J Dent* 39(3):218–224. <https://doi.org/10.1016/j.jdent.2010.12.007>
- Vernacchio L, Vezina RM, Mitchell AA (2007) Tolerability of oral xylitol solution in young children: implications for otitis media prophylaxis. *Int J Pediatr Otorhinolaryngol* 71(1):89–94. <https://doi.org/10.1016/j.ijporl.2006.09.008>
- Vernacchio L, Corwin MJ, Vezina RM, Pelton SI, Feldman HA, Coyne-Beasley T, Mitchell AA (2014) Xylitol syrup for the prevention of acute otitis media. *Pediatrics* 133(2):289–295. <https://doi.org/10.1542/peds.2013-2373>
- Wolnerhanssen BK, Meyer-Gerspach AC (2019) Health effects of sugar consumption and possible alternatives. *Ther Umsch* 76(3):111–116. <https://doi.org/10.1024/0040-5930/a001070>
- Wolnerhanssen BK, Meyer-Gerspach AC, Beglinger C, Islam MS (2019) Metabolic effects of the natural sweeteners xylitol and erythritol: a comprehensive review. *Crit Rev Food Sci Nutr*. <https://doi.org/10.1080/10408398.2019.1623757>
- Xu Y, Chi P, Bilal M, Cheng H (2019) Biosynthetic strategies to produce xylitol: an economical venture. *Appl Microbiol Biotechnol* 103(13):5143–5160. <https://doi.org/10.1007/s00253-019-09881-1>
- Yi EY, Kim YJ (2013) Xylitol inhibits in vitro and in vivo angiogenesis by suppressing the NF-kappaB and Akt signaling pathways. *Int J Oncol* 43(1):315–320. <https://doi.org/10.3892/ijo.2013.1917>
- Yuan X, Tu S, Lin J, Yang L, Shen H, Wu M (2020) Combination of the CRP mutation and ptsG deletion in *Escherichia coli* to efficiently synthesize xylitol from corncob hydrolysates. *Appl Microbiol Biotechnol* 104(5):2039–2050. <https://doi.org/10.1007/s00253-019-10324-0>
- Zhou G, Peng H, Wang YS, Huang XM, Xie XB, Shi QS (2019) Enhanced synergistic effects of xylitol and isothiazolones for inhibition of initial biofilm formation by *Pseudomonas aeruginosa* ATCC 9027 and *Staphylococcus aureus* ATCC 6538. *J Oral Sci* 61(2):255–263. <https://doi.org/10.2334/josnurd.18-0102>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.