

Pharmacologic Options for the Treatment of Sarcopenia

John E. Morley¹

Received: 5 May 2015 / Accepted: 5 June 2015
© Springer Science+Business Media New York 2015

Abstract Sarcopenia is now clinically defined as a loss of muscle mass coupled with functional deterioration (either walking speed or distance **or** grip strength). Based on the FRAX studies suggesting that the questions without bone mineral density can be used to screen for osteoporosis, there is now a valid simple questionnaire to screen for sarcopenia, i.e., the SARC-F. Numerous factors have been implicated in the pathophysiology of sarcopenia. These include genetic factors, mitochondrial defects, decreased anabolic hormones (e.g., testosterone, vitamin D, growth hormone and insulin growth hormone-1), inflammatory cytokine excess, insulin resistance, decreased protein intake and activity, poor blood flow to muscle and deficiency of growth derived factor-11. Over the last decade, there has been a remarkable increase in our understanding of the molecular biology of muscle, resulting in a marked increase in potential future targets for the treatment of sarcopenia. At present, resistance exercise, protein supplementation, and vitamin D have been established as the basic treatment of sarcopenia. High-dose testosterone increases muscle power and function, but has a number of potentially limiting side effects. Other drugs in clinical development include selective androgen receptor molecules, ghrelin agonists, myostatin antibodies, activin IIR antagonists, angiotensin converting enzyme inhibitors, beta antagonists, and fast skeletal muscle troponin activators. As sarcopenia is a major predictor of frailty, hip fracture,

disability, and mortality in older persons, the development of drugs to treat it is eagerly awaited.

Keywords Sarcopenia · Muscle loss · Frailty · Muscle function · Low muscle mass

Introduction

Sarcopenia was originally defined as the age-related loss of muscle mass [1]. Subsequently, it became obvious to clinicians that it was muscle quality, rather than muscle mass that determined the function of muscle [2, 3]. This led to the suggestion that it was muscle power (force x velocity) which should be utilized to determine the role of muscle in determining outcomes. It was suggested that this should be termed dynapenia [4]. From this developed the concept of a sarcopenia-disability cascade (Table 1). Each component of this cascade can be separately measured and theoretically would lead to worse outcomes.

However, in 2010, Cruz-Jentoft et al. [5] published the “European Consensus on Definition and Diagnosis of Sarcopenia.” They redefined sarcopenia as being muscle loss coupled with a decline in function (either walking speed or grip strength). This definition was validated as having a strong predictive ability of poor outcomes [6–8]. Subsequently, 4 other definitions of sarcopenia, all using gait speed and grip strength, as well as some measurement of low muscle mass have been published [9–12]. Each uses slightly different cut off points and 2 recognized the importance of having different cut offs for different ethnic groups. Woo et al. [13] compared each of these definitions and found that they had slightly different predictive abilities. Of the definitions, the Foundation of NIH (FNIH)

✉ John E. Morley
morley@slu.edu

¹ Division of Geriatric Medicine, Saint Louis University
School of Medicine, 1402 S. Grand Blvd., M238, St. Louis,
MO 63104, USA

Table 1 The sarcopenia-disability cascade

Term	Definition	Measurement
Sarcopenia	Loss of muscle mass not due to cachexia or peripheral vascular disease	Dual energy X-ray absorptiometry MRI/CT Ultrasound Bioelectrical impedance Midarm muscle circumference Calf circumference
Kratopenia	Loss of force i.e., strength	Isometric (dynametry) Isotonic
Dynapenia	Loss of power i.e., force × velocity	Walking speed Walking distance Stair climbing Jebson hand function
Frailty	Physical phenotype (fatigue, resistance, aerobic, illness, loss of weight)	CHS (fried) criteria FRAIL questionnaire Study of osteoporotic fractures criteria Canadian (Rockwood) criteria
Disability	Loss of activities of daily living (ADLs)	Katz ADLs Barthel index Functional index measure

The modern definition of sarcopenia is a combination for sarcopenia and dynapenia or kratopenia (grip strength)

Table 2 The SARC-F questionnaire for sarcopenia

Component	Question	Scoring
Strength	How much difficulty do you have in lifting and carrying 10 pounds?	None = 0 Some = 1 A lot or unable = 2
Assistance in walking	How much difficulty do you have walking across a room?	None = 0 Some = 1 A lot, use aids, or unable = 2
Rise from a chair	How much difficulty do you have transferring from a chair or bed?	None = 0 Some = 1 A lot or unable without help = 2
Climb stairs	How much difficulty do you have climbing a flight of ten stairs?	None = 0 Some = 1 A lot or unable = 2
Falls	How many times have you fallen in the last year?	None = 0 1–3 falls = 1 4 or more falls = 2

Scoring: 1–10 total points possible; 0–2 for each component; 0 = best, 10 = worst; 0–3 healthy; ≥ 4 is symptomatic for sarcopenia


sarcopenia criteria using gait speed, but not grip strength, had slightly better predictive value for poor outcomes.

Based on the parallels between osteoporosis and sarcopenia and the finding that the 6 FRAX questions without Bone Mineral Density are highly predictive of fracture risk [14], we developed a simple sarcopenic questionnaire to predict poor muscle function (Table 2)

[15]. This questionnaire has been shown to be a valid predictor of poor outcomes similar to that of the FNIH (walking speed) definition in both the United States and Asia [13, 16–18].

Sarcopenia has multiple causes and, as older persons develop a variety of diseases with increased production of cytokines, it may overlap with cachexia [19]. In this

Fig. 1 Aging, exercise, and muscle injury



review, we will first explore the physiological causes of sarcopenia with a special emphasis on potential pharmaceutical targets. We will then review the available and developing treatments for sarcopenia.

The Pathophysiology of Sarcopenia

When muscle contracts this activates mechanoreceptors, i.e., titin and dystroglycan, and causes muscle injury. The mechanoreceptors increase the activity of muscle growth factors (IGF1-Ea and muscle growth factor) which increase muscle protein synthesis and recruit satellite cells and motor units. This leads to muscle regeneration and increased muscle function (Fig. 1). With aging, there is increased muscle injury with a decrease in muscle regeneration and function. This is due to a decrease in muscle growth factors leading to a reduction in the protein synthesis/degradation cycle and the activation of satellite cells and motor units. Anatomically, with aging there is Type II fiber atrophy resulting in decreased muscle mass, strength, and power [20].

Old muscle shows fiber size heterogeneity and fiber grouping with an increase in myosin heavy chain [21]. This differs from cachexia where fiber size variability is not seen. This is similar to the histological changes seen with Amyotrophic Lateral Sclerosis. Sarcopenic patients have a reduction in the motor unit number index (MUNIX) which is intermediate between that seen in healthy older persons and in patients with Amyotrophic Lateral Sclerosis [22]. Further evidence of motor neuron degeneration is the increase in C-terminal agrin fragments in about a third of sarcopenic patients [23]. With aging, there is a 25 % loss of motoneurons leading to sprouting of small motor neurons


that innervate Type II fibers leading to an eventual loss of type II fibers [24]. Circulating levels of ciliary neurotrophic factor (CNTF), which stimulate motor unit formation, decline with aging [25, 26]. Older persons who have the null allele rs1800169 for CNTF have lower grip strength [27]. Axokine, a modified version of CNTF, was tried for weight loss due to its anorectic properties. The trials were suspended when subjects developed antibodies to CNTF.

Myokines

Besides CNTF, skeletal muscle produces a variety of myokines that can modulate muscle growth and repair (Fig. 2) [28–30]. Some of these, such as interleukin-6, may be predominantly produced by adipose tissue infiltrating muscle [31]. A number of these myokines have direct effects on muscle such as IGF-1, IGF binding proteins, myostatin, musclin, leukemia inhibitory factor and CXCL-1. Others such as VEGF-B, IL-8 and Follistatin-like 1 increase angiogenesis in muscles. Proteomics of muscle secretions should lead to the discovery of many more myokines that modulate muscle growth [30].

IL-15 is an inflammatory cytokine produced by muscle that increases contractile protein accumulation and causes myotube hypertrophy [32]. In vivo IL-15 reduces fast muscle fatigue and enhanced oxidative metabolism. Fatigue is an important component of the frailty phenotype that is separate from sarcopenia [33, 34]. IL-15 agonists or stimulants could be useful for the treatment of fatigue in older persons. Fatigue is, in part, related to a loss of muscle. A number of studies are examining the role of IL-15 agonists in advanced cancer (www.clinicaltrials.gov).

Fig. 2 Myokines and their target sites. *SPARC* secreted protein acidic and rich in cysteine, *IL* interleukin, *GDF* growth differentiation factor, *FGF* fibroblast growth factor, *BMP* bone morphogenetic protein, *LIF* leukemia inhibitory factor, *CXCL-1* chemokine ligand 1, *VEGF-B* Vascular endothelial growth factor, *CNTF* ciliary neurotrophic-derived factor



Genetics

Genes play a role in 65 % of the muscle mass and 50–80 % of muscle strength in older persons [35]. Studies on genes and muscle mass and strength have been rudimentary, with a number of results being controversial [36]. Angiotensin converting enzyme alleles have been shown to play a role in efficiency of muscle contraction. The alpha-actin 3 is responsible for anchoring the actin filaments to the 2-disk in fast twitch fibers. An ACTN3 gene deficiency is associated with reduced power and with endurance activity in men [37, 38]. Bradykinin increases muscle blood flow and the B₂R gene associated with high function is more frequent in endurance athletes. Other genes associated with muscle strength include CNTF, IL-15, collagen type, insulin growth factor II, myostatin, the vitamin D receptor, and the androgen CAG receptor. In older persons, expression of genes for insulin growth factor-1, myostatin, matrix metalloproteinase-2, ciliary neurotrophic factor, and myostatin correlated most optimally with training-induced strength gains [39]. Variants in the activin receptor 1B play an important role in human muscle strength [40]. Perilipin 2 is a protein associated with lipid droplets. Perilipin 2 is higher in older persons and is related to a decline in muscle strength and proteins associated with muscle atrophy, viz MURF1, and atrogin [41]. This suggests that expression of perilipin 2 may play a role in the development of obese sarcopenia [42, 43].

Mitochondria

The peroxisome proliferator-activated receptor γ coactivator 1 α (PGC-1 α) regulates mitochondrial biogenesis and function and regulates muscle fiber adaptation to exercise [44]. There is a reduction of PGC-1 α gene expression in old animals and older persons [45]. This reduction results in a low-grade inflammatory reaction with increased levels of IL-6 and TNF α . PGC-1 α activity decreases functional loss of mitochondrial enzymes in old animals and protects muscle from damage [46]. Biochemically, PGC-1 α inhibits FOXO and NF κ B and thus decreases autophagy and the ubiquitin–proteasome systems. PGC-1 α promotes mitochondrial biogenesis and fusion, thus maintaining ATP levels and reducing AMPK. Excess expression of PGC-1 α damages heart and muscle. It has been suggested that increasing PGC-1 α levels in sarcopenic tissue to physiological levels may be a key therapeutic approach to treating muscle wasting.

Mitochondrial dysfunction plays a major role in the pathogenesis of aging [47, 48]. Mitochondria control the production of cellular energy, free radical signaling, and can activate apoptotic pathways. The importance of bioenergetics in the development of sarcopenia is demonstrated by the correlation of ATP synthesis/oxygen consumption and walking speed in older persons [49]. Walking speed is a key component of the modern definitions of sarcopenia. In addition, with aging, there is often increased

protein 32 (TRIM32) directly control myofibril breakdown. MURF1 attacks the myosin binding protein and the myosin light chain eventually leading to destruction of the thick myosin filament. TRIM32 destroys desmin and then the Z-band and eventually the thin actin filament. At the same time TRIM32 directly inhibits PI3 K-AKT activity resulting in increased proteolysis. Myofibrils constitute the vast majority of muscle protein and their destruction leads to loss of muscle function [57, 58]. There is an increase in protein destruction and ubiquitination in sarcopenia. Obviously, inhibitors of TRIM32 and/or MURF1 represent attractive therapeutic targets for treating sarcopenia.

MicroRNAs

MicroRNAs (miRNA) are small molecules which regulate posttranscriptional gene function by silencing RNA. They are cleaved in the nucleus by Droscha and then exported into the cytoplasm, where they are processed by DICER and combined with AGO to form RNA-induced silencing complexes. These then repress translation of mRNAs. Satellite cells decline with aging [59]. miRNAs play a central role in satellite cell quiescence [60]. The miRNAs (miR-1, miR-208, and MiR486) regulate satellite cell renewal by modulating Pax7. Decreased skeletal muscle miRNA expression in older persons is associated with a decrease in the function of the IGF-1/PI₃K/AKT pathway [61]. Exercise modulates the response of a number of muscle-specific miRNAs. There is evidence that miRNAs are sensitive to a variety of drugs such as mu opioids [62] and drugs for Parkinson's disease [63]. There appears to be tremendous potential to treat sarcopenia by modulating miRNAs. As more is known about the role of miRNAs in modulating muscle growth, it will also become possible to modulate specific RNAs by mimicking the positive patterns with phosphorothiolated antisenses.

Electrical stimulation of both thighs for 9 weeks improved timed up and go test, walking speed and 5 time chair rise [64]. This training led to an increase in diameter and percent of fast fibers. There was a stimulation of IGF-1 isoforms with a reduction of MuRF-1 and atrogen-1 leading to a reduction in proteolysis. Electrical stimulation also produces an increase of satellite cells. Electrical stimulation increased miR-29 which would decrease fibrotic infiltration of muscle. Overall, this study strongly supports the concept of electrical muscle stimulation for treatment of sarcopenia.

Parabiosis

Studies with parabiotic mice have shown that the combination of a young and an old mouse leads to rejuvenation in

the muscle of the older mouse [65]. This was due to an increase in Notch signaling leading to an increase in satellite cells. The humoral agent responsible for this appears, in a large part, to be growth differentiation factor 11 (GDF11) [66]. Sinha et al. [67] have demonstrated that testosterone plays a permissive role in muscle mass and fiber cross-sectional area in parabiotic mice. These data suggest a role for GDF11 in treatment of sarcopenia.

Hibernating animals maintain their muscle structure during winter, raising the question of whether a circulating factor, in addition to shivering thermogenesis, is responsible for the protection of muscle during hibernation. The extensor digitorum longus muscle of the rat, when incubated with serum obtained from hibernating bears had a 40 % decrease in proteolysis, associated with a decline in cathepsin B and ubiquitin [68]. During hibernation, there is an increase in PGC-1 α , which is associated with a decrease in FOXO and MURF-1 [69]. Serum- and glucocorticoid-inducible kinase 1 (SGK1) has been shown to downregulate proteolysis, autophagy and increase protein synthesis in hibernating animals [70]. This bypasses the classical AKT-FOXO pathway. SGK-1 may represent an important therapeutic target to prevent atrophy-induced muscle loss.

Table 3 provides a list of potential targets for future drug development. Figure 4 gives an overview of the major factors so far demonstrated to be a component of the pathophysiology of sarcopenia. Many of these already have drugs available or under development. These and their physiological rationale will be discussed in the next section.

Management of Sarcopenia

The primary treatment of sarcopenia is resistance exercise [71–73]. As was shown by the LIFE study, aerobic exercise can also decrease functional decline in lower limb muscles [74]. Exercise has also been shown to be an important therapeutic approach to reversing frailty [75]. There is evidence to support that excess protein [1–1.2 g (kj day)] may also enhance muscle mass and, to a lesser extent, function [76–80]. This is particularly true for leucine enriched essential amino acids (whey protein) [81]. Essential amino acid supplementation prevents muscle mass loss due to bed rest [82]. A recent multicenter study has shown that whey protein together with vitamin D increased both muscle mass and stair climb [83]. There is some evidence for synergistic effects of exercise and protein to enhance muscle function [84–86]. Vitamin D supplementation increases muscle strength without increasing muscle mass or power [87]. Vitamin D is more effective in older persons and those with low vitamin D levels. It also decreases falls in persons who are vitamin D deficient [88].

testosterone in combination with a protein supplement decreased hospitalization [105].

In lower doses, testosterone increases protein synthesis resulting in an increase in muscle mass [106, 107]. In high doses, testosterone activates satellite cell recruitment and reduces adipose stem cells [108]. Testosterone effects on muscle cells bypass the WNT system and activate β -catenin [109]. This leads to increased myogenesis and cell cycling and decreased adipogenesis.

While testosterone as a therapeutic agent has been utilized since the 1940s, there is a fear that it will produce excessive side effects [110]. A meta-analysis of the controlled studies of testosterone in older males found no increase in mortality [111]. Whether or not it increases cardiovascular events, and particularly in the first 3 months after administration, remains controversial [112, 113]. Persons with diabetes mellitus have accelerated sarcopenia [114–116], and a recent study in diabetics found a decrease in mortality in diabetics receiving testosterone [117]. Nevertheless, this fear of negative effects from testosterone has driven the exploration for selective androgen receptor modulators (SARMs) which may be, theoretically, safer. At present, of the drugs developed and being developed for sarcopenia, testosterone remains the most efficacious and safest. In view of the fact that testosterone also increases bone mineral density and bone strength [118, 119] and osteoporosis often co-exists with sarcopenia (osteosarcopenia), it would seem that more clinical attention should be paid to the potential role of testosterone for treating sarcopenia. Two major trials are presently underway and these may help determine the place of testosterone in the management of sarcopenia, osteoporosis, and frailty (The Testosterone Trial in Older Men—www.clinicaltrials.gov and the T4DM trial—www.t4dm.org.au).

Anabolic Steroids/Selective Androgen Receptor Modulators (SARMs)

Nandrolone is an injectable anabolic steroid. It increased fiber area and muscle mass, but there is no evidence that it increased strength [120–122]. In three studies of persons with hip fracture, it had a nonstatistical improvement in functional status [123].

MK0773 (TFM-4AS-1) is a 4-aza steroidal drug that has androgen gene selectivity. In females, it increased IGF-1 as well as stair climbing power and gait speed [124]. This study was terminated because of an increased signal for cardiac failure. In women with sarcopenia, it increased muscle mass, bilateral leg press, and stair climbing power but not gait speed (www.clinicaltrials.gov). The study in males was reported at the 90th Endocrine Society in 2008. It showed anabolic effects of MK0773.

SARMs are androgen receptor ligands that bind to the androgen receptor with differing sensitivity compared to testosterone [125]. Steroidal SARMs were first developed in the 1940s. More recently, a number of nonsteroidal SARMs have been developed [126].

LGD-4033 is a nonsteroidal, orally active SARM. The phase I trial showed an increase in muscle mass, but no effect on fat mass in a 21-day trial [127]. BMS-564929 is also in phase I trials.

In a 12-week study, enobosarm increased total lean mass and stair climb [128]. In female patients with cancer, enobosarm increased lean mass compared to baseline, but not significantly compared to placebo [129]. In 2 phase 3 trials, it maintained body mass and improved stair climb in one of the 2 trials in patients with cancer [130].

Overall, these studies of SARMs have shown no advantage over testosterone.

Growth Hormone/Insulin Growth Factor-1

Rudman et al. [131] originally showed that growth hormone increased lean body mass in older men. The excitement created by their original data was dampened by finding that a growth hormone treatment for a year led to a variety of side effects such as carpal tunnel syndrome and gynecomastia [132]. Subsequently, growth hormone has been shown to increase muscle mass but not muscle strength in older persons [133]. A combination of growth hormone and testosterone increased muscle mass at 8 weeks and 1 repetition maximum strength only by week 17 [134]. Growth hormone, which produces its effects through the release of liver-derived IGF-1, also increased nitrogen retention [135]. Adverse effects include joint and muscle pain, edema, carpal tunnel syndrome, and hyperglycemia [136].

There are marked reductions of circulating IGF-1 with aging [137]. Both low and high levels of IGF-1 are associated with increased risk of cardiovascular disease. Similarly, there is limited evidence that circulating IGF-1 is associated with muscle power. A single small study of IGF-1 in older persons found an increase in side effects viz. orthostatic hypotension, gynecomastia, myositis, and edema [138].

Ghrelin

Ghrelin is produced from the fundus of the stomach. It increases food intake and growth hormone. These effects of ghrelin are due to the hypothalamic release of nitric oxide [139]. Ghrelin increased food intake and produced muscle mass gain in persons with cancer [140, 141]. The ghrelin agonist, anamorelin, increased food intake, and muscle mass, but not strength in persons with cancer cachexia

[142, 143]. Macimorelin is another ghrelin agonist under development.

Capromorelin, a ghrelin receptor agonist, was tested in older sarcopenic individuals [144]. It increased lean mass, tandem walk, and stair climb at the end of treatment for a year. MK-0677, which also activates the ghrelin receptor to increase growth hormone, was studied for 24 weeks in persons with hip fracture [145]. Over this period, it increased ability to stair climb and also decreased falls. The treatment was associated with an increase in heart failure.

Overall, while ghrelin agonists will increase food intake and muscle mass, it is unlikely that they will produce a significant effect on function in persons with sarcopenia.

Myostatin and Activin II Receptor Inhibitors

Myostatin or growth differentiation factor-8 is produced in skeletal muscle and prevents muscle growth and satellite cell production [146]. Myostatin activates the Activin IIR receptor to increase SMAD (Fig. 3). Lack of myostatin in animals leads to “double muscled” cows (Belgian Blue and Piedmontese). Heterozygote deletion of myostatin in whippets leads to an increase in running ability, while homozygotes are more muscular but not as good runners. A homozygote muscle deletion of myostatin in a young boy resulted in an increase in muscle mass [147].

In humans, creatine, together with resistance exercise, results in an amplification of the normal decrease in myostatin with resistance training alone [148]. Creatine produces a small increase in strength in persons with muscular dystrophy [149]. Myostatin monoclonal antibodies increase muscle mass in mice [150]. In humans with muscular dystrophy, a myostatin antibody (MYO-029) enhanced muscle mass [151]. Muscle fiber diameter increased in the 10 mg/kg dose. Side effects included urticaria and aseptic meningitis at high doses. Another myostatin antibody (AMG 745) increased lean body mass and decreased fat after 28 days in persons on androgen deprivation therapy for prostate cancer [152]. Diarrhea, confusion, and fatigue were more common in the persons receiving active drug. LY2495655 increased muscle volume and handgrip strength in persons with advanced cancer (www.clinicaltrials.gov). REGN1033 (GDF8 antibody) has reported promising effects on muscle at the sarcopenia meeting in Barcelona.

An activin II receptor ligand trap (ACE-011) increased bone mass and strength in monkeys [153]. ACE-031 in 48 postmenopausal women increased lean mass and thigh muscle volume after a single dose [154]. Another ligand trap ACE-083 is also under development. Side effects including telangiectasia, epistaxis, and changes in gonadotrophin levels resulted in the company stopping the development of these compounds.

Table 4 Approaches currently available or being developed to treat sarcopenia

Modality	Effect	Side effects
Resistance exercise	Increase muscle mass, strength, and power	Potential for falls; muscle injuries
Protein (essential amino acids)	Increase muscle mass; synergy with exercise to increase muscle strength and power	Minimal increased creatinine levels
Testosterone	Increase muscle mass, strength, power, and function	Fluid retention; increased hematocrit; short term worsening of sleep apnea; effects on prostate cancer; possible increase in cardiovascular events
Selective androgen receptor modulators (SARMS)	Increase muscle mass; small increase in power	Increased cardiac failure
Growth hormone	Increase nitrogen retention; increase muscle mass	Arthralgia; muscle pain; edema; carpal tunnel syndrome; hyperglycemia
Ghrelin agonists	Increased muscle mass and appetite	Fatigue; atrial fibrillation; dyspnea
Myostatin antibodies	Increased lean body mass and handgrip	Urticaria; aseptic meningitis; diarrhea; confusion; fatigue
Activin IIR antagonists	Increase thigh muscle volume, muscle mass, and 6-min walk distance	Acne; involuntary muscle contractions
Angiotensin converting enzyme inhibitor (perindopril)	Increased distance walked; decreased hip fracture	Hypotension; hyperkalemia; muscle cramps; numbness
Espindolol (B ₁ /B ₂ adrenergic receptor antagonist)	Maintains muscle mass; increased hand grip strength	?
Fast skeletal muscle troponin activators (Tirasentiv)	Improves muscle function	?

Inclusion body myositis is a rare autoimmune disorder. It occurs in persons 50 years and older. Its hallmark is amyloid inclusion bodies. Bimagrab is an activin receptor inhibitor. In persons with inclusion body myositis, bimagrab increased thigh muscle volume, lean muscle mass, and 6-min walking distance [155].

Espindolol (Mixed Agonist/Antagonist B1,B2, B3 Activity)

Espindolol is the S-enantiomer of pindolol. It increases muscle mass and decreases fat mass in older animals [156]. A phase II trial showed an increase in muscle mass and a decrease in fat mass [157]: It also increased handgrip strength.

Angiotensin Converting Enzyme Inhibitor (Perindopril)

Perindopril has been shown to increase distance walked in older persons with left ventricular systolic dysfunction [158]. It also improved 6-min walking distance in older persons with functional impairment [159]. There was, however, no enhancement in persons undergoing exercise training [160]. In addition, in the HYVET study perindopril decreased hip fracture [161].

Fast Skeletal Troponin Activators (Terasemtiv)

There are drugs which amplify motor neuron input, resulting in improved muscle power and muscle fatigability. Terasemtiv slowed the rate of decline in muscle strength [162]. In persons with peripheral vascular disease, terasemtiv increased work done in a bilateral heel raising test (www.cytokinetics.com/ck2017357).

Conclusion

Sarcopenia (loss of muscle mass and muscle function) is a strong predictor of frailty, disability, and mortality in older persons. At present resistance exercise is the primary treatment for sarcopenia. Supplementation with essential amino acids, creatine, and vitamin D may enhance the effect of resistance exercise. The effects of testosterone on muscle include increased muscle power and function. At present, the side effects of testosterone, though minimal in placebo controlled trials, remain a possible limitation to its use. No other drugs under development have been shown to be more potent than testosterone. All the drugs under development have their own set of side-effects and will clearly be more expensive than resistance exercise or injectable testosterone (Table 4). There are numerous

potential targets for enhancing muscle function, and the development of new drugs for sarcopenia represents a potentially exciting clinical area.

Conflict of interest The author declares that there were no funds received from any entity for the writing of this article. The author has received grants from Nestle (Purina) and Kemin that are not related to this work. The author is on the speaker's bureau and has been a consultant for Nutricia and received consultant fees from Nutricia, Sanofi-Aventis, Astellas, Boehringer-Ingelheim, Merck, and Akros, none of which have any influence on the writing of this article.

References

- Rosenberg IH, Roubenoff R (1995) Stalking sarcopenia. *Ann Intern Med* 123:727–728
- Clark BC, Manini TM (2008) Sarcopenia \neq dynapenia. *J Gerontol A Biol Sci Med Sci* 63:829–834
- Barbat-Artigas S, Rolland Y, Vellas B, Aubertin-Leheudre M (2013) Muscle quantity is not synonymous with muscle quality. *J Am Med Dir Assoc* 14:852.e1-7. doi:10.1016/j.jamda.2013.06.003
- Manini TM, Clark BC (2012) Dynapenia and aging: an update. *J Gerontol A* 67:28–40
- Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel JP, Rolland Y et al (2010) Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on sarcopenia in older people. *Age Ageing* 39:412–423
- Landi F, Cruz-Jentoft AJ, Liperoti R, Russo A, Giovannini S, Tosato M, Capoluongo E, Bernabei R, Onder G (2013) Sarcopenia and mortality risk in frail older persons aged 80 years and older: results from the ILSIRENTE study. *Age Ageing* 42:203–209
- Landi F, Liperoti R, Fusco D, Mastropalo S, Quattrocioni D, Proia A, Tosato M, Bernabei R, Onder G (2012) Sarcopenia and mortality among older nursing home residents. *J Am Med Dir Assoc* 13:121–126
- Lee WJ, Liu LK, Peng LN, Lin MH, Chen LK, ILAS Research Group (2013) Comparisons of sarcopenia defined by IWGS and EWGSOP criteria among older people: results from the I-Lan longitudinal aging study. *J Am Med Dir Assoc* 14:528.e1-7
- Fielding RA, Vellas B, Evans WJ, Bhasin S, Morley JE, Newman AB, Abellan van Kan G et al (2011) Sarcopenia: An undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International Working Group on Sarcopenia. *J Am Med Dir Assoc* 12:249–256
- Morley JE, Abbatecola AM, Argiles JM, Baracos V, Bauer J, Bhasin S, Cederholm T, Coats AJ, Cummings SR, Society on Sarcopenia, Cachexia and Wasting Disorders Trialist Workshop et al (2011) Sarcopenia with limited mobility: An international consensus. *J Am Med Dir Assoc* 12:403–409
- Dam TT, Peters KW, Fragala M, Cawthon PM, Harris TB, McLean R, Shardell M, Alley DE, Kenny A, Ferrucci L, Guralnik J, Kiel DP, Kritchevsky S, Vassileva MT, Studenski S (2014) An evidence-based comparison of operational criteria for the presence of sarcopenia. *J Gerontol A* 69:584–590
- Chen LK, Liu LK, Woo J, Assantachai P, Auyeung TW, Bahyah KS, Chou MY, Chen LY, Hsu PS, Krairit O, Lee JS, Lee WJ, Lee Y, Ck Liang, Limpawattana P, Lin CS, Peng LN, Satake S,

- Suzuki T, Won CW, Wu CH, Wu SN, Zhang T, Zeng P, Akishita M, Arai H (2014) Sarcopenia in Asia: consensus report of the Asian Working Group for Sarcopenia. *J Am Med Dir Assoc* 15:95–101
13. Woo J, Leung J, Morley JE (2014) Defining sarcopenia in terms of incident adverse outcomes. *J Am Med Dir.* doi:[10.1016/j.jamda.2014.11.013](https://doi.org/10.1016/j.jamda.2014.11.013)
 14. Kanis JA, McCloskey E, Johansson H, Oden A, Leslie WD (2012) FRAX[®] with and without bone mineral density. *Calcif Tissue Int* 90:1–13
 15. Malmstrom TK, Morley JE (2013) SARC-F: a simple questionnaire to rapidly diagnose sarcopenia. *J Am Med Dir Assoc* 14:531–532
 16. Woo J, Leung J, Morley JE (2014) Validating the SARC-F: a suitable community screening tool for sarcopenia? *J Am Med Dir Assoc* 15:630–634
 17. Cao L, Chen S, Zou C, Ding X, Gao L, Liao Z, Liu G, Malmstrom TK, Morley JE, Flaherty JH, An Y, Dong B (2014) A pilot study of the SARC-F scale on screening sarcopenia and physical disability in the Chinese older people. *J Nutr Health Aging* 18:277–283
 18. Malmstrom TK, Simonsick EM, Ferrucci L, Miller DK, Morley JE (2015) SARC-F: a symptom score to predict persons with sarcopenia at risk for poor functional outcomes. *J Cachexia Sarc Muscle.* doi:[10.1002/jcsm.12048](https://doi.org/10.1002/jcsm.12048)
 19. Argiles JM, Anker SD, Evans WJ, Morley JE, Fearon KC, Strasser F, Muscaritoli M, Baracos VE (2010) Consensus on cachexia definitions. *J Am Med Dir Assoc* 11:229–230
 20. Purves-Smith FM, Sgarioto N, Hepple RT (2014) Fiber typing in aging muscle. *Exerc Sport Sci Rev* 42:45–52
 21. Hepple RT (2003) Sarcopenia: a critical perspective. *Sci Aging Knowl Environ* 45:31
 22. Drey M, Grosch C, Neuwirth C, Bauer JM, Sieber CC (2013) The motor unit number index (MUNIX) in sarcopenic patients. *Exp Gerontol* 48:381–384
 23. Hettwer S, Dahinden P, Kucsera S, Farina C, Ahmed S, Fariello R et al (2013) Elevated levels of a C-terminal agrin fragment identifies a new subset of sarcopenia patients. *Exp Gerontol* 48:69–75
 24. Drey M, Krieger B, Sieber CC, Bauer JM, Hettwer S, Bertsch T, DISARCO Group (2014) Motoneuron loss is associated with sarcopenia. *J Am Med Dir Assoc* 15:435–439
 25. McNeil CJ, Doherty TJ, Stashuk DW, Rice CL (2005) Motor unit number estimates in the tibialis anterior muscle of young, old, and very old men. *Muscle Nerve* 31:461–467
 26. Guillet C, Auguste P, Mayo W, Kreher P, Gascan H (1999) Ciliary neurotrophic factor is a regulator of muscular strength in aging. *J Neurosci* 19:1257–1262
 27. Arking DE, Fallin DM, Fried LP, Li T, Beamer BA, Xue QL, Chakravarti A, Walston J (2006) Variation in the ciliary neurotrophic factor gene and muscle strength in older Caucasian women. *J Am Geriatr Soc* 54:823–826
 28. Demontis F, Piccirillo R, Goldberg AL, Perrimon N (2013) The influence of skeletal muscle on systemic aging and lifespan. *Aging Cell* 12:943–949
 29. Iizuka K, Machida T, Hirafuji M (2014) Skeletal muscle is an endocrine organ. *J Pharmacol Sci* 125:125–131
 30. Yoon JH, Kim J, Song P, Lee TG, Suh PG, Ryu SH (2012) Secretomics for skeletal muscle cells: a discovery of novel regulators? *Adv Biol Regul* 52:340–350
 31. Pal M, Febbraio MA, Whitham M (2014) From cytokine to myokine: The emerging role of interleukin-6 in metabolic regulation. *Immunol Cell Biol* 92:331–339
 32. Pistilli EE, Quinn LS (2013) From anabolic to oxidative: reconsidering the roles of IL-15 and IL-15R α in skeletal muscle. *Exerc Sport Sci Rev* 41:100–106
 33. Morley JE, von Haehling S, Anker SD, Vellas B (2014) From sarcopenia to frailty: a road less travelled. *J Cachexia Sarcopenia Muscle* 5:5–8
 34. Morley JE, Vellas B, van Kan GA, Anker SD, Bauer JM, Bernabei R, Cesari M, Chumlea WC, Doehner W, Evans J et al (2013) Frailty consensus: a call to action. *J Am Med Dir Assoc* 14:392–397
 35. Garatachea N, Lucia A (2013) Genes and the ageing muscle: a review on genetic association studies. *Age* 35:207–233
 36. Stewart CEH, Rittweger J (2006) Adaptive processes in skeletal muscle: molecular regulators and genetic influences. *J Musculoskelet Neuronal Interact* 6:783–786
 37. Seto JT, Quinlan KG, Lek M, Zheng XF, Garton F, MacArthur DG et al (2013) ACTN3 genotype influences muscle performance through the regulation of calcineurin signaling. *J Clin Invest* 123:4255–4263
 38. Seto JT, Chan S, Turner N, MacArthur DG, Raftery JM, Berman YD et al (2011) The effect of α -actinin-3 deficiency on muscle aging. *Exp Gerontol* 46:292–302
 39. Dennis RA, Zhu H, Kortebein PM, Bush HM, Harvey JF, Sullivan DH, Peterson CA (2009) Muscle expression of genes associated with inflammation, growth, and remodeling is strongly correlated in older adults with resistance training outcomes. *Physiol Genomics* 38:169–175
 40. Windelinckx A, De Mars G, Huygens W, Peeters MW, Vincent B, Wijmenga C, Lambrechts D et al (2011) Comprehensive fine mapping of chr12q12-14 and follow-up replication identify activin receptor 1B (ACVR1B) as a muscle strength gene. *Eur J Hum Genet* 19:208–215
 41. Conte M, Vasuri F, Trisolino G, Bellavista E, Santoro A, Degiovanni A et al (2013) Increased Plin2 expression in human skeletal muscle is associated with sarcopenia and muscle weakness. *PLoS ONE* 8(8):e73709
 42. Rolland Y, Lauwers-Cances V, Crostini C, Abellan van Kan G, Janssen I, Morley JE, Vellas B (2009) Difficulties with physical function associated with obesity, sarcopenia, and sarcopenic-obesity in community-dwelling elderly women. *EPIDOS Study* 89:1895–1900
 43. Baumgartner RN, Wayne SJ, Waters DL, Janssen I, Gallagher D, Morley JE (2004) Sarcopenic obesity predicts instrumental activities of daily living disability in the elderly. *Obes Res* 12:1995–2004
 44. Arnold A-S, Egger A, Handschin C (2011) PGC-1 α and myokines in the aging muscle—a mini-review. *Gerontology* 57:37–43
 45. Ling C, Poulsen P, Carlsson E, Ridderstrale M, Almgren P, Wojtaszewski J, Beck-Nielsen H, Groop L, Vaag A (2004) Multiple environmental and genetic factors influence skeletal muscle PGC-1 α and PGC-1 β gene expression in twins. *J Clin Invest* 114:1518–1526
 46. Sandri M, Lin J, Handschin C, Yang W, Zp Arany, Lecker SH, Goldberg AL, Spiegelman BM (2006) PGC-1 α protects skeletal muscle from atrophy by suppressing foxO3 action and atrophy-specific gene transcription. *Proc Natl Acad Sci USA* 103:16260–16265
 47. Hepple RT (2014) Mitochondrial involvement and impact in aging skeletal muscle. *Front Aging Neurosci* 6:211. doi:[10.3389/fnagi.2014.00211](https://doi.org/10.3389/fnagi.2014.00211)
 48. Marzetti E, Calvani R, Cesari M, Buford TW, Lorenzi M, Behnke BJ, Leeuwenburgh C (2013) Mitochondrial dysfunction and sarcopenia of aging: From signaling pathways to clinical trials. *Int J Biochem Cell Biol* 45:2288–2301
 49. Coen PM, Jubrias SA, Distefano G, Amati F, Mackey DC, Glynn NW et al (2013) Skeletal muscle mitochondrial energetics are associated with maximal aerobic capacity and walking speed in older adults. *J Gerontol A* 68:447–455

50. Wang CH, Kao CH, Chen YF, Wei YH, Tsai TF (2014) Cisd2 mediates lifespan: Is there an interconnection among Ca²⁺ homeostasis, autophagy, and lifespan? *Free Radic Res* 48: 1109–1114
51. Wu CY, Chen YF, Wang CH, Kao CH, Zhuang HW, Chen CC, Chen LK et al (2012) A persistent level of Cisd2 extends healthy lifespan and delays aging in mice. *Hum Mol Genet* 21: 3956–3968
52. Lawrenson L, Poole JG, Kim J, Brown C, Patel P, Richardson RS (2003) Vascular and metabolic response to isolated small muscle mass exercise: effect of age. *Am J Physiol Heart Circ Physiol* 285:H1023–H1031. doi:10.1152/ajpheart.00135.2003
53. Donato AJ, Uberoi A, Wray DW, Nishiyama S, Lawrenson L, Richardson RS (2006) Differential effects of aging on limb blood flow in humans. *Am J Physiol Heart Circ Physiol* 290:H272–H278
54. Seals DR, Jablonski KL, Donato AJ (2011) Aging and vascular endothelial function in humans. *Clin Sci* 120:357–375
55. Behnke BJ, Delp MD, Dougherty PJ, Musch TI, Poole DC (2005) Effects of aging on microvascular oxygen pressures in rat skeletal muscle. *Respir Physiol Neurobiol* 146:259–268
56. Cohen S, Nathan JA, Goldberg AL (2015) Muscle wasting in disease: molecular mechanisms and promising therapies. *Nature* 14:58–74
57. Cohen S, Brault JJ, Gygi SP, Glass DJ, Valenzuela DM, Geratner C, Latres E, Goldberg AL (2009) During muscle atrophy, thick, but not thin, filament components are degraded by MuRF1-dependent ubiquitylation. *J Cell Biol* 185:1083–1095
58. Cohen S, Zhai B, Gygi SP, Goldberg AL (2012) Ubiquitylation by Trim32 causes coupled loss of desmin, Z-bands, and thin filaments in muscle atrophy. *J Cell Biol* 198:575–589
59. Verdijk LB, Koopmans R, Schaart G, Meijer K, Savelberg HHCM, van Loon LJC (2007) Satellite cell content is specifically reduced in type II skeletal muscle fibers in the elderly. *Am J Physiol Endocrinol Metab* 292:E151–E157
60. McGregoe RA, Poppitt SD, Cameron-Smith D (2014) Role of microRNAs in the age-related changes in skeletal muscle and diet or exercise interventions to promote healthy aging in humans. *Ageing Res Rev* 17:25–33
61. Rivas DA, Lessard SJ, Rice NP, Lustgarten MS, So K, Goodyear LJ, Parnell LD, Fielding RA (2014) Diminished skeletal muscle microRNA expression with aging is associated with attenuated muscle plasticity and inhibition of IGF-1 signaling. *FASEB J* 28:4133–4147
62. Barbierato M, Zusso M, Skaper SD, Giusti P (2015) MicroRNAs: emerging role in the endogenous mu opioid system. *CNS Neurol Disord Drug Targets* 14:239–250
63. Alieva AKH, Filatova EV, Karabonov AV, Illarioshkin SN, Limborska SA, Shadrina MI, Slominsky PA (2015) miRNA expression is highly sensitive to a drug therapy in Parkinson's disease. *Parkinsonism Relat Disord* 21:72–74
64. Kern H, Barberi L, Löffler S, Sbardella S, Burggraf S, Fruhmann H et al (2014) Electrical stimulation counteracts muscle decline in seniors. *Front Aging Neurosci*. doi:10.3389/fnagi.2014.00189
65. Conboy IM, Conboy MJ, Wagers AJ, Girma ER, Weissman IL, Rando TA (2005) Rejuvenation of aged progenitor cells by exposure to a young systemic environment. *Nature* 433(7027):760–764
66. Sinha M, Jang YC, Oh J, Khong D, Wu EY, Manohar R, Miller C et al (2014) Restoring systemic GDF11 levels reverses age-related dysfunction in mouse skeletal muscle. *Science* 344(6184):649–652
67. Sinha I, Sinha-Hikim AP, Wagers AJ, Sinha-Hikim I (2014) Testosterone is essential for skeletal muscle growth in aged mice in a heterochronic parabiosis model. *Cell Tissue Res* 357(3):815–821
68. Fuster G, Busquets S, Almendro V, Lopez-Soriano FJ, Argiles JM (2007) Antiproteolytic effects of plasma from hibernating bears: a new approach for muscle wasting therapy? *Clin Nutr* 26:658–661
69. Bodine SC (2013) Hibernation: The search for treatments to prevent disuse-induced skeletal muscle atrophy. *Exper Neurol* 248:129–135
70. Andres-Mateos E, Brinkmeier H, Burks TN, Mejias R, Files DC, Steinberger M, Soleimani A et al (2013) Activation of serum/glucocorticoid-induced kinase 1 (SGK1) is important to maintain skeletal muscle homeostasis and prevent atrophy. *EMBO Mol Med* 5:80–91
71. Valenzuela T (2012) Efficacy of progressive resistance training interventions in older adults in nursing homes: a systematic review. *J Am Med Dir Assoc* 13:418–428
72. Cadone Cadore EL, Izquierdo M (2013) New strategies for the concurrent strength-, power-, and endurance-training prescription in elderly individuals. *J Am Med Dir Assoc* 14:623–624
73. Singh NA, Quine S, Clemson LM, Williams EJ, Williamson DA, Stavrinou TM et al (2012) Effects of high-intensity progressive resistance training and targeted multidisciplinary treatment of frailty on mortality and nursing home admissions after hip fracture: a randomized controlled trial. *J Am Med Dir Assoc* 13:24–30
74. Pahor M, Guralnik JM, Ambrosius WT, Blair S, Bonds DE, Church TS et al (2014) Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *JAMA* 311(23):2387–2396
75. Yamada M, Arai H, Sonoda T, Aoyama T (2012) Community-based exercise program is cost-effective by preventing care and disability in Japanese frail older adults. *J Am Med Dir Assoc* 13:507–511
76. Bauer J, Biolo G, Cederholm T, Cesari M, Cruz-Jentoft AJ, Morley JE et al (2013) Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE study group. *J Am Med Dir Assoc* 14:542–559
77. Malafarina V, Uriz-Otano F, Iniesta R, Gil-Guerrero L (2013) Effectiveness of nutritional supplementation on muscle mass in treatment of sarcopenia in old age: a systematic review. *J Am Med Dir Assoc* 14:10–17
78. Boirie Y (2013) Fighting sarcopenia in older frail subjects: protein fuel for strength, exercise for mass. *J Am Med Dir Assoc* 14:140–143
79. Tieland M, van de Rest O, Dirks ML, van der Zwaluw N, Mensink M, van Loon LJ et al (2012) Protein supplementation improves physical performance in frail elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 13:720–726
80. Cermak NM, de Groot LC, van Loon LJ (2013) Perspective: Protein supplementation during prolonged resistance type exercise training augments skeletal muscle mass and strength gains. *J Am Med Dir Assoc* 14:71–72
81. Gryson C, Ratel S, Rance M, Penando S, Bonhomme C, Le Ruyet P, Duclos M, Boirie Y, Walrand S (2014) Four-month course of soluble milk proteins interacts with exercise to improve muscle strength and delay fatigue in elderly participants. *J Am Med Dir Assoc* 15:958.e1-9
82. Ferrando AA, Paddon-Jones D, Hays NP, Kortebein P, Ronsen O, Williams RH, McComb A, Symons TB, Wolfe RR, Evans W (2010) EAA supplementation to increase nitrogen intake improves muscle function during bed rest in the elderly. *Clin Nutr* 29:18–23
83. Bauer JM, Verlaan S, Bautmans I, Brandt K, Donini LM, Maggio M, et al. (2015) Effects of a vitamin D and leucine-enriched whey protein nutritional supplement on measures of

- sarcopenia in older adults. The PROVIDE study: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir* (in press)
84. Morley JE, Argiles JM, Evans WJ, Bhasin S, Cella D, Deutz NE et al (2010) Nutritional recommendations for the management of sarcopenia. *J Am Med Dir Assoc* 11:391–396
 85. Van Wetering C, Hoogendoorn M, Broekhuizen R, Geraerts-Keeris GJ, de Munck DR, Rutten-van Molken MP et al (2010) Efficacy and costs of nutritional rehabilitation in muscle-wasted patients with chronic obstructive pulmonary disease in a community-based setting: a prespecified subgroup analysis of the INTERCOM trial. *J Am Med Dir Assoc* 11:179–187
 86. Tieland M, Dirks ML, van der Zwaluw N, Verdijk LB, van de Rest O, de Groot LC, van Loon LJ (2012) Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 13:713–719
 87. Beaudart C, Buckinx F, Rabenda V, Gillain S, Cavalier E, Slomian J et al (2014) The effects of vitamin D on skeletal muscle strength, muscle mass, and muscle power: a systematic review and meta-analysis of randomized controlled trials. *J Clin Endocrinol Metab* 99:4336–4345
 88. Murad MH, Elamin KB, Abu Elnour NO, Elamin MB, Alkhatib AA, Fatourehchi MM et al (2011) Clinical review: the effect of vitamin D on falls: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2011(96):2997–3006
 89. Morley JE (2011) Should frailty be treated with testosterone? *Aging Male* 14:1–3
 90. Morley JE, Fe Kaiser, Perry HM 3rd, Patrick P, Morley PM, Stauber PM, Vellas B, Baumgartner RN, Garry PJ (1997) Longitudinal changes in testosterone, luteinizing hormone, and follicle-stimulating hormone in healthy older men. *Metabolism* 46:410–413
 91. Baumgartner RN, Di Waters, Gallagher D, Morley JE, Garry PJ (1999) Predictors of skeletal muscle mass in elderly men and women. *Mech Ageing Dev* 107:123–136
 92. Morley JE, Perry HM 3rd, Kaiser FE, Kraenzle D, Jensen J, Houston K, Mattammal M, Perry HM Jr (1993) Effects of testosterone replacement therapy in old hypogonadal males: a preliminary study. *J Am Geriatr Soc* 41:149–152
 93. Page ST, Amory JK, Bowman FD, Anawalt BD, Matsumoto AM, Bremner WJ et al (2005) Exogenous testosterone (T) alone or with finasteride increases physical performance, grip strength, and lean body mass in older men with low serum T. *J Clin Endocrinol Metab* 90:1502–1510
 94. Sih R, Morley JE, Kaiser FE, Perry HM 3rd, Patrick P, Ross C (1997) Testosterone replacement in older hypogonadal men: A 12-month randomized controlled trial. *J Clin Endocrinol Metab* 82:1661–1667
 95. Wittert GA, Chapman IM, Haren MT, Mackintosh S, Coates P, Morley JE (2003) Oral testosterone supplementation increases muscle and decreases fat mass in healthy elderly males with low-normal gonadal status. *J Gerontol A* 58:618–625
 96. Bhasin S, Woodhouse L, Casaburi R, Singh AB, Mac RP, Lee M, Yarasheski KE et al (2005) Older men are as responsive as young men to the anabolic effects of graded doses of testosterone on the skeletal muscle. *J Clin Endocrinol Metab* 90:678–688
 97. Srinivas-Shankar U, Roberts SA, Connolly MJ, O'Connell MD, Adams JE, Oldham JA et al (2010) Effects of testosterone on muscle strength, physical function, body composition, and quality of life in intermediate-frail and frail elderly men: a randomized, double-blind, placebo-controlled study. *J Clin Endocrinol Metab* 95:639–650
 98. Kenny AM, Kleppinger A, Annis K, Rathier M, Browner B, Judge JO et al (2010) Effects of transdermal testosterone on bone and muscle in older men with low bioavailable testosterone levels, low bone mass, and physical frailty. *J Am Geriatr Soc* 58:1134–1143
 99. Travison TG, Basaria S, Storer TW, Jette AM, Miciek R, Farwell WR et al (2011) Clinical meaningfulness of the changes in muscle performance and physical function associated with testosterone administration in older men with mobility limitation. *J Gerontol A Biol Sci Med Sci* 66:1090–1099
 100. Caminiti G, Volterrani M, Iellamo F, Marazzi G, Massaro R, Miceli M et al (2009) Effect of long-acting testosterone treatment on functional exercise capacity, skeletal muscle performance, insulin resistance, and baroreflex sensitivity in elderly patients with chronic heart failure: a double-blind, placebo-controlled, randomized study. *J Am Coll Cardiol* 54:919–927
 101. Iellamo F, Volterrano M, Caminiti G, Karam R, Massaro R, Fini M et al (2010) Testosterone therapy in women with chronic heart failure: a pilot double-blind, randomized, placebo-controlled study. *J Am Coll Cardiol* 56:1310–1316
 102. Malkin CJ, Pugh PJ, West JN, van Beek EJ, Jones TH, Channer KS (2006) Testosterone therapy in men with moderate severity heart failure: a double-blind randomized placebo controlled trial. *Eur Heart J* 27:57–64
 103. Stout M, Tew GA, Doll H, Zwierska I, Woodroffe N, Channer KS, Saxton JM (2012) Testosterone therapy during exercise rehabilitation in male patients with chronic heart failure who have low testosterone status: a double-blind randomized controlled feasibility study. *Am Heart J* 164:893–901
 104. Morley JE, Perry HM 3rd (2003) Androgens and women at the menopause and beyond. *J Gerontol A Biol Sci Med Sci* 58:M409–M416
 105. Chapman IM, Visvanathan R, Hammond AJ, Morley JE, Field JB, Tai K et al (2009) Effect of testosterone and a nutritional supplement, alone and in combination, on hospital admissions in undernourished older men and women. *Am J Clin Nutr* 89:880–889
 106. Wolfe R, Ferrando A, Sheffield-Moore M, Urban R (2000) Testosterone and muscle protein metabolism. *Mayo Clin Proc* 75(Suppl):S55–S59
 107. Ferrando AA, Sheffield-Moore M, Paddon-Jones D, Wolfe RR, Urban RJ (2003) Differential anabolic effects of testosterone and amino acid feeding in older men. *J Clin Endocrinol Metab* 88:358–362
 108. Kovacheva EL, Hikim AP, Shen R, Sinha I, Sinha-Hikim I (2010) Testosterone supplementation reverses sarcopenia in aging through regulation of myostatin, c-Jun NH2-terminal kinase, Notch, and Akt signaling pathways. *Endocrinology* 151:628–638
 109. Harin MT, Siddiqui AM, Armbrrecht HJ, Kevorkian RT, Kim MJ, Haas MJ, Mazza A, Kumar VB, Green M, Banks WA, Morley JE (2011) Testosterone modulates gene expression pathways regulating nutrient accumulation, glucose metabolism and protein turnover in mouse skeletal muscle. *Int J Androl* 34:55–68
 110. Morley JE (2013) Scientific overview of hormone treatment used for rejuvenation. *Fertil Steril* 99:1807–1813
 111. Carona G, Maseroli E, Rastrelli G, Isidori AM, Sforza A, Mannucci E, Maggi M (2014) Cardiovascular risk associated with testosterone-boosting medications: a systematic review and meta-analysis. *Expert Opin Drug Saf* 13:1327–1351
 112. Mogentaler A, Miner MM, Caliber M, Guay AT, Khera M, Traish AM (2015) Testosterone therapy and cardiovascular risk: advances and controversies. *Mayo Clin Proc* 90:224–251
 113. Borst SE, Shuster JJ, Zou B, Ye F, Jia H, Wokhlu A, Yarrow JF (2014) Cardiovascular risks and elevation of serum DHT vary by route of testosterone administration: a systematic review and meta-analysis. *BMC Med* 12:211–215

114. Morley JE, Malmstrom TK, Rodriguez-Manas L, Sinclair AJ (2014) Frailty, sarcopenia and diabetes. *J Am Med Dir Assoc* 15:853–859
115. Landi F, Onder G, Bernabei R (2013) Sarcopenia and diabetes: two sides of the same coin. *J Am Med Dir Assoc* 14:540–541
116. Leenders M, Verdijk LB, van der Hoeven L, Adam JJ, van Kranenburg J, Nilwik R, van Loon LJ (2013) Patients with type 2 diabetes show a greater decline in muscle mass, muscle strength, and functional capacity with aging. *J Am Med Dir Assoc* 14:585–592
117. Muraleedharan V, March H, Kapoor D, Channer KS, Jones TH (2013) Testosterone deficiency is associated with increased risk of mortality and testosterone replacement improves survival in men with type 2 diabetes. *Eur J Endocrinol* 169:725–733
118. Nieschlag E (2015) Current topics in testosterone replacement of hypogonadal men. *Best Pract Res Clin Endocrinol Metab* 29:77–90
119. Irwig MS (2014) Bone health in hypogonadal men. *Curr Opin Urol* 24:608–613
120. Sharma S, Arneja A, McLean L, Duerksen D, Leslie W, Sciberras D, Lertzman M (2008) Anabolic steroids in COPD: a review and preliminary results of a randomized trial. *Chron Respir Dis* 5:169–176
121. Frisoli A Jr, Chaves PH, Pinheiro MM, Vi Szejnfeld (2005) The effect of nandrolone decanoate on bone mineral density, muscle mass, and hemoglobin levels in elderly women with osteoporosis: a double-blind, randomized, placebo-controlled clinical trial. *J Gerontol A* 60:648–653
122. Macdonald JH, Marcora SM, Jibani MM, Kumwenda MJ, Ahmed W, Lemmey AB (2007) Nandrolone decanoate as anabolic therapy in chronic kidney disease: a randomized phase II dose-finding study. *Nephron Clin Pract* 106:c125–c135
123. Farooqi V, van den Berg ME, Cameron ID, Crotty M (2014) Anabolic steroids for rehabilitation after hip fracture in older people. *Cochrane Database Syst Rev* 10:CD008887
124. Papanicolaou DA, Ather SN, Zhu H, Zhou Y, Lutkiewicz J, Scott BB, Chandler J (2013) A phase IIA randomized, placebo-controlled clinical trial to study the efficacy and safety of the selective androgen receptor modulator (SARM), MK-0773 in female participants with sarcopenia. *J Nutr Health Aging* 17:533–543
125. Mohler ML, Bohl CE, Jones A, Coss CC, Narayanan R, He Y et al (2009) Nonsteroidal selective androgen receptor modulators (SARMs): dissociating the anabolic and androgenic activities of the androgen receptor for therapeutic benefit. *J Med Chem* 52:3598–3617
126. Bhasin S, Jasuja R (2009) Selective androgen receptor modulators as function promoting therapies. *Curr Opin Clin Nutr Metab Care* 12:232–240
127. Basario S, Collins L, Dillon EL, Orwoll K, Storer TW, Micek R et al (2013) The safety, pharmacokinetics, and effects of LGD-4033, a novel nonsteroidal oral, selective androgen receptor modulator, in healthy young men. *J Gerontol A* 68:87–95
128. Dalton JT, Barnette KG, Bohl CE, Hancock ML, Rodriguez D, Dodson ST et al (2011) The selective androgen receptor modulator GTx-024 (enobosarm) improves lean body mass and physical function in healthy elderly men and postmenopausal women: results of a double-blind, placebo-controlled phase II trial. *J Cachexia Sarcopenia Muscle* 2:153–161
129. Dobs AS, Boccia RV, Croot CC, Gabriel NY, Dalton JT, Hancock ML et al (2013) Effects of enobosarm on muscle wasting and physical function in patients with cancer: a double-blind, randomized controlled phase 2 trial. *Lancet Oncol* 14:335–345
130. Steiner MS (2013) Enobosarm, a selective androgen receptor modulator, increases lean body mass in advanced non-small cell lung cancer patients in two pivotal, international phase 3 trials. *J Cachexia Sarcopenia Muscle* 4:69. doi:10.1007/s13539-013-0123-9
131. Rudman D, Feller AG, Nagraj HS, Gergans GA, Lalitha PY, Goldberg AF et al (1990) Effects of human growth hormone in men over 60 years old. *N Engl J Med* 323:1–6
132. Cohn L, Feller AG, Draper MW, Rudman IW, Rudman D (1993) Carpal tunnel syndrome and gynaecomastia during growth hormone treatment of elderly men with low circulating IGF-1 concentrations. *Clin Endocrinol (Oxf)* 39:417–425
133. Kim MJ, Morley JE (2005) The hormonal fountains of youth: myth or reality? *J Endocrinol Invest* 28(11):5–14
134. Blackman MR, Sorkin JD, Munzer T, Bellantoni MF, Busby-Whitehead J, Stevens TE et al (2002) Growth hormone and sex steroid administration in healthy aged women and men: a randomized controlled trial. *JAMA* 288:2282–2292
135. Kaiser FE, Silver AJ, Morley JE (1991) The effect of recombinant human growth hormone on malnourished older individuals. *J Am Geriatr Soc* 39:235–240
136. Liu H, Bravata DM, Olkin I, Nayak S, Roberts B, Am Garber, Hoffman AR (2007) Systematic review: the safety and efficacy of growth hormone in the healthy elderly. *Ann Intern Med* 146:104–115
137. Giannoulis MG, Martin FC, Nair KS, Umpleby AM, Sonksen P (2012) Hormone replacement therapy and physical function in healthy older men. Time to talk hormones? *Endocr Rev* 33:314–377
138. Sullivan DH, Carter WJ, Warr WR, Williams LH (1998) Side effects resulting from the use of growth hormone and insulin-like growth factor-1 as combined therapy to frail elderly patients. *J Gerontol A* 53:M183–M187
139. Gaskin FS, Farr SA, Banks WA, Kumar VB, Morley JE (2008) Ghrelin-induced feeding is dependent on nitric oxide. *Peptides* 24:913–918
140. Argiles JM, Stemmler B (2013) The potential of ghrelin in the treatment of cancer cachexia. *Expert Opin Biol Ther* 13:67–76
141. Miyazaki T, Tanaka N, Hirai H, Yokobori T, Sano A, Sakai M et al (2012) Ghrelin level and body weight loss after esophagectomy for esophageal cancer. *J Surg Res* 176:74–78
142. Garcia JM, Boccia RV, Graham CD, Yan Y, Duus EM, Allen S, Friend J (2015) Anamorelin for patients with cancer cachexia: an integrated analysis of two phase 2, randomized, placebo-controlled, double-blind trials. *Lancet Oncol* 16:108–116
143. Temel J, Bondarde S, Jain M, Yun Y, Duus E, Allen S et al (2013) Efficacy and safety results from a phase II study of anamorelin HC21, a ghrelin receptor agonist, in NSCLC patients. *J Cachexia Sarcopenia Muscle* 4:334
144. White HK, Petrie CD, Landschulz W, MacLean D, Taylor A, Lyles K et al (2009) Effects of an oral growth hormone secretagogue in older adults. *J Clin Endocrinol Metab* 94:1198–1206
145. Adunsky A, Chandler J, Heyden N, Lutkiewicz J, Scott BB, Berd Y et al (2011) MK-0677 (ibutamoren mesylate) for the treatment of patients recovering from hip fractures: a multicenter, randomized, placebo-controlled phase IIb study. *Arch Gerontol Geriatr* 53:183–189
146. Elkina Y, vono Haehling S, Anker SD, Springer J (2011) The role of myostatin in muscle wasting: an overview. *J Cachexia Sarcopenia Muscle* 2:143–151
147. Schuelke M, Wagner KR, Stolz LE, Hubner C, Riebel T, Komen W, Braun T, Tobin JF, Lee SJ (2004) Myostatin mutation associated with gross muscle hypertrophy in a child. *N Engl J Med* 350:2682–2688
148. Saremi A, Gharakhanloo R, Sharghi S, Gharraati MR, Larijani B, Omidfar K (2010) Effects of oral creatine and resistance training on serum myostatin and GASP-1. *Mol Cell Endocrinol* 317:25–30

149. Walter MC, Reilich P, Lochmuller H, Kohnen R, Schlotter B, Hautmann H, Dunkl E et al (2002) Creatine monohydrate in myotonic dystrophy: a double-blind, placebo-controlled clinical study. *J Neurol* 249:1717–1722
150. Tsuchida K (2008) Targeting myostatin for therapies against muscle-wasting disorders. *Curr Opin Drug Discov Dev* 11:487–494
151. Wagner KR, Fleckenstein JL, Amato AA, Barhn RJ, Bushby K, Eagle M, Florence JM, King WM et al (2008) A phase I/II Trial of MYO-029 in adult subjects with muscular dystrophy. *Ann Neurol* 63:561–571
152. Padhi D, Higano CS, Shore ND, Sieber P, Rasmussen E, Smith MR (2014) Pharmacological inhibition of myostatin and changes in lean body mass and lower extremity muscle size in patients receiving androgen deprivation therapy for prostate cancer. *J Clin Endocrinol Metab* 99:E1967–E1975
153. Fajardo RJ, Manoharan RK, Pearsall RS, Davies MV, Marvell T, Monnell TE et al (2010) Treatment with a soluble receptor for activin improves bone mass and structure in the axial and appendicular skeleton of female cynomolgus macaques (*Macaca fascicularis*). *Bone* 46:64–71
154. Attie KM, Brogstein NG, Yang Y, Condon CH, Wilson DM, Pearsall AE et al (2013) A single ascending-dose study of muscle regulator ACE-031 in healthy volunteers. *Muscle Nerve* 47:416–423
155. Amato AA, Sivakumar K, Goyal N, David WS, Salajegheh M, Praestgaard J, Lach-Trifileff E, Trendelenburg AU, Laurent D, Glass DJ, Roubenoff R, Tseng BS, Greenberg SA (2014) Treatment of Sporadic inclusion body myositis with bimagrumab. *Neurology* 83:2239–2246
156. Potsch MS, Tschirner A, Palus S, von Haehling S, Doehner W, Beadle J, Coats AJ, Anker SD, Springer J (2014) The anabolic catabolic transforming agenda (ACTA) espidolol increases muscle mass and decreases fat mass in old rats. *J Cachexia Sarcopenia Muscle* 5:149–158
157. Steward Coats AJ, Srinivasan Surendran J, Chicarmana H, Vangipuram SR, Bhatt NN, Jain M et al (2011) The ACT-ONE trial, a multicenter, randomized, double-blind, placebo-controlled dose-finding study of the anabolic/catabolic transforming agent, MT-102 in subjects with cachexia related to stage III and IV non-small cell lung cancer and colorectal cancer: study design. *J Cachexia Sarcopenia Muscle* 2:201–207
158. Hutcheon SD, Gillespie ND, Crombie IK, Struthers AD, McMurdo ME (2002) Perindopril improves six minute walking distance in older patients with left ventricular systolic dysfunction: a randomized double blind placebo controlled trial. *Heart* 88:373–377
159. Sumukadas D, Witham MD, Struthers AD, McMurdo ME (2007) Effect of perindopril on physical function in elderly people with functional impairment: a randomized controlled trial. *CMAJ* 177:867–874
160. Sumukadas D, Band M, Miller S, Cvorov V, Witham M, Struthers A, McConnachie A, Lloud SM, McMurdo M (2014) Do ACE inhibitors improve the response to exercise training in functionally impaired older adults? A randomized controlled trial. *J Gerontol A* 69:736–743
161. Peters R, Beckett N, Burch L, de Vernejoul MC, Liu L, Duggan J et al (2010) The effect of treatment based on diuretic (indapamide) ± ACE inhibitor (perindopril) on fractures in the hypertension in the very elderly trial (HYVET). *Age Ageing* 39:609–616
162. Malik F (2013) Fast skeletal muscle troponin activators and their application to disease-preclinical characterization. In: Abstracts, 7th Cachexia Conference, Kobe. p 50