Endocrine Care

# Vitamin D<sub>3</sub> Supplementation at 4000 International Units Per Day for One Year Results in a Decrease of Positive Cores at Repeat Biopsy in Subjects with Low-Risk Prostate Cancer under Active Surveillance

David T. Marshall, Stephen J. Savage, Elizabeth Garrett-Mayer, Thomas E. Keane, Bruce W. Hollis, Ronald L. Horst, Linda H. Ambrose, Mark S. Kindy, and Sebastiano Gattoni-Celli

Departments of Radiation Oncology (D.T.M., L.H.A., S.G.-C.), Urology (S.J.S., T.E.K.), Medicine (Biostatistics) (E.G.-M.), Pediatrics (B.W.H.), and Neurosciences (M.S.K.), Medical University of South Carolina, Charleston, South Carolina 29425; Ralph H. Johnson Veterans Affairs Medical Center (S.J.S., M.S.K., S.G.-C.), Charleston, South Carolina 29401; and Department of Animal Science (R.L.H.), Iowa State University, Ames, Iowa 50011

Context: We wanted to investigate vitamin D in low-risk prostate cancer.

**Objectives:** The objective of the study was to determine whether vitamin  $D_3$  supplementation at 4000 IU/d for 1 yr is safe and would result in a decrease in serum levels of prostate-specific antigen (PSA) or in the rate of progression.

**Design:** In this open-label clinical trial (Investigational New Drug 77,839), subjects were followed up until repeat biopsy.

**Setting:** All subjects were enrolled through the Medical University of South Carolina and the Ralph H. Johnson Veterans Affairs Medical Center, both in Charleston, SC.

**Patients and Other Participants:** All subjects had a diagnosis of low-risk prostate cancer. Fifty-two subjects were enrolled in the study, 48 completed 1 yr of supplementation, and 44 could be analyzed for both safety and efficacy objectives.

Intervention: The intervention included vitamin D<sub>3</sub> soft gels (4000 IU).

**Main Outcome Measures:** Adverse events were monitored throughout the study. PSA serum levels were measured at entry and every 2 months for 1 yr. Biopsy procedures were performed before enrollment (for eligibility) and after 1 yr of supplementation.

**Results:** No adverse events associated with vitamin  $D_3$  supplementation were observed. No significant changes in PSA levels were observed. However, 24 of 44 subjects (55%) showed a decrease in the number of positive cores or decrease in Gleason score; five subjects (11%) showed no change; 15 subjects (34%) showed an increase in the number of positive cores or Gleason score.

**Conclusion:** Patients with low-risk prostate cancer under active surveillance may benefit from vitamin D<sub>3</sub> supplementation at 4000 IU/d. (*J Clin Endocrinol Metab* 97: 2315–2324, 2012)

The hypothesis that vitamin D could provide anticancer benefits was recognized from epidemiological studies decades ago; Apperly (1) reported the association between

ISSN Print 0021-972X ISSN Online 1945-7197 Printed in U.S.A.

Copyright © 2012 by The Endocrine Society

latitude and cancer mortality based on sun exposure. After being forgotten for decades, the relationship between vitamin D and cancer was resurrected through basic science

doi: 10.1210/jc.2012-1451 Received February 19, 2012. Accepted March 22, 2012. First Published Online April 25, 2012

Abbreviations: IOM, Institute of Medicine; MUSC, Medical University of South Carolina; 25(OH)D, 25-hydroxyvitamin D; 1,25(OH)<sub>2</sub>D, 1,25-dihydroxyvitamin D; 1,25(OH)<sub>2</sub>D<sub>3</sub>, 1,25 dihydroxyvitamin D <sub>3</sub>; PSA, prostate-specific antigen; VDR, vitamin D receptor.

and further epidemiological studies. Abe *et al.* (2), while studying the metabolism of vitamin D in myeloid leukemia cells, made the important observation that 1,25 dihydroxyvitamin D<sub>3</sub> [1,25(OH)<sub>2</sub>D<sub>3</sub>] actually induced the differentiation of these neoplastic cells. It was later determined that this differentiation was due to the regulation of c-myc by 1,25(OH)<sub>2</sub>D<sub>3</sub> (3). A few years later, it was hypothesized that inadequate vitamin D status, resulting from limited solar exposure at higher latitudes, was a risk factor for the development of various noncutaneous malignancies (4–6). Since that time, literally thousands of studies encompassing basic science, clinical trials, and epidemiology have been published with respect to vitamin D and cancer.

Prostate cancer is of great interest from the standpoint of vitamin D interaction because prostate cells express the vitamin D receptor (VDR), vitamin D-25-hydroxylase, 25-hydroxy-vitamin D-1- $\alpha$ -hydroxylase, and the 25-hydroxyvitamin D-24-hydroxylase (7–11). Recent observational studies investigating the association between vitamin D status, as defined by circulating 25-hydroxyvitamin D [25(OH)D], and prostate cancer have produced variable results ranging from protection to increased risk (12– 15). Although one in six men will be diagnosed with prostate cancer during his lifetime, only one in 36 men will die of prostate cancer (American Cancer Society: Learn about Cancer; http://www.cancer.org/Cancer/ProstateCancer/ DetailedGuide/prostate-cancer-key-statistics). Therefore, many indolent prostate cancers are monitored over time in an effort to determine which progress into clinically important, more aggressive cancers. Thus, it provides an excellent model in which to study the effects of enhancing vitamin D status and related changes in tumor progression over an extended period of time. Also, treatment with vitamin D<sub>3</sub> is without any known potential side effects other than hypercalcemia and/or hypercalcuria, which rarely occur and are easily corrected.

Digital rectal examination and measurement of the serum level of prostate-specific antigen (PSA) remain the standard of care for prostate cancer screening, despite significant shortcomings (17). There is general agreement that the introduction of PSA testing led to a measurable decrease in U.S. death rates from prostate cancer (18), but two large randomized trials of PSA screening gave mixed

Variable	Baseline	Follow-up	P value			
Age						
Mean [range]	65 [49, 78]					
Race, n (%)	(					
Caucasian	37 (71%)					
AA	14 (27%)					
Asian	1 (2%)					
25(OH)D (ng/ml serum)			<0.00013			
Mean (sd) [range]	32.8 (13.3) [11.7, 75.0]	66.2 (14.5) [35.7, 95.4]	<0.00001			
PTH (pg/ml serum)			0.000028			
Mean (SD) [range]	45.5 (18.9) [7.6, 117.1]	36.2 (15.3) [8.9, 71.4]	0.00002			
$M_{\text{paper}}(r_{\text{paper}}) [r_{\text{paper}}]$		12 09 /2 72) [C E 20 C]	<0.00001			
1.25(OH) D (n - 10) (ng/ml corum)	0.25 (5.52) [2.2, 15.7]	13.08 (3.73) [0.5, 20.0]	<0.00001			
$M_{\text{pan}}$ (sp) [range]	37 8 (15 2) [17 3 65 7]	<i>1</i> 6 6 (19 6) [22 5 107 5]	0.02			
Months from baseline to follow-up biopsy	57.8 (15.2) [17.5, 65.7]	40.0 (19.0) [22.9, 107.9]	0.02			
Mean (sp) [range]	20.2 (9.8) [12, 63]					
Gleason n(%)	20.2 (3.0) [12, 03]					
0	0	15 (34%)	$0.08^{b}$			
6	50 (100%)	23 (52%)				
7	0	4 (9%)				
8	0	2 (5%)				
Missing	2	8				
Number of positive cores (of 12), n (%)						
0	0 (0%)	15 (34%)				
1	24 (48%)	9 (20%)	0.03 <sup>b</sup>			
2	11 (22%)	7 (16%)				
3	7 (14%)	5 (11%)				
4	4 (8%)	5 (11%)				
5	3 (6%)	3 (7%)				
6	12 (%)	0 (0%)				

AA, African-American.

<sup>a</sup> P values for comparison of pre- and post-25(OH)D and PTH based on paired t test.

<sup>b</sup> P values for comparison of pre- and post-Gleason score and number of positive cores based on the nonparametric sign test.

## **TABLE 1.** Characteristics of supplemented subjects (n = 52)

jcem.endojournals.org 2317

results for improvements in overall survival (19, 20). Nonetheless, the latest analysis of the Surveillance, Epidemiology, and End Results program reveals a very favorable 10-yr outcome in the PSA era, after conservative management of early-stage, well-differentiated prostate cancer (21). Additional longitudinal studies have revealed that the risk of dying from prostate cancer after 20 yr was only 30% with a Gleason score of 6 (22). Therefore, a considerable number of patients diagnosed with lowgrade, low-risk prostate cancer may not require definitive treatment (surgery, radiation, or a combination thereof), especially considering possible side effects such as urinary incontinence, bladder dysfunction, rectal irritation, and erectile dysfunction. Thus, the generally indolent nature of prostate cancer and the risk of side effects from surgery or radiation therapy have led to active surveillance as an option for early-stage, low-risk prostate cancer, in which definitive treatment is deferred to a later date or even indefinitely. However, active surveillance is essentially a monitoring regimen that does not specifically address the issue of how to treat low-risk disease.

It has been recently reported that low serum levels of 25(OH)D (<50 nmol/liter equivalent to <20 ng/ml) strongly correlate with death from prostate cancer (12), raising the possibility that vitamin D deficiency over time would favor the progression of subclinical prostate cancer to clinical disease (23). Additional studies have shown that LNCaP prostate cancer cells treated with 1,25(OH)<sub>2</sub>D<sub>3</sub> undergo apoptotic cell death. Vitamin D-induced apoptosis correlates with down-regulation of Bcl-2 expression and appears to be caspase dependent by primarily triggering the mitochondrial pathway of cell death (24, 25). Furthermore, neoplastic progression in prostate tissue appears to be associated with loss of 25-hydroxyvitamin D<sub>3</sub>  $1\alpha$ -hydroxylase activity (26). However, the latest recommendations from the Institute of Medicine (IOM) concerning dietary reference intake for calcium and vitamin D (27) emphasize that the role of vitamin D in non-bonerelated health issues such as cancer remains unresolved because of the conflicting nature of the available evidence. These doubts have been reaffirmed in a recent article (28). It appears that clinical studies involving robust and sustained, but nontoxic, vitamin D<sub>3</sub> supplementation would shed some clarity on this important issue. Therefore, we sought to determine whether daily supplementation with 4000 IU vitamin D<sub>3</sub> for 1 yr was safe and of clinical benefit in patients electing active surveillance for low-risk prostate cancer.

## **Subjects and Methods**

#### Subjects

Eligible subjects had histologically confirmed diagnosis of adenocarcinoma of the prostate (Gleason score  $\leq 6$ , PSA  $\leq 10$ ,

clinical stage T1c or T2a) and had elected to be managed through active surveillance. For the purpose of eligibility, these additional criteria were verified: serum creatinine 2.0 mg/dl or less, serum phosphate (measured as phosphorus) greater than 2.3 and less than 4.8 mg/dl, and serum calcium greater than 8.5 and less than 10.5 mg/dl. Exclusion criteria were the following: any concurrent malignancy, except nonmelanoma skin cancer; history of sarcoidosis; vitamin  $D_3$  supplementation at 1000 IU/d or greater; history of hypercalcemia; or use of lithium as a medication. This clinical trial was approved by the Institutional Review Board of the Medical University of South Carolina (MUSC) and the Veterans Affairs Medical Center (www.ClinicalTrials.gov Identifier: NCT01045109).

#### Objectives

The objectives of this study were to determine whether vitamin  $D_3$  (4000 IU/d for 1 yr) would be safe and result in a measurable decrease of serum PSA levels in a significant number of enrolled subjects and be associated with a stabilization or improvement of their disease assessed through histological examination of repeat biopsy specimens (Gleason score and number of positive cores), obtained at the end of the study as part of the standard medical care for active surveillance.

#### Study medication

Food and Drug Administration Investigational New Drug no. 77,839 was granted for clinical studies designed to assess the effect of vitamin  $D_3$  (4000 IU/d for 1 yr) on subjects with early-stage prostate cancer. The vitamin  $D_3$  (cholecalciferol) was manufactured by J. R. Carlson Laboratories, Inc. (Arlington Heights, IL).

**TABLE 2.** Characteristics of historical control subjects (n = 19)

			Р
Variable	Baseline	Follow-up	value
Age			
Mean (sd) [range]	66 (7) [50, 81]		
Race, n (%)			
Caucasian	15 (79%)		
AA	4 (21%)		
Asian	0 (0%)		
Months from baseline			
to follow-up			
biopsy			
Mean (sp) [range]	18.4 (8.0) [10, 42]		
Gleason			
0	0 (0%)	2 (11%)	1.00 <sup>a</sup>
6	18 (95%)	13 (68%)	
7	1 (5%)	4 (21%)	
Number of positive			
cores (of 12),			
n (%)			
0	0 (0%)	2 (11%)	0.12 <sup>a</sup>
1	11 (58%)	4 (21%)	
2	5 (26%)	4 (21%)	
3	2 (11%)	4 (21%)	
4	0 (0%)	5 (26%)	
5	1 (5%)	0 (0%)	

AA, African-American.

<sup>a</sup> *P* values for comparison of pre- and post-Gleason score and number of positive cores based on the nonparametric sign test.

#### Methods

Serum levels of vitamin  $D_3$  (cholecalciferol) were determined by HPLC; 25(OH)D, 1,25-dihydroxyvitamin D [1,25(OH)<sub>2</sub>D], and PTH were determined by RIA, as previously described (29, 30). Serum levels of PSA were measured in the Clinical Laboratories of MUSC by the Siemens Advia Centaur Immunoassay System (Chemiluminescence Technology). Assessment of tissue biopsy specimens, before enrollment and after study completion, was part of the standard of care for each subject and was performed by pathologists who were blinded to the study.

#### Clinical trial study design

The open-label study enrolled 52 eligible subjects. Forty-eight subjects completed the study and were included in the safety analysis; these subjects had complete PSA laboratory results for inclusion in the evaluation of changes in PSA. Forty-four subjects had both baseline and repeat biopsy (as part of their standard of care) to compare the number of positive cores and Gleason score with baseline, after completing vitamin D<sub>3</sub> supplementation. All subjects had study visits every 2 months for 1 yr to measure serum levels of 25(OH)D, PSA, phosphorus, and PTH, plus complete blood count, basic metabolic panel, and urinary calcium to creatinine ratio (to rule out any potential toxicities from vitamin D<sub>3</sub> supplementation). In addition, circulating levels of vitamin D<sub>3</sub> (cholecalciferol) and  $1,25(OH)_2D$  (calcitriol) were measured at baseline and exit in 19 subjects for whom extra serum samples were available.

#### Historical control subjects

From an institutional (MUSC) database of more than 700 patients diagnosed with prostate cancer within the last 5 yr, we identified 19 control subjects who were under active surveillance, underwent repeat biopsies, and met all eligibility criteria of the open-label clinical trial, except that they received no vitamin  $D_3$  supplementation. We selected only patients who had their follow-up biopsy at 10 months or later after their initial positive biopsy to ensure comparability with the timing of the follow-up visit of the patients in the open-label trial. All subjects who met eligibility criteria were included. From these control subjects, we abstracted information on their serum PSA levels and assessment of tissue biopsy specimens.

## **Statistical analysis**

For PSA values, before comparison of means or paired differences by *t* test, a natural log transformation was performed to stabilize variance. As a result, hypothesis tests were based on differences in log(PSA+1). Paired *t* tests and signed rank tests were used to assess differences in serum levels of 25(OH)D, log PSA, PTH, cholecalciferol, and 1,25(OH)<sub>2</sub>D at enrollment and



**FIG. 1.** Entry *vs.* exit (after treatment) serum values of 25(OH)D, PTH, and PSA. Each *black point* represents an individual subject. A–C, Entry values are read off the horizontal x-axis (baseline); exit values are read off the vertical y-axis (exit). D–F, Change in serum values of 25(OH)D, PTH, and PSA, respectively. The *marked empty square* ([X]) in each panel represents the mean entry and posttreatment value (A–C) and the mean change in value for each parameter (D–F). Note the significant increase in 25(OH)D serum values (P < 0.00001), the significant decrease in PTH serum values (P = 0.00002), and the overall stability of PSA serum values after vitamin D<sub>3</sub> supplementation (P = 0.27).

at the end of the study. Changes from baseline to follow-up in Gleason score and number of positive scores of biopsies were assessed using a sign test. Fisher's exact test was used to test for differences in proportions.

Random effects models, which included random effects per patient, were used to assess changes in the number of cores over time and to test that the resulting slope was different from zero. The log of number of cores + 1 was used as the outcome to adhere to assumption of the regression (*i.e.* linearity of the trend and homoscedasticity). Due to differentials in times between baseline and follow-up biopsy, we had concerns regarding influence of follow-up biopsies that occurred relatively late (*i.e.* 48 months or longer after the baseline biopsy). Regression models were estimated including and excluding data points greater than 24, greater than 36, and greater than 48 months from baseline to determine how inferences were affected. The inferences were unchanged: the *P* value remained significant, and the magnitude of the slope changes were only slightly dependent on the inclusion or exclusion of the later biopsy data.

# Results

### **Study population**

Eligible subjects were enrolled through the Departments of Radiation Oncology and Urology at MUSC and through the Urology Clinic of the Ralph H. Johnson Veterans Affairs Medical Center. Fifty-two subjects were enrolled in this open-label study. Four subjects were removed from the study



**FIG. 2.** Comparison of repeat biopsy outcome in vitamin D<sub>3</sub>-supplemented subjects. Estimates (in percent) of progression (A) and improvement (B) for all analyzable subjects (n = 44, y-axis) and by entry values of serum 25(OH)D (<20 ng/ml vs. >20 ng/ml, x-axis) are shown. *Bars* are exact 95% confidence intervals (CI). Progression (CI in *parentheses*): 34% of all subjects (0.20, 0.50); 45% of vitamin D-deficient subjects (0.17, 0.77); and 30% of nondeficient subjects (0.16, 0.49). Improvement (CI in *parentheses*): 55% of all subjects (0.39, 0.70); 36% of vitamin D-deficient subjects (0.11, 0.69); and 61% of nondeficient subjects (0.42, 0.77).

due to exclusionary criteria: a diagnosis of colorectal cancer, noncompliance, elevated serum PSA (>10 ng/ml), and voluntary withdrawal. Forty-eight subjects completed 1 yr of vitamin  $D_3$  supplementation. Forty-four subjects who completed supplementation and underwent 12-core biopsy at baseline and at the end of supplementation could be analyzed for both study objectives (Table 1).

The 19 historical controls fit all eligibility criteria for the open-label trial and are described in Table 2. As can be seen from the age, race, and baseline distributions of Gleason scores and number of positive cores, the control population was similar to the open-label study population.

## Safety and efficacy

Supplementation with vitamin D<sub>3</sub> at 4000 IU/d for 1 yr appeared to correct all cases of vitamin D deficiency or insufficiency, although the magnitude of the response varied among subjects. No adverse events linked to vitamin D<sub>3</sub> supplementation at 4000 IU/d were observed. Table 1 and Fig. 1 show that exit values of serum 25(OH)D (expressed as mean  $\pm$  sD in nanograms per milliliter) were significantly higher than entry values (66.2  $\pm$  14.5 vs. 32.8  $\pm$  13.3, respectively; P < 0.00001). As a result of supplementation, serum levels of PTH (expressed as mean  $\pm$  sD in picograms per milliliter) were significantly lower at the end of the study (exit: 36.2  $\pm$  15.3 vs. entry:

> $45.5 \pm 18.9$ ; P = 0.00002), confirming that secondary hyperparathyroidism can be a consequence of vitamin D insufficiency or deficiency. Figure 1, D and E, also shows the change in serum values of 25(OH)D and PTH (exit minus baseline value), respectively, as a function of baseline levels for each of 48 analyzable subjects enrolled in the study and suggests that subjects with lower baseline levels tend to have larger increases. However, serum PSA levels (expressed as median  $\pm$  sp in nanograms per milliliter) remained relatively constant, and changes do not appear to be related to the baseline PSA level (entry:  $4.1 \pm 1.8 vs.$  exit:  $4.5 \pm 2.7$ ; P =0.27), as shown in Fig. 1, C and E. Furthermore, serum PSA levels did not seem to correlate with the repeat biopsy outcome (see below).

> At baseline, the mean number of positive cores ( $\pm$ SD) for the 44 analyzable subjects was 2.1  $\pm$  1.4; however, at repeat biopsy the corresponding mean value decreased to 1.7  $\pm$  1.6. To deter-



**FIG. 3.** Comparison of repeat biopsy outcome between vitamin  $D_3$ -supplemented subjects and historical controls. A, Supplemented subjects appeared to have a lower rate of progression (P = 0.0514) compared with baseline. B, Supplemented subjects appeared to have a significantly higher rate of improvement (P = 0.0257) compared with baseline. Bars are exact 95% confidence intervals (CI).

mine whether individual subjects improved or progressed, we compared the number of positive cores and Gleason grade at baseline and repeat biopsy (Fig. 2). By these criteria, 34% of these subjects progressed (Fig. 2A), defined as an increase in the number of positive cores or in Gleason score; one of the subjects progressed due to an increase in Gleason score (from 6 to 8 in one core), despite a decrease in the number of positive cores (from five to two), and was classified as having had a progression. Five subjects (11%) showed no change. However, more than half of the subjects (55%) showed improvement (Fig. 2B), defined as a decrease in positive cores and no increase in Gleason score at repeat biopsy. Positive core data were also analyzed according to baseline levels of 25(OH)D, subdividing subjects into two groups using 20 ng/ml as a cutoff serum value (Fig. 2). These results suggest that subjects who are vitamin D deficient at baseline [25(OH)D <20 ng/ml serum] may benefit less than the other subjects from vitamin D<sub>3</sub> supplementation, although this inference is premature because of the small number of subjects involved in these subsets analyses.

We also compared the number of positive cores and Gleason grade at baseline and repeat biopsy in the supplemented group and the control group (Fig. 3). Using the same criteria, 63% of the subjects in the control group progressed (because of an increase in the number of positive cores or in Gleason score), compared with 34% in the supplementation group (P = 0.05), as seen in Fig. 3A. Furthermore, in the historical control group, four patients (21%) had improvements in biopsy results, three showed no differences (16%), and 12 (63%) progressed. Comparing the proportions who responded in each group (Fig. 3B), a significant difference was found (P = 0.025).

Figure 4, A and B, shows graphical displays of the positive cores over time in which baseline biopsy occurs at time 0. The baseline and the repeat biopsy for each individual are connected by gray lines. The fitted regression lines (in *black*) indicate that, over time, the number of positive cores tends to decrease in the supplementation group (Fig. 4A), whereas it tends to increase in the untreated control group (Fig. 4B). The slope of the regression line is not significantly different from 0 in the control group (P = 0.19), whereas it is significantly different in the supplementation group (P = 0.017). The slopes of the two fitted regression lines are sig-

nificantly different from one another (P = 0.02). Data represented in Fig. 4A were analyzed using three different thresholds: follow-up through 24, 36, and 48 months, respectively (*i.e.* all data from time points beyond the threshold are not used in the model estimation). Interestingly, the slopes are identical with four decimal places when using the 24- and 48-month time thresholds (b = -0.0146), whereas the slope when using a threshold of 36 months is slightly flatter (-0.0125). All of these values are significant at the P < 0.01 level, evidence of a decline in the number of positive cores over time.

We compared serum levels of 25(OH)D (in nanograms per milliliter) at the end of the study between responders (66.8  $\pm$  14.5) and nonresponders (65.5  $\pm$  14.6), but the values were not significantly different (P = 0.77), suggesting that other factor(s), in addition to the level of vitamin D repletion, affect the response to supplementation.

Circulating values of vitamin D<sub>3</sub> (cholecalciferol) and  $1,25(OH)_2D$  (calcitriol) were measured in 19 subjects for whom extra serum samples were available, and we found significant differences between baseline and exit values of calcitriol (37.8 ± 15.2 *vs.* 46.6 ± 19.6 pg/ml, respectively; P = 0.023) but no obvious correlation with positive core data. However, there was a more significant increase from baseline in circulating levels of cholecalciferol (6.2 ± 3.3)



**FIG. 4.** Trajectories of number of positive cores over time in subjects supplemented with vitamin  $D_3$  (A) and untreated historical control subjects (B). A and B, *Gray lines* represent individual patient data, and *thick solid black lines* are the fitted estimate including all data within 48 months of baseline. The *gray circles* at the ends of some of the *gray lines* refer to subjects with an increase of Gleason score at repeat biopsy or a decrease of Gleason score from baseline. A, The *thick solid black lines* are the fitted estimates including all data for 24-, 36-, and 48-month follow-up period, respectively. The slopes are identical between the 24- and 48-month time thresholds (b = -0.0146), whereas the slope of the 36-month threshold is slightly flatter (-0.0125). All corresponding *P* values are significant: 0.0090 (24 months); 0.0098 (36 months); 0.00093 (48 months). B, The slope is not significantly different from 0 (*P* = 0.17). However, the slopes in A and B (based on all follow-up data) are significantly different from each other (*P* = 0.002).

*vs.*  $13.1 \pm 3.7$  ng/ml, respectively; P < 0.0001), indicating that these levels are much more responsive to supplementation than calcitriol (Fig. 5).

# Discussion

Increasing numbers of patients diagnosed with low-risk prostate cancer are electing active surveillance due to the relative indolent nature of the disease as well as the risks of side effects from definitive treatment (surgery or radiation therapy). This approach includes regular monitoring of the disease through measurements of serum PSA levels, digital rectal examination, and repeat prostate biopsy in an attempt to identify those who are at risk for progressing to clinically relevant disease. To date, it is unclear what other parameters can help clinicians either identify those at risk for progression or provide treatment measures to prevent progression of disease.

The results of our clinical study suggest that serial measurements of PSA serum levels (every 2 months for 1 yr) are a relatively poor predictor of progression in patients with low-risk prostate cancer. However, the combination of active surveillance and vitamin D<sub>3</sub> supplementation at 4000 IU/d resulted in a decreased number of positive cores at repeat biopsy in more than half of patients diagnosed with low-risk prostate cancer. These subjects (responders) are all eligible to remain on active surveillance and do not require definitive treatment (e.g. surgery or radiation therapy). Therefore, this regimen may decrease the chances of overtreatment for patients with low-risk prostate cancer who, based on the results of the repeat biopsy, respond to the combination and remain stable or improve. Conversely, vitamin D<sub>3</sub> supplementation did not benefit 35-40% of subjects (nonresponders) for reasons yet to be investigated. Nevertheless, the combination of active surveillance and vitamin D<sub>3</sub> supplementation may also help identify those patients (nonresponders) who are more likely to need definitive treatment.

The results of this clinical study indicate that changes in circulating levels vitamin  $D_3$  and 25(OH)D are quite substantial. In contrast, the much smaller differences between baseline and exit values of circulating  $1,25(OH)_2D$  sug-

gest that renal production of 1,25(OH)<sub>2</sub>D may be less responsive to vitamin D<sub>3</sub> supplementation and does not necessarily correlate with the tissue levels of calcitriol; furthermore, the prostate 25-hydroxyvitamin D-1- $\alpha$ -hydroxylase, unlike the renal enzyme, is not regulated by PTH and calcium (31). In turn, this suggests that vitamin D action on prostate tissue is through an autocrine/paracrine mechanism, rather than the classic renal endocrine pathway. The mechanism of action is likely to be mediated through the delivery of cholecalciferol and 25(OH)D to prostatic cells, with subsequent activation to 25(OH)D and 1,25(OH)<sub>2</sub>D by CYP2R1 and CYP27B1, respectively (9, 10). A similar mechanism has been described for the control of cathelicidin production by human monocytes and macrophages, for which circulating calcitriol levels are of little consequence (32). The increased circulating levels of cholecalciferol may be of considerable importance because its free concentration would be high enough to enter the cell at an accelerated rate to be hydroxylated first to 25(OH)D and subsequently to  $1,25(OH)_2D$ . It should also be noted that the rate of 25-hydroxylation of vitamin D<sub>3</sub> decreases dramatically with higher initial cir-



**FIG. 5.** Change in serum levels of cholecalciferol (*left panel*) and calcitriol (*right panel*) between baseline (visit 1) and exit (visit 8). *Black lines* and *open circles* represent individual values at baseline and exit; *thick lines* and *closed circles* show mean values at baseline and exit. Testing that there is a difference in visit 1 vs. visit 8 values, both measures show significance, although cholecalciferol levels (in nanograms per milliliter of serum) increased more dramatically than  $1,25(OH)_2D$  levels (in picograms per milliliter of serum) (calcitriol: P = 0.023 based on paired *t* test; P = 0.016, based on nonparametric signed rank test; cholecalciferol: P < 0.0001 based on paired *t* test; P = 0.0003 based on nonparametric signed rank test).

culating 25(OH)D levels. This has been previously reported in a recent study in which production of 25(OH)D decreases when circulating 25(OH)D levels approach 40 ng/ml, probably through feedback inhibition (30).

Comparison of the outcomes of repeat biopsy, between supplemented subjects and historical controls, suggests that supplementation with vitamin  $D_3$  at 4000 IU/d may benefit patients with early-stage, low-risk prostate cancer on active surveillance. However, we realize that there are many limitations to the use of historical controls for comparison purposes. The historical control cohort analyzed in this study was small and there may be additional differences in this group, leading to potential biases. Furthermore, we could not measure levels of circulating vitamin D in these control subjects. Therefore, it is essential to validate the effectiveness of vitamin  $D_3$  supplementation in active surveillance by conducting a larger-scale, randomized clinical trial.

This study was not intended to investigate any mechanism by which vitamin D may interfere with growth, viability, or survival of prostate cancer cells. However, it has been shown that calcitriol is capable of inducing apoptotic death of prostate cancer cells (24, 25). Also, high VDR expression in human prostate cancer is associated with a reduced risk of lethal disease (8). Furthermore, it has been recently reported that in mice a vitamin  $D_3$ -deficient diet combined with low VDR expression results in a tissue environment favoring early procarcinogenic events that enhance prostate cancer risk (33). Therefore, our working hypothesis is that vitamin  $D_3$ supplementation at 4000 IU/d may benefit patients with low-risk prostate cancer by increasing the prostate tissue levels of 1,25(OH)<sub>2</sub>D<sub>3</sub>, which in turn will induce apoptosis of cancer cells. We also propose to use positive cores at repeat biopsy as a progression biomarker to assess the potential benefit of treatment options that specifically target early-stage, lowrisk prostate cancer.

Prevention of cancer by vitamin D in humans has relied mostly on observational studies (34). It should be noted that these types of studies all compare the lowest with the highest levels found within a population that is not supplemented and do not have a comparable group with high enough 25(OH)D levels to inhibit cancer progression (35), as we have observed in this study. Using vitamin D to prevent cancer was recently declared ineffective based on three randomized controlled trials (28).

Two of the three trials were probably not successful because of either too long intervals between supplementation doses (36) or providing an insufficient amount of vitamin  $D_3$  in the treatment arm, which essentially becomes another placebo group (37). Trivedi et al. (36) administered an oral dose of 100,000 IU vitamin D<sub>3</sub> on a quarterly basis for 5 yr, which proved to be effective on skeletal homeostasis but was ineffective on cancer prevention. The Women's Health Initiative study, which involved thousands of subjects, was positive for colorectal cancer vs. circulating 25(OH)D levels based on their nested case-control data (37). However, the interventional arm of the study was negative because of insufficient supplementation: in this study, 400 IU/d of vitamin  $D_3$  for 7 yr as a treatment would have been unable to significantly raise systemic levels of 25(OH)D. In fact, the actual vitamin D intake in this large study was approximately 280 IU/d if their reported compliance rate is taken into account (37). Furthermore, this study did not perform any posttreatment 25(OH)D analyses on the subjects and implied that a 280 IU/d vitamin D supplement would be an effective dose, which is not likely (38). The third randomized controlled trial (39) was a vitamin D-based skeletal study; however, a secondary analysis of the data demonstrated a strong preventive effect on cancer. Our interventional trial is similar to that of Lappe et al. (39), in that both studies used a daily dose of vitamin D<sub>3</sub> that would increase and maintain a stable circulating level of 25(OH)D during the study period, although the dosage used in our study (4000 IU/d) resulted in much higher systemic circulating levels of 25(OH)D. It is worth noting that in both the study by Lappe et al. (39) and our interventional trial, baseline 25(OH)D levels were 70-80 nmol and increased significantly after supplementation to produce the prevention/ treatment effects on cancer. According to the recent IOM report, our mean basal circulating 25(OH)D level of 80 nmol far exceeded the recommended level of 50 nmol (27). Furthermore, the IOM report went as far as to say that circulating 25(OH)D levels exceeding 125 nmol could be a risk factor for developing cancer. Our study resulted in 25(OH)D levels reaching 175 nmol with no clear adverse events but with potential oncologic benefits. We therefore hypothesize that there are different vitamin D requirements or thresholds for different biological systems (e.g. skeletal vs. prostate).

It is important to point out the absence of any measurable toxicity of the vitamin  $D_3$  treatment used in this clinical study. Supplementation with 4000 IU/d for 1 yr elevated circulating 25(OH)D to a range recently recommended by The Endocrine Society practice guidelines (40). These guidelines, compared with those of the Food and Nutritional Board of the IOM (16), recommend a 2- to 3-fold increase in vitamin D intake, with a tolerable upper intake level of 10000 IU/d. The IOM report also concluded that circulating 25(OH)D levels were unrelated to various neoplasias, including prostate cancer, and in fact higher levels of circulating 25(OH)D could make the cancer worse. These conclusions of the IOM report are not supported by our current data. Our next objective is to conduct a definitive, randomized trial to validate the effectiveness of vitamin  $D_3$ supplementation in active surveillance, using an intervention strategy that is extremely cost effective and easy to implement.

# Acknowledgments

The contents of this paper do not represent the views of the Department of Veterans Affairs or the United States Government. We thank Drs. Leander Cannick, Jason Zauls, Kyle Russo, and Lewis Cooper (Medical University of South Carolina Department of Radiation Oncology) for helping with the recruitment of eligible subjects.

Address all correspondence and requests for reprints to: Sebastiano Gattoni-Celli, M.D., Strom Thurmond Biomedical Research Building, Room 338C, 114 Doughty Street, Charleston, South Carolina 29403. E-mail: gattonis@musc.edu.

This work was supported in part by Gateway for Cancer Research Grant G-06-068), the Health Services Research and Development Program of the Department of Veterans Affairs, Veterans Affairs Merit Award CX000163, the Biostatistics Shared Resource, Hollings Cancer Center's Support Grant P30 CA138313 (National Institutes of Health/National Center for Research Resources Grant UL1 RR029881), and South Carolina Clinical and Translational Research Institute Grant UL1 RR029882.

Disclosure Summary: B.W.H. is an academic consultant to the DiaSorin Corp. The other authors have nothing to disclose.

# References

- 1. Apperly FL 1941 The relation of solar radiation to cancer mortality in North America. Cancer Res 1:191–195
- Abe E, Miyaura C, Sakagami H, Takeda M, Konno K, Yamazaki T, Yoshiki S, Suda T 1981 Differentiation of mouse myeloid leukemia cells induced by 1 alpha, 25-dihydroxyvitamin D<sub>3</sub>. Proc Natl Acad Sci USA 78:4990–4994
- Reitsma PH, Rothberg PG, Astrin SM, Trial J, Bar-Shavit Z, Hall A, Teitelbaum SL, Kahn AJ 1983 Regulation of myc gene expression in HL-60 leukaemia cells by a vitamin D metabolite. Nature 306:492– 494
- 4. Garland CF, Garland FC 1980 Do sunlight and vitamin D reduce the likelihood of colon cancer? Int J Epidemiol 9:227–231
- Garland FC, Garland CF, Gorham ED, Young JF 1990 Geographic variation in breast cancer mortality in the United States: a hypothesis involving exposure to solar radiation. Prev Med 19:614–622
- 6. Lefkowitz ES, Garland CF 1994 Sunlight, vitamin D, and ovarian cancer mortality rates in U.S. women. Int J Epidemiol 23:1133–1136
- Miller GJ, Stapleton GE, Ferrara JA, Lucia MS, Pfister S, Hedlund TE, Upadhya P 1992 The human prostatic carcinoma cell line LNCaP expresses biologically active, specific receptors for 1α,25-dihydroxyvitamin D<sub>3</sub>. Cancer Res 52:515–520
- Hendrickson WK, Flavin R, Kasperzyk JL, Fiorentino M, Fang F, Lis R, Fiore C, Penney KL, Ma J, Kantoff PW, Stampfer MJ, Loda M, Mucci LA, Giovannucci E 2011 Vitamin D receptor protein expression in tumor tissue and prostate cancer progression. J Clin Oncol 29:2378–2385
- Ellfolk M, Norlin M, Gyllensten K, Wikvall K 2009 Regulation of human vitamin D<sub>3</sub> 25-hydroxylases in dermal fibroblasts and prostate cancer LNCaP cells. Mol Pharmacol 75:1392–1399
- Schwartz GG, Whitlatch LW, Chen TC, Lokeshwar BL, Holick MF 1998 Human prostate cells synthesize 1,25dihydroxyvitamin D<sub>3</sub> from 25-hydroxyvitamin D<sub>3</sub>. Cancer Epidemiol Biomarkers Prev 7:391–395
- Deeb KK, Luo W, Karpf AR, Omilian AR, Bshara W, Tian L, Tangrea MA, Morrison CD, Johnson CS, Trump DL 2011 Differential vitamin D 24-hydroxylase/CYP24A1 gene promoter methylation in endothelium from benign and malignant human prostate. Epigenetics 6:994–1000
- Tretli S, Hernes E, Berg JP, Hestvik UE, Robsahm TE 2009 Association between serum 25(OH)D and death from prostate cancer. Brit J Cancer 100:450–454
- 13. Fang F, Kasperzyk JL, Shui I, Hendrickson W, Hollis BW, Fall K, Ma J, Gaziano JM, Stampfer MJ, Mucci LA, Giovannucci E 2011 Prediagnostic plasma vitamin D metabolites and mortality among patients with prostate cancer. PLoS One 6:e18625
- Yin L, Raum E, Haug U, Arndt V, Brenner H 2009 Meta-analysis of longitudinal studies: serum vitamin D and prostate cancer risk. Cancer Epidemiol 33:435–445
- 15. Tuohimaa P, Tenkanen L, Ahonen M, Lumme S, Jellum E, Hallmans G, Stattin P, Harvei S, Hakulinen T, Luostarinen T, Dillner J, Lehtinen M, Hakama M 2004 Both high and low levels of blood vitamin D are associated with a higher prostate cancer risk: a longitudinal, nested case-control study in the Nordic countries. Int J Cancer 108:104–108

- 16. Ross AC, Manson JE, Abrams SA, Aloia JF, Brannon PM, Clinton SK, Durazo-Arvizu RA, Gallagher JC, Gallo RL, Jones G, Kovacs CS, Mayne ST, Rosen CJ, Shapses SA 2011 The 2011 report on dietary reference intake for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. J Clin Endocrinol Metab 96:53–58
- 17. Thompson IM, Pauler DK, Goodman PJ, Tangen CM, Lucia MS, Parnes HL, Minasian LM, Ford LG, Lippman SM, Crawford ED, Crowley JJ, Coltman Jr CA 2004 Prevalence of prostate cancer among men with a prostate-specific antigen level < or = 4.0 ng per milliliter. N Engl J Med 350:2239–2246
- Ries LAG, Melbert D, Krapcho M, Stinchcomb DG, Howlader N, Horner MJ, Mariotto A, Miller BA, Feuer EJ, Altekruse SF, Lewis DR, Clegg L, Eisner MP, Reichman M, Edwards BK, eds. 2009 Surveillance Epidemiology and End Results (SEER) Cancer Statistics Review, 1975–2006. National Cancer Institute. Bethesda, MD. http:// seer.cancer.gov/csr/1975\_2006/, based on November 2008 SEER
- Andriole GL, Crawford ED, Grubb 3rd RL, Buys SS, Chia D, Church TR, Fouad MN, Gelmann EP, Kvale PA, Reding DJ, Weissfeld JL, Yokochi LA, O'Brien B, Clapp JD, Rathmell JM, Riley TL, Hayes RB, Kramer BS, Izmirlian G, Miller AB, Pinsky PF, Prorok PC, Gohagan JK, Berg CD; PLCO Project Team 2009 Mortality results from a randomized prostate-cancer screening trial. N Engl J Med 360:1310–1319
- 20. Schröder FH, Hugosson J, Roobol MJ, Tammela TL, Ciatto S, Nelen V, Kwiatkowski M, Lujan M, Lilja H, Zappa M, Denis LJ, Recker F, Berenguer A, Määttänen L, Bangma CH, Aus G, Villers A, Rebillard X, van der Kwast T, Blijenberg BG, Moss SM, de Koning HJ, Auvinen A; ERSPC Investigators 2009 Screening and prostate-cancer mortality in a randomized European study. N Engl J Med 360:1320–1328
- Lu-Yao GL, Albertsen PC, Moore DF, Shih W, Lin Y, DiPaola RS, Barry MJ, Zietman A, O'Leary M, Walker-Corkery E, Yao SL 2009 Outcomes of localized prostate cancer following conservative management. JAMA 302:1202–1209
- Albertsen PC, Hanley JA, Fine J 2005 20-Year outcomes following conservative management of clinically localized prostate cancer. JAMA 293:2095–2101
- Schwartz GG 2005 Vitamin D and the epidemiology of prostate cancer. Semin Dial 18:276–289
- Blutt SE, McDonnell TJ, Polek TC, Weigel NL 2000 Calcitriolinduced apoptosis in LNCaP cells is locked by overexpression of Bcl-2. Endocrinology 141:10–17
- Guzey M, Kitada S, Reed JC 2002 Apoptosis induction by 1α,25-dihydroxyvitamin D<sub>3</sub> in prostate cancer. Mol Cancer Ther 1:667–677
- 26. Hsu JY, Feldman D, McNeal JE, Peehl DM 2001 Reduced  $1\alpha$ -hydroxylase activity in human prostate cancer cells correlates with decreased susceptibility to  $25(OH)D_3$ -induced growth inhibition. Cancer Res 61:2852-2856
- 27. Institute of Medicine 2011 Dietary references intakes for calcium and vitamin D. Washington, DC: National Academies Press
- Manson JE, Mayne ST, Clinton SK 2011 Vitamin D and prevention of cancer: ready for prime time? N Engl J Med 364:1385–1387

- 29. Hollis BW, Kamerud JQ, Selvaag SR, Lorenz JD, Napoli JL 1993 Determination of vitamin D status by radioimmunoassay with an <sup>125</sup>I-labeled tracer. Clin Chem 39:529–533
- Hollis BW, Johnson D, Hulsey TC, Ebeling M, Wagner CL 2011 Vitamin D supplementation during pregnancy: double blind, randomized clinical trial of safety and effectiveness. J Bone Miner Res 26:2341–2357
- 31. Young MV, Schwartz GG, Wang L, Jamieson DP, Whitlatch LW, Flanagan JN, Lokeshwar BL, Holick MF, Chen TC 2004 The prostate 25-hydroxyvitamin D-1α-hydroxylase is not influenced by parathyroid hormone and calcium: implications for prostate cancer chemoprevention by vitamin D. Carcinogenesis 25:967–971
- 32. Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzic SR, Ochoa MT, Schauber J, Wu K, Meinken C, Kamen DL, Wagner M, Bals R, Steinmeyer A, Zugel U, Gallo RL, Eisenberg D, Hewison M, Hollis BW, Adams JS, Bloom BR, Modlin RL 2006 Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. Science 311:1770–1773
- 33. Kovalenko PL, Zhang Z, Yu JG, Li Y, Clinton SK, Fleet JC 2011 Dietary vitamin D and vitamin D receptor level modulate epithelial cell proliferation and apoptosis in the prostate. Cancer Prev Res 4:1617–1625
- 34. Giovannucci E 2011 The epidemiology of vitamin D and cancer Risk. In: Feldman D, Pike JW, Adams JS, eds. Vitamin D. 3rd ed. Academic Press, San Diego, CA; 1569–1590
- Krishnan AV, Feldman D 2011 Vitamin D and prostate cancer. In Feldman D, Pike JW, Adams JS, eds. Vitamin D. 3rd ed. Academic Press, San Diego, CA; 1675–1709
- 36. Trivedi DP, Doll R, Khaw KT 2003 Effect of four monthly oral vitamin D<sub>3</sub> (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomized double-blind controlled trial. BMJ 326:469
- 37. Wactawski-Wende J, Kotchen JM, Anderson GL, Assaf AR, Brunner RL, O'Sullivan MJ, Margolis KL, Ockene JK, Phillips L, Pottern L, Prentice RL, Robbins J, Rohan TE, Sarto GE, Sharma S, Stefanick ML, Van Horn L, Wallace RB, Whitlock E, Bassford T, Beresford SA, Black HR, Bonds DE, Brzyski RG, Caan B, Chlebowski RT, Cochrane B, Garland C, Gass M, Hays J, Heiss G, Hendrix SL, Howard BV, Hsia J, Hubbell FA, Jackson RD, Johnson KC, Judd H, Kooperberg CL, Kuller LH, LaCroix AZ, Lane DS, Langer RD, Lasser NL, Lewis CE, Limacher MC, Manson JE; Women's Health Initiative Investigators 2006 Calcium plus vitamin D supplementation and the risk of colorectal cancer. N Engl J Med 354:684–696
- Vieth R, Cole DE, Hawker GA, Trang HM, Rubin LA 2001 Wintertime vitamin D insufficiency is common in young Canadian women, and their vitamin D intake does not prevent it. Eur J Clin Nutr 55:1091–1097
- 39. Lappe JM, Travers-Gustafson D, Davies KM, Recker RR, Heaney RP 2007 Vitamin D and calcium supplementation reduces cancer risk: results of a randomized trial. Am J Clin Nutr 85:1586–1591
- 40. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM 2011 Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. J Clin Endocrinol Metab 96:1911–1930