



Photo © by Dr Vasquez, 2015

Review • Survey • Nutrition • Dietary Intervention • Public Health • Clinical Care • Pesticides

Survey Reports Improved Health After Avoiding Genetically Modified Foods

Part 1: Health Concerns—GMOs, Bt-toxin, and Roundup®

Part 2: Survey Results

Part 3: Focus on Digestive Disorders

Jeffrey M Smith MBA, Director of Institute for Responsible Technology

Abstract: A survey of 3,256 respondents, primarily residing in the United States, reveal improvements in a wide range of health symptoms following the removal or reduced consumption of genetically engineered foods, also called genetically modified organisms or GMOs. The changes are consistent with reports by physicians and others about improvements accompanying a switch to largely non-GMO and organic diets. The conditions that were most frequently reported as showing improvement include: Digestive: 85.2%, Fatigue, low energy: 60.4%, Overweight or obesity: 54.6%, Clouding of consciousness, “brain fog”: 51.7%, Food allergies or sensitivities: 50.2%, Mood problems, such as anxiety or depression: 51.1%, Memory, concentration: 48.1%, Joint pain: 47.5%, Seasonal allergies: 46.6%, Gluten sensitivities: 42.2%, Insomnia: 33.2%, Other skin conditions (not eczema): 30.9%, Hormonal problems: 30.4%, Musculoskeletal pain: 25.2%, Autoimmune disease: 21.4%, Eczema: 20.8%, and Cardiovascular problems, including high blood pressure: 19.8%. Mechanisms by which GMOs may contribute to digestive disorders—the most frequently reported symptoms improved by GMO avoidance—is discussed. Three possible modes of action by GMOs are evaluated: 1) the disruptive and unpredictable nature of the process of genetic modification itself, which can introduce or elevate allergens, toxins, and anti-nutrients; 2) possible allergenic and toxic effects of Bt toxin, the insecticide produced within most genetically engineered corn varieties grown in the United States (US); and 3) the health impacts of glyphosate-based herbicides, such as Roundup®, which are sprayed on and absorbed into most genetically engineered food crops. The studies in these areas support several potential causative pathways leading to digestive disorders and may help explain why these and other related diseases have been rising in parallel with the increased acreage of GMOs and the application of Roundup® on these crop acres.

Part 1: Introduction—Discussion of Health Concerns—GMOs, Bt-toxin, and Roundup® Widespread Use of Untested GM Foods and Related Pesticide Chemicals

The process of genetic modification (GM) involves the transfer or rearrangement of genetic material within or between species’ DNA using laboratory techniques. These

laboratory techniques are distinct from natural methods, such as hybridization, that alter the genome through sexual reproduction.

Most of the currently commercialized crops known as genetically modified organisms (GMOs) have had non-plant genes inserted into their DNA. Such genes are usually taken from bacteria or viruses, to confer a

Updates: The most complete version of and any updates to this article are available at the website of *International Journal of Human Nutrition and Functional Medicine* ® <http://intjhumnutrfunctmed.org/> ISSN 2378-4881

Copyrights: Copyright © by author(s) and International College of Human Nutrition and Functional Medicine ® www.ICHNFM.org

Free access: Freely available and distributable; all content, text, and image rights reserved by author(s) and ICHNFM.

Citation: Smith JM. Survey Reports Improved Health After Avoiding Genetically Modified Foods. *Int J Hum Nutr Funct Med* 2017; article in review

particular trait. Eleven genetically modified (GM) food crops are currently grown for commercial consumption. The six major GM crops are soy, corn, cotton, canola, sugar beets, and alfalfa, all of which are used as food for humans and animals. Cottonseed and canola are also processed into food-grade oils and sugar beets are refined to make sugar.

All six major GMOs are engineered to be herbicide tolerant (HT), i.e. to survive spray applications of herbicide (vernacular: “weed killer”). HT crops comprise 89% of all GMOs grown in the US. By far, the most widely grown HT variety of crops is called “Roundup[®] Ready” (RR), produced by Monsanto Company to withstand field treatments of Roundup[®] herbicide with glyphosate as the active ingredient, which is absorbed into the crop.¹ The food portion of RR crops contains high residue levels of glyphosate.² As of 2016, 94% of soybeans grown in the US were RR.³ Other varieties of GMOs are engineered to withstand the herbicide glufosinate, and more recently, Dicamba and 2,4-D.

Some varieties of corn and cotton have genes from *Bacillus thuringiensis*, a soil bacteria variety, inserted into their cells, which produce a toxic insecticide called *Bt* toxin. As of 2016, 76% of corn grown in the US is both *Bt*-producing (*Bt*) and HT. Corn with only the *Bt* trait comprises just 3% of the US corn acreage, while HT-only corn comprises 13%. For cotton, 80% are both *Bt* and HT, 4% are *Bt* only, and 9% are HT only.⁴

Varieties of GMO zucchini, yellow squash, and papaya varieties have virus genes inserted into them, which are designed to provide resistance to infections from specific plant viruses. Two newly approved crops, apples and potatoes, were engineered using double stranded RNA technology, which suppresses expression of the gene that causes the food to oxidize and discolor (i.e. turn brown when sliced). A small amount of GMO apples was commercially released for the first time in 2016 and according to the potatoes’ developer, J. R. Simplot, GMO potatoes have been sold via supermarkets.⁵ Numerous other types of GMO crops have been developed and many have been subject to field trials.

Background

The US Food and Drug Administration (FDA) policy regarding GMOs, implemented in 1992 and still in force, allows GMO makers to determine on their own if their foods are “Generally Recognized as Safe” (GRAS). If they are deemed to be GRAS, the FDA does not require any safety studies or labels.

The FDA also does not require that companies submit any data to the agency, but offers companies a voluntary pre-market consultation. There are no requirements or safety testing standards. Documentation

provided by the GMO makers is typically summary in nature with no raw data.⁶

At the end of the consultation, the FDA releases a letter to the company acknowledging that it is the responsibility of the GMO maker to determine that their foods are safe, and that the voluntary consultation process simply confirms that the company made that determination. In one letter to Monsanto regarding their MON810 *Bt* corn, the FDA Regulatory Affairs Manager wrote, “Based on the safety and nutritional assessment *you* have conducted, it is our understanding that *Monsanto* has concluded that corn products derived from this new variety are not materially different in composition, safety, and other relevant parameters from corn currently on the market, and that the genetically modified corn does not raise issues that would require premarket review or approval by FDA.” (Emphasis added.) *The letter does not state that the FDA declares that the GMO is safe.*⁷

The agency justified this hands-off approach in the policy document by declaring that it “wasn’t aware of any information” showing that GMOs were different “in any meaningful or uniform way”⁸ — a direct contradiction to the opinions voiced in the memos from their scientists.

In 1998 the FDA was forced to turn over tens of thousands of pages of internal memos related to GMOs due to a lawsuit filed by the Alliance for Bio-integrity. The memos⁹ revealed that agency scientists who were tasked to help create the FDA policy on GMOs had repeatedly warned their superiors that GMO foods were quite different than foods created from traditional breeding. The technology, they said, could create serious side effects, such as allergies, toxins, antibiotic-resistant diseases, and nutritional problems. They urged their superiors to require rigorous long-term tests.

The same set of FDA documents also revealed that the White House had instructed the agency to promote biotechnology.¹⁰ Then Vice-President Dan Quayle said it was necessary to “resist the spread of unnecessary regulation” in order to keep America the “world leader in biotechnology.”¹¹

The person who oversaw the GMO policy for the FDA was a political appointee, Michael Taylor, the former outside attorney for Monsanto, later the company’s vice president of government and regulatory affairs, and later the Deputy Commissioner for Foods at the FDA.

Although several other nations require some limited safety data, even those requirements are widely criticized as inadequate, obsolete, and inappropriately reliant on unpublished research provided by the GMO producers themselves. There are very few safety studies that would be rigorous enough to be called “academic.” Although proponents of GMOs often point to compilations of hundreds of studies, the vast majority of

**See associated Editorial
and Comments attached
with this article**

these are considered commercial in nature. They look at data for market considerations, but rarely have relevant designs for safety evaluations. Furthermore, analysis of industry “safety” studies has revealed research protocols apparently designed to hide evidence of harm.¹²

No human clinical trials and no post-market surveillance on health outcomes related to GMO consumption have been conducted. In 2002, Health Canada announced that it would monitor the health of Canadians to see if GMOs adversely affected health, but by the next year, according to CBC TV news, they “abandoned that research less than a year later saying it was ‘too difficult to put an effective surveillance system in place.’” The reporter added, “So at this point, there is little research into the health effects of genetically modified food. So will we ever know for sure if it’s safe?”¹³ In 1998, it was discovered that the United Kingdom’s Food Standards Agency had asked supermarket executives for the purchasing data from the 30 million consumers using loyalty cards, so they could see if those eating GMOs had higher rates of cancer, birth defects, childhood allergies, or hospital admissions. When the data collection plans were made public, the government, which had told the public that GMOs were safe, withdrew the program.¹⁴

In the face of insufficient pre- and post-marketing safety studies, extra attention must be paid to reports from individuals and/or their healthcare providers about potential reactions to the inclusion or exclusion of GMOs from their diet. Although correlation does not equal causation, any correlations between national health statistics and GMO consumption should be carefully investigated. Otherwise, even a significant rise in disease rates related to GMOs will easily go unnoticed.

Survey design informed by physician and personal reports

On May 8, 2009, the American Academy of Environmental Medicine (AAEM) published their policy paper on GMOs, which included a review of several peer-reviewed safety studies. Several animal studies, according to their policy paper, reveal a long list of disorders, including “infertility, immune dysregulation, accelerated aging, dysregulation of genes associated with cholesterol synthesis, [faulty] insulin regulation, cell signaling, and protein formation, and changes in the liver, kidney, spleen and gastrointestinal system.” The policy concludes, “There is more than a casual association between GM foods and adverse health effects. There is causation as defined by Hill’s Criteria in the areas of strength of association, consistency, specificity, biological gradient, and biological plausibility.”¹⁵ The AAEM called on the US government to implement an immediate moratorium on all GM foods and urged physicians to prescribe non-GMO diets for all patients.

“Physicians are probably seeing the effects in their patients,” said AAEM past-president Jennifer Armstrong, MD, “but need to know how to ask the right questions.”¹⁶ According to David Schubert, PhD, of the Salk Institute, the patients at greatest risk from consuming GMOs are the very young. “Children are the most likely to be adversely affected by toxins and other dietary problems.”¹⁷

Starting in November 2009, the author of this paper (JMS) began interviewing physicians and other healthcare providers who advise their patients to switch to a non-GMO and/or organic diet, asking what outcomes they observed. Audiences at numerous medical and healthcare conferences, as well as at more than 100 public lectures, were informally surveyed from the stage. Audience members shared which symptoms or conditions improved after the dietary changes. Commonly, after individual audience members shared their stories, numerous others raised their hands to indicate that they too experienced similar improvements.

The selection of health conditions used in the formal survey reported herein was based primarily on the thousands of responses by audience members, as well as numerous private conversations and email exchanges with individuals and healthcare practitioners.

Three mechanisms by which GMOs may cause health problems

Reviewed herein are three main mechanisms by which GMOs might produce or exacerbate the conditions listed in the survey: 1) the generic side-effects of the GMO transformation process, 2) the *Bt* toxin found in GMO corn and cotton plants and 3) the herbicides—particularly glyphosate-based herbicides (GBHs)—that are sprayed on most GMOs.

Collateral effects of genetic engineering

The process of creating a GMO crop results in significant damage to the host organism, with hundreds or thousands of mutations possible throughout the plants’ genome.¹⁸ A GM plant’s total DNA can be 2-4% different from that of its natural parent.¹⁹ In addition, up to 5% of its natural genes can alter their levels of protein expression because of a single insertion.²⁰

These changes in the genetic sequence and expression can impact numerous other compounds and phytochemicals that make up a plant. For example, Monsanto’s data on cooked GM soybeans shows as much as seven times the level of trypsin inhibitor, a natural soy allergen, and a doubling of soy lectin, an anti-nutrient that can potentially block nutrient absorption.²¹ Monsanto’s MON810 *Bt* corn has 43 genes that are significantly altered in their expression levels. One of these, which produces an allergenic protein called gamma zein, is normally switched off in corn. In Monsanto’s GMO variety, however, the allergenic protein is expressed.²² In

addition, GM corn and soy produce higher amounts of lignin.²³

The most comprehensive comparison of a GMO crop and its non-GMO equivalent to date, conducted by Antoniou et al., showed that Monsanto's RR corn has 117 proteins and 91 small molecule biochemicals that are significantly different from natural corn. For example, there was an increase of several polyamines in the GMO corn, including putrescine and cadaverine. In addition to being responsible for the foul odor of rotting flesh, these two substances, according to co-author Dr. Michael Antoniou, "enhance the effects of histamine, thus heightening allergic reactions, and both have been implicated in the formation of carcinogenic substances called nitrosamines with nitrite in meat products."²⁴

Unanticipated changes in proteins

1. Amino acid sequence

Side-effects from the process of genetic engineering can result in unexpected changes in the amino acid sequence of the transgene product. For example, the transgene construct inserted into RR soybeans was designed to create a fixed length of RNA transcript. Instead, the portion of the transgene that was supposed to determine the length of the transgene (NOS terminator) failed to function correctly. This resulted in overly long RNA transcripts *that do not exist in nature* with some potentially able to produce proteins that are different from the intended/targeted result. Most GMOs use the same inefficient NOS terminator.

2. Glycosylation

Proteins produced in transgenic organisms may be modified by addition of materials, such as sugars, in unpredictable ways. The binding of sugar (glycosylation) to proteins can convert a benign protein, such as naturally produced by beans, into an immunoreactive and potentially allergenic protein, when produced in transgenic peas.²⁵

3. Misshaped proteins

The shape of a protein is critical to its function. The process of genetic engineering may result in unpredicted alterations of protein shape and size, with potentially dangerous effects. In a proteomics analysis of MON810, for example, researchers discovered that seed storage proteins in the Bt corn were truncated, which they described "as a major concern."²⁶

One reason proteins can be misshaped is if they are folded improperly. When the polypeptides produced from inserted foreign genes fold after synthesis in the potentially different cellular environment of the GMO, that new environment may have characteristics (e.g. altered pH or a lack of needed "chaperone" proteins) that affects folding in ways different from their native context.

Likewise, the new cellular context may cause them to denature more frequently.

Unanticipated effects from altered RNA

It is now understood that certain types of small RNA molecules can have a direct and significant impact on gene expression through a process known as RNA interference (RNAi). RNAi usually results in reducing expression of certain genes, which in turn can lead to decreases or increases in expression of others. This may impact an organism's function and health. The process of engineering small RNA molecules into GMOs also has the potential of unintentionally producing additional small RNAs that can interfere with the function of genes that are not being targeted. Since small RNAs in food have been found to survive digestion and enter the body of the consumer,²⁷ ingestion of the intended and unintended small RNA molecules in the GMO can alter gene function with unknown consequences.

Gene transfer from GMOs

Transgenes may horizontally transfer to humans or other organisms. While GMO DNA has been identified in various organs and the blood of animals fed GMOs, the studies were unable to determine if they were integrated into the host cells' genome. Netherwood et al²⁸ confirmed that part of the RR soybean transgene transferred and integrated into bacteria living inside human intestines. It is not known whether the transformed bacteria actively expressed the transgenic protein. If it is expressed, then our gut flora may be compromised by GMOs and forced to produce GMO proteins continuously inside our digestive tract.

Evidence of adverse effects from GMO process

A striking example of the damage caused by the generic GMO process was highlighted in experiments by Arpad Pusztai. Commissioned by the United Kingdom (UK) government to design testing protocols for GMO food safety, Pusztai and his team came up with a system to better identify the impacts of the inserted transgene as well as the unintended consequences due to the process of genetic engineering.²⁹ To demonstrate the protocol, they used a GM potato engineered with a gene from the snowdrop plant that produces a protein with insecticidal properties called galanthus nivalis lectin (GNA). Pusztai and his colleagues conducted extensive research on GNA for nearly seven years and found it to be harmless to rats.


As part of their feeding trial, groups of rats were fed diets with GMO potatoes, natural potatoes of the same type, or natural potatoes spiked with added GNA in the same amount produced by the GMO potato.³⁰

The GMO potatoes adversely affected virtually every organ system of young rats—with most changes found after just 10 days. The diet with added non-GMO GNA, however, did not produce such harm. This

demonstrated that effects from the GMO process—other than the lectin itself—were toxic to the animals. The impacts of the GMO potato included the following:³¹

- The young rats’ brains, livers, and testicles were generally smaller, suggesting disruption of normal growth processes due to either malabsorption of nutrients or unknown toxins.
- White blood cells responded to a challenge more slowly, indicating immune system damage; organs related to the immune system, including the thymus and the spleen showed changes.
- The animals had enlarged pancreases and intestines, and partial atrophy of the liver.⁴
- In all cases, the GM potato created proliferative cell growth in the stomach and in both the small and large intestines; the lining was significantly thicker than controls (see Figure 1). Although no tumors were detected, such growth may indicate a precancerous condition.

Figure 1. The digestive tract of rats fed GM potatoes engineered to produce GNA lectin showed excessive cell growth compared to rats fed non-GMO potatoes. Another group of rats fed non-GMO potatoes plus the GNA lectin did not exhibit the cell growth. This suggests that the process of genetically engineering the potato, and not the lectin, was the cause.



Stomach

The stomach lining of GM-fed rats showed proliferative cell growth.

Intestines

The excessive cell growth was also found in the wall of the small intestines (crypts) in rats fed GM potatoes.

Photos provided by Stanley Ewen.

The studies conducted by the makers of GMOs have not used the same rigorous approach as Pusztai. His approach includes the third feeding group in which animals consume non-GMO crops spiked with just the protein produced in the GMO. Thus, their studies cannot evaluate which of the side-effects are due to the specific gene product and which are due to the generic transformation process itself. Industry studies do not generally test for the type of health effects found in Pusztai’s rats, leaving us without valuable and necessary information regarding the biochemical products, physiological effects, and clinical consequences of GMO foods.

Consumption of the Bt Toxin insecticide

Two types of commercialized crops, corn and cotton, are engineered to produce an insecticide called *Bt* toxin. The

only product from cotton that we consume directly is cottonseed oil, which is generally void of proteins and would therefore not be a source of dietary *Bt* toxin. *Bt* corn, however, can expose us to the toxin via fresh corn and corn products, such as corn chips, polenta, and tortillas.

Bt toxin is produced from *Bacillus thuringiensis* soil bacteria. In its natural state, the toxin creates small holes in insects’ gut walls, killing them. It is believed that the gut bacteria within the insect move through the “leaky gut” and then kill the insect.³²

Genetic engineers have inserted various altered *Bt* toxin genes directly into the DNA of corn and cotton plants so the crops produce the toxin in every cell. To justify this addition to our food supply, biotech companies and the US Environmental Protection Agency (EPA) claimed that *Bt* toxin and *Bacillus thuringiensis* in their natural forms were used as natural methods of pest control for years, with no impact on humans or mammals of any type and were therefore only dangerous to certain insects.³³ However, several peer-reviewed published studies contradict this assertion.

Studies in mice showed that natural *Bt* toxin provoked systemic and mucosal immune system responses as powerful as cholera toxin. Furthermore, exposure to *Bt* toxin sensitized the mice so that their immune system responded to formerly harmless substances.³⁴

A 2008 Italian government study found that *Bt* corn provoked profound immune responses in mice.³⁵ Monsanto’s own rat studies with *Bt* corn also showed toxicity and immune responses.³⁶ Another mouse study confirmed that *Bt* toxin is cytotoxic.³⁷

A 2012 article in *Journal of Applied Toxicology*³⁸ “documented that modified *Bt* toxins [from GM plants] are not inert on human cells, but can exert toxicity.” In vitro and in generally higher concentrations than that which is expected to be produced in average *Bt* corn, researchers found that *Bt* toxin disrupts the cell membrane of human cells in just 24 hours, causing fluid to leak through the cell walls.

In the US, farmworkers exhibited antibody responses to *Bt* toxin and hundreds of people in the Pacific Northwest, who were inadvertently sprayed with *Bt* when it was used to kill gypsy moths, exhibited allergic and flu-like symptoms. Some workers had to go to the hospital.³⁹

Numerous reports, including medical investigations and hospital records, show that thousands of agricultural workers in India exposed to *Bt* cotton varieties reported skin rashes and other health symptoms.^{40,41}

US EPA regulators assumed that *Bt* toxin would be broken down in the stomach. However, in a 2011 Canadian study conducted at Sherbrooke Hospital, researchers discovered *Bt* toxin in the blood of 93% of the

pregnant women and 69% of non-pregnant women.⁴² A mouse study confirmed that *Bt* toxin is cytotoxic,⁴³ it therefore might also damage human blood cells. Since fetuses do not have fully developed blood brain barriers, it is possible that the toxin reaches the fetal brain.

In 2001, the EPA's Scientific Advisory Panel, which included leading experts in the US, pointed to these early mouse and farmworker studies and stated that they "suggest that *Bt* proteins could act as antigenic and allergenic sources."⁴⁴ The EPA disregarded the warning, reregistered the *Bt* crops, and continues to claim that *Bt* toxin has no impact on humans or mammals.

They also claim that the *Bt* toxin engineered into plants is the same as that which was sprayed. However, industry submissions and published papers establish that the genetically engineered *Bt* toxin in plants is structurally different⁴⁵ from the natural *Bt* toxin used in spray applications to crops. Whereas the spray version creates a protoxin that is fully activated after entering the alkaline environment of the insect's gut, the plant version is designed to be immediately toxic. The genetically engineered *Bt* toxin (the plant version) has properties of known allergens, fails the World Health Organization's allergenicity decision tree criteria⁴⁶ and is produced in concentrations thousands of times higher than the spray version. Most notably, while the spray version can be washed off the plant and biodegrades quickly in sunlight, the plant version is encapsulated within the plant cells, remains intact, and cannot be removed by washing.

Increased herbicide use and residues on food

Herbicide-tolerant crops comprise 89% of all US grown GMOs. These plants are engineered to allow specific herbicides to be sprayed in high amounts without damaging the GM plant. Although GMO companies had publicly predicted GMOs would reduce herbicide use, the opposite has occurred—pesticide chemical use has increased as a result of GMO crop use. In fact, overuse of these herbicides has resulted in "superweeds," which have developed resistance to the herbicide. Farmers often spray higher quantities of the herbicides to kill these "superweeds." According to Benbrook, statistics from the United States Department of Agriculture (USDA) reveal that herbicide tolerant crops led to an increase in herbicide use of 527 million pounds over the first 16 years.⁴⁷ Use of Roundup® and other GBHs has increased 100-fold since the late 1970s.^{48,49} The allowable glyphosate residues on GMO crops have also increased substantially, as government regulations have been relaxed to allow higher use of pesticides to accommodate industry goals.⁵⁰

A 2013 paper in the journal *Entropy* examined the potential effects of glyphosate ingestion. Examining the biochemical impacts on two key metabolic pathways, as well as its broad-spectrum chelating effects, the authors speculate potential causal mechanisms that link it to "most

of the diseases and conditions associated with a Western diet, which include gastrointestinal disorders, obesity, diabetes, heart disease, depression, autism, infertility, cancer and Alzheimer's disease."⁵¹ (Emphasis added.) Although interesting, it remains to be demonstrated whether the suggested association between glyphosate ingestion and this very wide range of human disease takes place at real-world levels of exposure.

Numerous studies in the past several years, ranging from in vivo and in vitro, to occupational exposure analyses, have implicated Roundup®, or its active ingredient glyphosate, in cancer, birth defects, endocrine disorders, Parkinson's, and damage to beneficial gut bacteria.⁵²

In order to postulate how applications of Roundup® or other GBHs and residues on GMO crops might cause or exacerbate specific disorders, it is necessary to identify glyphosate's possible modes of action in the body.

Glyphosate as carcinogen

The International Agency for Research on Cancer (IARC) of the World Health Organization, which is responsible for classifying chemicals as carcinogens, classified glyphosate and glyphosate-based herbicides as class 2A carcinogens—"Probably carcinogenic." They confirmed that glyphosate causes cancer in animals, it creates mutations in human DNA, and where it is sprayed, there have been spikes in cancer among the exposed populations. IARC also determined that glyphosate is genotoxic and creates oxidative stress.⁵³

Glyphosate as chelator

Glyphosate, the active ingredient in Roundup®, was patented as a powerful mineral chelator in 1964, a decade before Monsanto patented it as an herbicide. It binds with cations, including zinc, manganese, cobalt, aluminum, calcium, magnesium, arsenic, iron, selenium, cobalt, chromium and arsenic. Living organisms—plants and animals (including humans)—rely on minerals for numerous metabolic pathways to function. When bound with glyphosate, the minerals cannot be utilized.⁵⁴ Glyphosate applications may therefore result in symptoms of mineral deficiency, even though the minerals are present, because the bound ("chelated") mineral is not biologically available to perform its biochemical and physiological functions. Whether glyphosate disturbs nutrient mineral homeostasis at real world levels of ingestion that could then result in disease remains an urgent factor to be investigated.

Glyphosate inhibits critical metabolic pathways

Monsanto has long described glyphosate's herbicidal mode of action as its ability to block plants' shikimate pathway. They identify the mechanism as direct inhibition of excitatory postsynaptic currents (EPSPs) by binding to

the active site of that enzyme. Other researchers identify glyphosate's tendency to strongly bind with cobalt and manganese as the mechanism for inhibiting the shikimate pathway. In either case, this shikimate pathway produces the aromatic amino acids—tryptophan, tyrosine, and phenylalanine, which are needed for plant survival.⁵⁵

Monsanto has claimed that glyphosate is safe for humans and other mammals because they lack the shikimate pathway. However, critical gut bacteria in humans and mammals also possess the shikimate pathway to produce these essential aromatic amino acids, which are the building blocks for producing proteins.

The production of serotonin, for example, requires tryptophan, which is a product of the shikimate pathway. There is insufficient research to determine whether the amount of tryptophan produced by gut bacteria contributes significantly to the production of serotonin. If so, when there is insufficient tryptophan in the food, glyphosate's suppression of the shikimate pathway may reduce overall serotonin levels.

Serotonin might also be impacted in other ways. Up to 90% of the serotonin in the human body is produced in the gut by enterochromaffin (EC) cells. It is now well established that a large proportion of the serotonin produced by EC cells is dependent on stimulation from certain gut bacteria.⁵⁶ Gut bacterial dysbiosis resulting from sufficient amounts of glyphosate ingestion could therefore give rise to inadequate amounts of serotonin production by EC cells. This could lead to health and behavioral problems.

No studies have evaluated the impact of glyphosate on gut-produced and microbe-produced serotonin.

Regardless, the safety determination must not ignore the fact that other metabolic pathways could also be disrupted by glyphosate's ability to bind with nutrient metals. Minerals are critical components in numerous pathways acting as vital elements of many enzymatic functions, and glyphosate's ability to chelate is potent and thus potentially damaging to health.

Glyphosate as antibiotic

Glyphosate was patented as a broad-spectrum biocide, which preferentially kills the types of bacteria that are considered beneficial in the gut of humans and animals. For example, *Lactobacillus* and *Bifidobacteria* are particularly sensitive to glyphosate and are killed in the presence of even low concentrations. Unfortunately, bacteria considered potentially pathogenic in the human gut are more resistant to glyphosate including *Clostridium difficile*, *Escherichia coli*, and *Salmonella*.⁵⁷ Studies are needed to evaluate the extent of antibiotic activity that occurs in the human gut due to real-world levels of glyphosate exposure.

Glyphosate as mitochondrial toxin

Glyphosate can disturb vital functions of the mitochondria. This may be due in part to its ability to bind with manganese, which is needed by the mitochondria for proper functioning. Further research is needed to establish whether the amount of glyphosate residues in food are sufficient to contribute to mitochondrial dysfunction in humans.⁵⁸

Glyphosate as teratogen

Glyphosate and GBHs can interfere with the retinoic acid pathway in fetal development. Studies have shown teratogenic effects.⁵⁹ However, these studies used high, unrealistic doses, so further studies using realistic doses are needed.

Glyphosate as endocrine disruptor and estrogen mimicker

Several studies have shown that glyphosate influences hormones and can act as an endocrine disruptor. For the most part, these studies use animal models and human cells with levels of glyphosate that are higher than that which is expected to be consumed as residues on food. Of course, many endocrine disrupting chemicals can exert an influence—even a greater influence—at minute levels. There are no regulatory requirements to test these low levels because government policies have not kept pace with the body of research on low-dose impacts. Some research of note, using both small and high dosages, includes:

- Glyphosate can interfere with the action of aromatase, which determines the ratio of estrogen and testosterone.⁶⁰
- Rats fed Roundup® (R) in the drinking water over 24 months, and also those fed GMO RR corn, showed changes in their sex hormones. “In females, the androgen/estrogen balance in serum was modified by GM maize and Roundup® treatments... For male animals at the highest R treatment dose, levels of estrogens were more than doubled.”⁶¹
- Adjuvants or surfactants used with herbicides are generally considered inert, but the “inert” ingredient in the full formulation of Roundup® can also exert low dose hormonal effects. According to Defarge, et al., “Aromatase activity was decreased both by the co-formulants alone (polyethoxylated tallow amine-POEA and alkyl polyglucoside-APG) and by the formulations, from concentrations 800 times lower than the agricultural dilutions; while G [glyphosate] exerted an effect only at 1/3 of the agricultural dilution. *It was demonstrated for the first time that endocrine disruption by GBH could not only be due to the declared active ingredient but also to co-formulants.*” (Emphasis added.)
- Seralini observed hormonal effects observed in rats fed GBHs, but it is unclear whether the impacts were due to the glyphosate, the adjuvants, the GMO

transformation process, or a combination of them all.⁶²

- At parts per billion, glyphosate attached to estrogen receptors in human breast cancer cells and triggered growth,⁶³ although a follow-up study was unable to replicate this result.⁶⁴

Overlapping and synergistic factors

A two-year feeding study highlights how negative impacts of GMOs may be due to both the GMO process and the added herbicide.⁶⁵ The researchers fed rats RR corn that had been sprayed with Roundup®, RR corn without Roundup®, or Roundup® without the corn. The Roundup® alone was tested at a range of doses, including a very low dose of a level that would be permitted in drinking water. Animals in all three treatment categories—GMO alone, Roundup® alone, and GMO corn with Roundup®—suffered organ damage, especially to the liver and kidney but also to the pituitary gland. Increased tumor incidence and mortality were observed in most treatment groups. In particular, a statistically significant increase in mammary tumor incidence occurred in the lowest dose Roundup® group. In this study, it was clear that compared to controls, rats fed GMO RR corn, Roundup®, and the combination of the two, were all harmed. Most animal studies that test GM crops do not include multiple feeding groups, making it unclear whether the causative factor is the GMO, the *Bt* toxin, or the Roundup®.

In the study above, a follow up molecular analysis (transcriptome – gene function profile) of liver and kidney tissues clearly confirmed the damage for the lowest dose Roundup®-fed group.⁶⁶ A further molecular analysis involving proteomics (protein type profile) and metabolomics (metabolite profile) found that this lowest-dose Roundup®-fed group suffered from non-alcoholic fatty liver disease (NAFLD),⁶⁷ an increasingly common condition that can lead to a more serious disease, non-alcoholic steatohepatitis, and ultimately to cirrhosis. NAFLD is also a risk factor for liver cancer. This study has identified long-term low-dose exposure to Roundup® as a previously unrecognized risk factor for NAFLD and its associated complications.

Several animal studies show health impacts

The online publication *GMO Myths and Truths* provides an excellent summary of the research results on GMOs, using laboratory and farm animals. The publication also counters several arguments proposed by GMO advocates that attempt to downplay the findings. The categories of problems discovered, as listed below, are from studies⁶⁸ conducted both by independent researchers and by GM industry employees or contractors.

- Severe organ damage and increased rates of large tumors and mortality⁶⁹
- Altered blood biochemistry, multiple organ damage, and potential effects on male fertility⁷⁰

- Stomach lesions and unexplained mortality⁷¹
- Immune response and allergic reaction⁷²
- Immune disturbances⁷³
- Enlarged lymph nodes and immune disturbances⁷⁴
- Disturbed liver, pancreas and testes function⁷⁵
- Liver aging⁷⁶
- Disturbed enzyme functioning in kidney and heart⁷⁷
- Higher density of uterine lining⁷⁸
- Severe stomach inflammation and heavier uteri⁷⁹
- Liver and kidney toxicity⁸⁰
- Changed level of fats in blood and signs of liver and kidney toxicity⁸¹
- Toxic effects on liver and kidneys and altered blood biochemistry⁸²
- Enlarged liver⁸³
- Disturbances in digestive system and changes to liver and pancreas⁸⁴
- Excessive growth in the lining of the gut⁸⁵
- Intestinal abnormalities⁸⁶
- Altered blood biochemistry and gut bacteria, and immune response⁸⁷
- Altered gut bacteria and organ weights⁸⁸
- Less efficient feed utilization and digestive disturbance⁸⁹

Consumer exposure

Consumers are exposed to GM foods and GM DNA through consumption of the whole food crop, derivatives of the crop, and to a lesser extent through milk, meat, and eggs from animals that have been fed GM feed. GMO crops can be eaten raw (e.g. GMO papaya and zucchini) or cooked (e.g. edamame, corn, corn chips, tortillas, zucchini, squash, etc.). Therefore, any legitimate safety testing must include both raw and cooked samples, including samples cooked in a variety of real-world means including baking and frying.

GMO product derivatives include highly processed components, such as sugar, corn sweeteners, soy protein-based products and the oils from soy, corn, cottonseed, and canola. For these, the DNA and proteins derived from the transgene are often removed during processing. While processing may reduce the health risk, it does not ensure that GMO derivatives are always as safe as their non-GMO counterparts. The composition of these processed ingredients may be altered due to the changes that take place in the plant as a result of the disruptive GMO transformation process. This could result in the presence of novel toxins and allergenic substances.

GMO DNA has been detected in milk from animals fed GMOs.⁹⁰ It is not clear whether the *Bt* toxin consumed by the animals continues to be intact after consumption and digestion, and if it will be active in meat or dairy products. According to a study by Aris and Leblanc, “there may be a high risk of exposure through consumption of contaminated meat.”⁹¹ They cite studies showing that “trace amounts of the Cry1Ab toxin were

detected in the gastrointestinal contents of livestock fed on GM corn."⁹²

Glyphosate residues are found in food from RR crops sprayed with a GBH. In 1999, Monsanto claimed that residue levels of up to 5.6 mg/kg in GM-soy represent "...extreme levels, and far higher than those typically found."⁹³ But authors of a 2014 study demonstrated that "Seven out of the 10 GM-soy samples we tested, however, surpassed this 'extreme level'" (of glyphosate [plus its derivative AMPA]).⁹⁴ Their average residue level was 9.0 mg/kg.

Glyphosate is water soluble; therefore it is unlikely that soybean *oil* (rather than whole soy) contains residues. According to a report from China, however, tests verified the presence of aminomethylphosphonic acid (AMPA), the primary breakdown derivative of glyphosate in soybean oil.⁹⁵ AMPA exhibits some toxic properties and has a similar structure and profile to glyphosate.⁹⁶ Another study by Böhn et al., showed significant levels of glyphosate residues in GMO soybeans.⁹⁷

According to Monsanto's radiolabel studies submitted to the EPA and later obtained by Anthony Samsel, glyphosate is found in animal tissues.⁹⁸ A study using an ELISA detection method found glyphosate residues in breast milk,⁹⁹ while two studies using mass spec did not.¹⁰⁰

If GMOs and GBHs have detrimental effects, it is logical to conclude that animals fed a diet almost exclusively of GMOs would be measurably different than animals raised on diets free from GMOs. According to Heinemann's extensive review of the scientific literature, studies reveal the presence of "DNA and protein unique to GM plants within animals and animal products." Furthermore, "There is compelling evidence that animals provided with feed containing GM ingredients can react in a way that is unique to an exposure to GM plants. This is revealed through metabolic, physiological or immunological responses in exposed animals."¹⁰¹ No studies have been conducted to determine how these differences could exert an influence on the health of humans consuming the animal products.

The consumption of residues of GBHs is not limited to RR crops. It is not uncommon for some non-GM crops, such as wheat, barley, rice, wine grapes, sunflowers, rye, oats, and sugarcane, to be given a pre-harvest application of GBH as a desiccant to dry the crop, accelerate ripening/maturation (as the plant dies), and/or kill weeds. To accommodate this practice, the EPA raised the allowable levels of glyphosate residues on more than 160 crops.¹⁰² The actual amount of glyphosate residues in food, however, is not yet measured by the FDA, even though they monitor levels of all other commonly used pesticides. The FDA announced plans to monitor glyphosate residues, but the program was suspended and then later declared active once again. However, we do not yet have a measure of human glyphosate exposure in the

US. According to independent food testing, common US foods ranged from a low of 8.02 ppb in Goldfish Crackers Colors by Pepperidge Farm, to 1,125.3 ppb in Original Cheerios by General Mills.¹⁰³ The EPA asserts that these levels are below permitted residue levels and thus pose no health concerns for consumers. However, others challenge that level as unscientific. For example, the amount of Roundup® consumed by rats in their drinking water that caused non-alcoholic fatty liver was 0.1 ppb containing 0.05ppb of glyphosate. On a per body weight basis the rats ingested 4 nanograms (4 thousand millionths of a gram) of glyphosate per kilogram body weight. This is 437,500 times lower than US permitted levels.¹⁰⁴ And based on glyphosate levels detected in US citizens, Americans probably consume about 1000-fold more glyphosate than the amount responsible for the liver disease in rats.¹⁰⁵ Thus it is possible that ingestion of glyphosate from foods at the residue levels detected could cause harm over the long term.

Organic crops are not allowed to be sprayed with GBH. Therefore, if GBH is one of the causative factors in the health problems reported, we would expect better outcomes for those who switch to organic, rather than those who switch to non-GMO foods that are still sprayed.

The main source of exposure to glyphosate is through food. However, secondary sources include air, rain, water, and drift from agricultural and homeowner use.¹⁰⁶

Part 2: Survey Results

Methods

Survey participation was requested from November 2014 through August 2015. The questionnaire was emailed to 180,716 members of the database of the Institute for Responsible Technology (IRT). There were 3,256 responses—a response rate of 1.8%.

Reporting bias

This is a self-selecting survey of a non-representative sample of the population. IRT is a leading advocacy group that educates people on the health dangers of GMOs. The results of this survey are therefore limited to a population that is already aware of GM crops and has been exposed to information about the negative health impacts. Some percentage of the respondents may be biased towards attributing health improvements to the elimination of GMOs based on expectations. On the other hand, this population will have a higher percentage of people who have become educated about GM food risks, eliminated them, and may have noticed a change as a result.

There is also an expected bias on a per-disease basis. People will more likely identify an actual connection between their diet and a chronic condition if that condition is normally associated with a dietary response. Gastrointestinal disorders and food allergies are more likely to be considered and evaluated in terms of

reaction to diet compared to cancer, for example, or infertility. Furthermore, the more serious diseases, once discovered, are often treated with pharmaceuticals, surgery, and other treatments that might overshadow or mask the impact of the diet.

The Questionnaire

Based on the first question in the survey, respondents offered results from dietary changes for themselves (80%), their spouse (4.6%), their child (4.2%), and their patient (1%). In addition, 2.2% provided information on their pet, 3% on their livestock, and 7% on “other,” but these were not tallied in the results below. Most respondents did not choose to indicate their zip code or location, but of the 1,870 who did, 1,620 were in the US and 98 in Canada. Thus, the vast majority of respondents were located in the US, which is consistent with the demographics of the email list used to solicit responses.

The survey was designed to identify the relative frequency of conditions that improved with a non-GMO diet, and the degree of reported improvement. It was not intended to identify what percentage of the population would improve on a non-GMO diet.

The survey was composed of six questions. The second question was, “What symptoms or conditions have you seen improve since switching to a non-GMO Diet?” Response choices for each of 28 symptoms or conditions were as follows:

1. Some Mild Improvement
2. Moderate Improvement
3. Significant Improvement
4. Nearly Gone
5. Complete Recovery
6. N/A Not Applicable*

*“Not applicable” was the pre-checked default response, required by the survey system used.

Table 1 shows the percentages of respondents who indicated any improvements, 1-5.

Competing co-factors

There are no laws requiring GMO foods to be labeled as such in the US. Avoiding them, therefore, usually involves a strategy that can introduce other possible co-factors that may be responsible for the reported improved health outcomes. Because organic foods are not allowed to intentionally contain GMOs, switching to organic foods is a popular strategy to avoid GMOs. However, numerous toxic chemicals are also considered prohibited substances according to the organic standards. Thus, improvements in health may be due in part or in full to the elimination of these other products.

Most GMOs are found in processed foods. This is because derivatives of soy, corn, cottonseed, canola, and sugar beets are common ingredients in processed foods. Many people choose to avoid GMOs by reducing consumption of processed foods. Therefore, health

improvements may be related to the benefits of unprocessed foods.

Some healthcare practitioners recommend elimination of GMOs along with other dietary instructions at the same time, such as eliminating gluten or dairy, which may contribute to or account for the health improvements.

The questionnaire included the question: “In addition to eliminating GMOs from your diet, were there other changes that you made at the same time or thereafter? Check all that apply.” The number and percentage of respondents that checked each change shown in Table 2.

Table 1. Survey Question #2 Responses

Health Condition Improved	% Reported
Digestive problems	85.2%
Fatigue	60.4%
Overweight or obesity	54.6%
Clouding of consciousness (brain fog)	51.7%
Mood problems/anxiety/depression	51.1%
Food allergies or sensitivities	50.2%
Memory and concentration	48.1%
Joint pain	47.5%
Seasonal allergies	46.6%
Gluten sensitivities	42.2%
Insomnia	33.2%
Other skin conditions	30.9%
Hormonal problems	30.4%
Musculoskeletal pain	25.2%
Autoimmune disease	21.4%
Eczema	20.8%
Cardiovascular problems and high blood pressure	19.8%
Asthma	14.8%
Menstrual problems	13.1%
Diabetes	10.6%
Other mental disorders	7.9%
Underweight	6.5%
Cancer	4.8%
Kidney disease	4.5%
Infertility	3.8%
Autism spectrum	2.6%
Alzheimer’s disease	2.4%
Parkinson’s disease	1.4%

Table 2. Survey Question #4 Responses

Dietary Change	Number	Percent
Organic diet	2,424	74.4%
Reduced processed foods	2,182	67.0%
Stopped drinking soda, or other sweetened beverages	1,503	46.2%
Gluten-free	1,099	33.8%

Eliminated dairy products	755	23.2%
Raw	381	11.7%
Vegetarian	372	11.4%
Vegan	228	7.0%
No other changes	226	6.9%

Without carefully controlled human clinical trials, it may be impossible to assess how much, if any, the GMO component of the diet is causing health problems. The following evidence does, however, support the notion that GMOs (and their associated pesticide content) are a contributor to health conditions:

1. Most of the reported improvements in humans correlate with the categories of health impacts of GMOs, glyphosate, and GBH on animals in carefully controlled feeding trials, which exclude other confounding dietary factors.
2. Based on informal surveys and conversations referred to above, farmers and veterinarians describe improvements in livestock that are switched to non-GMO soy, corn, or both. In livestock, there are generally no other dietary changes and the reported improvements, e.g. gastrointestinal, immune, irritable or aggressive behavior, fatigue level, skin health, etc. are similar or identical to those reported by individuals and their practitioners.
3. The categories of reported improvements also correlate with many of the diseases and conditions that increased in parallel with the expanded exposure of GMOs and their associated herbicides in the US population (See Figures 2 through 5).
4. The reporting of numerous health improvements was consistent across various dietary strategies. For example, gastrointestinal health improved for a large percentage of respondents, irrespective of whether their strategies were to switch to organic, reduce processed foods, or favor brands labeled non-GMO, etc. The non-GMO component was common to all.
5. The characteristics of GMOs, and the agricultural toxins found in the foods, can plausibly explain the conditions linked to their consumption.

Part 3: Focus on Digestive Disorders

No survey can, on its own, demonstrate causality. This survey does, however, provide data that can be analyzed in conjunction with other evidence to support the argument that GMOs promote particular disorders. Below we examine additional data that may explain or highlight a causal relationship between GMO consumption and the most frequently cited health improvement in the survey—digestive problems.

With 85.2% of survey respondents reporting mild to total recovery, digestive issues are by far the number one category of self-reported benefits from a non-GMO diet. The breakdown of responses is as follows: Some

Mild improvement (5.9%), Moderate Improvement (11.3%), Significant Improvement (29.1%), Nearly Gone (22.2%), Complete Recovery (16.6%).

Digestive disorders in general include inflammatory bowel, Crohn's disease, irritable bowel, acid reflux, diarrhea, constipation, ulcerative colitis, bloating, and gas.

According to CDC data, incidence of inflammatory bowel disease (IBD), Crohn's, and ulcerative colitis all rose in parallel with the percentage of GMO corn acreage planted in the US since 1996 and the amount of GBH sprayed on GMO corn and soy acreage. The corresponding graphs shown in Figures 2 through 5 are reproduced below with permission from Nancy Swanson.

Figure 2. IBD diagnoses and acres of Bt corn¹⁰⁷




Figure 3. Function Bowel disorders and acres of Bt corn¹⁰⁸




Figure 4. IBD and glyphosate applied to corn and soy: Correlation between inflammatory bowel disease and glyphosate applications to US corn and soy crops.¹⁰⁹





Figure 5. Deaths due to intestinal infection and glyphosate applied to corn and soy: Correlation between age-adjusted intestinal infection deaths and glyphosate applications to US corn and soy crops.¹¹⁰



Digestion and GMOs

The digestive tract is the first and largest point of contact with food. According to Arpad Pusztai, who was commissioned by the UK government to create a protocol for testing the safety of GMOs, the digestive tract can reveal numerous reactions to consumed toxins and should therefore be the first subject of a GMO food risk assessment. In his research using rats, Dr. Pusztai discovered that the stomach and intestinal lining of these animals fed GMO potatoes showed altered architecture and potentially precancerous cell growth compared to controls. His study design further revealed that the changes were not due to the protein expressed from the transgene, implicating the generic process of the GMO transformation process as the cause.¹¹¹

In another study on rats fed GMO peas, the rat intestines were heavier, possibly indicating excessive cell growth (a hallmark of cancer), but researchers ultimately failed to examine the intestines for cell growth.¹¹²

The FlavrSavr tomato, engineered for delayed ripening and rotting, was approved and marketed in the US in 1994 but quickly withdrawn by 1997. Scientists conducted rat feeding studies on two lines of the tomato. In the line that was not commercialized, 7 of 20 rats developed stomach lesions. According to Pusztai, the type of stomach lesions linked to the tomatoes “could lead to life-endangering hemorrhage, particularly in the elderly who use aspirin to prevent [blood clots].”¹¹³ Studies on the FlavrSavr did not look beyond the stomach to evaluate other possible impact on the intestines.

A study by Carman and colleagues found that pigs fed typical transgenic corn and soy-containing feed in the US had significantly higher incidence of severe inflammation of the stomach than controls. In fact, compared to a non-GMO diet, the stomachs of 32% of the GMO-fed pigs were scored in a category of severe inflammation—a classification that was above and beyond any of the inflammation ratings for animal feeds a non-GMO diet.¹¹⁴

With the exception of Pusztai’s rat studies, it is difficult to tell whether the cause of the digestive problems seen in animal studies was due to the GMO transgene product itself, the added herbicide, or the disruption of plant host gene function resulting in novel toxins or allergens.

In a study conducted by Monsanto, GMO soy was found to contain 27% more trypsin inhibitor than its natural isogenic counterpart.¹¹⁵ Additional data from Monsanto’s study that was not published, was later discovered by medical writer, Barbara Keeler, in the archives of the *Journal of Nutrition*. It demonstrated that the trypsin inhibitor in GMO soy was resistant to denaturing from heat. After cooking GMO soy meal twice, according to a review by Pusztai and Bardocz, “one of the soybean lines (61-67-1) appeared to have almost seven times as much trypsin inhibitor per mg sample dry weight as the parent. Indeed, the values of this GM soybean approached that found in untoasted seed samples. Even the other GM line (40-3-2) contained three times as much trypsin inhibitor as the non-GM line.” They concluded “heat treatment appeared to have a far lesser denaturing effect on the trypsin inhibitor content of the GM lines.”¹¹⁶

Trypsin, a pancreatic protease, catalyzes the hydrolysis of proteins into smaller peptides for digestion and reduces the allergenicity of the proteins. By blocking the effects of trypsin, trypsin inhibitors can inhibit the digestion of proteins and enhance the allergenic properties of proteins.

In addition, the toasted GM soy meal contained nearly twice the amount of a lectin, which may interfere with assimilation of nutrients.¹¹⁷

The pancreas of mice fed RR soy exhibited profound changes.¹¹⁸ Starting in month two, production of alpha-amylase, a major pancreatic enzyme that degrades carbohydrates, dropped by an average of 77%. In months

five and eight, it was 75% and 60% lower than controls. This reduced production of alpha-amylase was confirmed in the rough endoplasmic reticulum, the Golgi apparatus, and within zymogen granules.

One-month-old GMO-fed mice produced less zymogens than those fed a non-GMO diet, but the differences became negligible as they aged. The size of these granules was consistently smaller in GMO-fed mice, with the biggest difference being 39%, in month five. The pancreatic modifications disappear after removing the GMO soy from the diet.

Digestion and Bt toxin

The possible role of *Bt* toxin as a causative agent in digestive disorders is not difficult to extrapolate. Its mode of action as an insecticide is to kill insects by creating holes in the walls of their guts. Studies on human embryonic kidney cells demonstrate that *Bt* toxin creates similar micropores.¹¹⁹ Increased intestinal permeability in humans is known to be causally associated with a variety of autoimmune, inflammatory, allergic and pain-related conditions.

While the amount of *Bt* toxin required to cause micropores in human cell cultures in dishes was higher than the amount of toxin that is supposed to be produced in *Bt* corn and thus ingested, we cannot discount the potential for *Bt* toxin to cause holes in the human digestive tract for a number of reasons:

1. The environment within the stomach may be quite different than the lab simulation and the type of cells used in the experiment are not those found in the gut. The quantity needed to disrupt the cell integrity may be less (or more) in vivo.
2. The amount of *Bt* toxin produced by corn can vary, depending on environmental and other factors. According to an investigation by Professor Terje Traavik on *Bt* corn in the Philippines, the expression of *Bt* toxin in one single corn cobb varied per kernel up to 64-fold. The range may have actually been larger, but the amount was measured at the limit of detection. This particular corn, however, may have also been atypically unstable. It was associated with a strong smell and health conditions that afflicted the nearly 100 people living adjacent to the cornfield.¹²⁰
3. The *Bt* toxin gene may transfer to the DNA of bacteria living within the gut. Although this was never studied for *Bt* corn, research on RR soybeans confirmed that part of the RR gene, including its viral promoter, transferred and integrated into the DNA of gut bacteria of ileostomy subjects. Although the transformed bacteria survived exposure to glyphosate, it is not clear whether this was due to the intact functionality of the transgene or to the bacteria's natural immunity to this antibiotic. If the *Bt* toxin gene transferred and continued to function from within gut bacteria, the amount of the toxin produced by the gut

flora could well exceed the level produced in corn. Further, the exposure could be constant, 24 hours per day. Unfortunately, this is hypothetical because no studies have investigated this potential risk.

As discussed above, numerous studies of *Bt* toxin elicit an immune response. Histamine is the major paracrine stimulant of gastric acid. Thus, elevated immune reactivity might contribute to digestive disorders through histamine production. For example, histamine is involved in the secretion of gastric acid.¹²¹

Mice fed potatoes engineered to produce the *Bt* toxin developed abnormal and damaged microvilli, as well as proliferative cell growth in the lower part of their small intestines (ileum).¹²²

Digestion and Glyphosate

A study on fairly high levels of Roundup® exposure in carnivorous fish revealed remarkable adverse effects throughout the digestive system, including “disruption of mucosal folds and disarray of microvilli structure” in the intestinal wall, along with an exaggerated secretion of mucin throughout the alimentary tract.¹²³

Reduced digestive enzymes

Although the relevance of fish model to human health is limited, it is remarkable that the activity of protease, lipase, and amylase, important enzymes involved in the digestion of proteins, fats, and carbohydrates, were all decreased in the esophagus, stomach, and intestine of these fish following exposure to glyphosate.¹²⁴

Enzymes secreted by the pancreas are responsible for the breakdown of food so that it can be absorbed through the walls of the small intestine into the bloodstream. Any restriction of these enzymes may result in impaired digestion and a shortfall of nutrient assimilation. If carbohydrates are not properly degraded in the small intestine, (as may occur with reduced alpha-amylase), they may be broken down by bacteria in the large intestine, which can produce gas. If protein digestion is inhibited, which may occur with reduced zymogens, it can increase the chance of allergic reactions to protein fragments. The pancreas may also be forced to produce and excrete more protein digesting enzymes, possibly putting undue pressure on the organ.

Ultimately, if the digestive system is not functioning properly, then food particles are not broken down as quickly or as completely. This can create several problems:

- **Nutritional deficiencies:** If a person is not properly absorbing and gaining sufficient nutrition from the foods consumed, overall health, including the immune system, can suffer.
- **Dysbiosis:** With poor digestion, proteins can remain intact for longer than normal periods in the gastrointestinal (GI) tract. This can result in the larger, undigested food particles becoming the

“food” of pathogenic gut bacteria, leading to overgrowth, which can further compromise digestion and immunity.

- **Inflammation:** When proteins putrefy, they can also release excess hydrogen sulfide (as toxic as cyanide gas) which irritates and inflames the mucous membranes.
- **Autoimmunity:** Undigested proteins also have a greater likelihood of provoking autoimmune reactions, in which the immune system attacks parts of the body.

Intestinal inflammation also appears to reduce production of cholecystokinin (CCK) and this reduction in CCK, in turn, reduces the digestive enzymes produced by the pancreas, as well as the bile produced in the liver. Without sufficient enzyme levels, digestion is slowed down, particularly digestion of proteins; without sufficient bile, fat and fat-soluble vitamins cannot be digested and absorbed efficiently.

This can become a vicious cycle: Larger food particles can result in bacterial overgrowth, which in turn can further irritate the lining of the intestines, further lowering digestive capacity both directly and through reduced CCK levels. Lowered digestive capacity results in increased large food particles.

Glyphosate altering gut bacteria

Glyphosate has been shown to reduce the population of healthy bacterial varieties in the digestive tract and promote overgrowth of dangerous pathogenic bacteria, according to in vitro research with poultry¹²⁵ and cattle models.¹²⁶

The implications for health may be quite profound and complex. For example, *Bifidobacterium* strains, which are often used as probiotics, reduce the cytokines that provoke inflammation. *Bifidobacterium* is one of the types of bacteria that are easily killed by glyphosate.¹²⁷ The result could be an increase in inflammation, which is now recognized as central to the disease process for numerous diseases.

On the other hand, “the highly pathogenic bacteria,” such as those that produce *Salmonella* or *Clostridium botulinum* (the botulism toxin), “are highly resistant to glyphosate.” Furthermore, some of the beneficial bacteria that are killed normally keep some of the pathogenic bacteria population in check. Researchers in Germany, for example, suggest that glyphosate use kills lactic-acid producing bacteria in the gut of cattle, allowing the bacteria that produce deadly botulism to flourish. This might explain the increase in chronic botulism in cattle.¹²⁸ Cases of Sudden Infant Death Syndrome have also been linked to the botulism toxin.¹²⁹

Bacterial pathogens can activate zonulin, a protein that modulates the permeability of the tight junctions between cells of the wall of the digestive tract. Activation of zonulin can induce a breakdown of the tight

junctions in cells lining the gut, leading to increased intestinal permeability or “leaky gut”.¹³⁰ Indeed, some of the same bacteria whose growth is stimulated through glyphosate exposure, i.e. *Clostridium botulinum*,¹³¹ *Clostridium perfringens*,¹³² and *Salmonella*,¹³³ have been found to provoke diseases in humans; these are not benign bacteria.

An experimental study using two intestinal cell lines showed that glyphosate could adversely affect mucosal barrier integrity. The authors concluded that at higher doses “glyphosate significantly disrupts the barrier properties of cultured intestinal cells.”¹³⁴

By inhibiting the Shikimate pathway, glyphosate might reduce the production of tryptophan and serotonin. Because serotonin is important for intestinal motility, a deficiency could have consequences for digestive health. According to a review article by Sikander, et al., “Altered serotonin signaling may lead to both intestinal and extra intestinal systems in IBS [irritable bowel syndrome].”¹³⁵

Digestive disorders as gateways to other conditions

Digestive disorders not only create symptoms themselves, they can act as causative factors for other conditions. Lack of proper breakdown and assimilation of nutrients can lead to nutritional deficiencies, which can result in a myriad of health problems. A hyperpermeable gut is linked to numerous inflammatory and metabolic disorders, ranging from allergy to depression and autoimmunity. Altered gut bacteria can impact detoxification, immune function, and the availability of key nutrients. Therefore, many of the other symptoms listed in the survey may have originated in the gut.

Conclusion

GMOs are pervasive in the diet of people living in the US and several other nations. Although presumed safe or GRAS by the US government and GMO producers, published studies point to numerous physiological responses in animals and cell lines that challenge this assumption. They demonstrate changes or even damage to the immune system, reproductive system, vital organs (especially liver and kidney), digestive system, and endocrine system.

Survey results of 3,256 people reporting improvements in at least some health conditions, after switching to a non-GMO, diet suggest GMOs may be contributing to health conditions. Many of the conditions that improved in the survey participants are similar to the health issues found in lab animals fed GMOs or the associated herbicide Roundup®. Other dietary factors, such as increased consumption of organic food, reduction of processed food, etc., may also play a role in the health improvements. Thus, future research should exclude confounding factors as much as possible.

Digestive issues were by far the most common problem reported by respondents as improved when

GMOs were removed from the diet. GMOs can negatively impact digestion through several possible modes of action and digestive disorders in turn, can lead to numerous other health issues.

Future research is also warranted to clarify if the generic GMO transformation process, *Bt* toxin, and/or the glyphosate/GBH residues are contributing to or causing health problems and to definitively determine the causative pathways of potential harm in the human body.

It is clear that more research needs to be done. However, given the prevalence of data correlating GMOs and glyphosate/Roundup® with health issues, and evidence that a switch to non-GMO organic diets contributes to recovery (improved health), the author believes the precautionary principle dictates that healthcare practitioners should advise patients to avoid exposure by switching to organic foods. Furthermore, practitioners are encouraged to document the impacts and publish case studies.

Acknowledgements

The author (JMS) would like to acknowledge the many people who contributed to the article, as editor, scientific or medical advisor, content contributor, contributor to the survey design, survey manager, etc. These include:

- Sara Jennings
- Nancy Swanson PhD
- Stephanie Seneff PhD
- Sayer Ji
- Stanley Ewen
- Tom O'Bryan DC
- Michelle Perro MD
- Alex Vasquez DC ND DO
- Terri Ward MS NTP CGP
- Don Huber PhD
- Anthony Samsel PhD
- and others



About the Author:

Jeffrey M. Smith is the Executive Director of the Institute for Responsible Technology (IRT), a US-based nonprofit that educates policy makers, media, healthcare practitioners, and the public about the health risks of GMOs and their associated pesticides. IRT also exposes the unscientific methods used by the promoters of GMOs to hide evidence of harm. Mr. Smith is the author of *Genetic Roulette: The Documented Health Risks of Genetically Engineered Foods and Seeds of Deception*. He is director of the film *Genetic Roulette—The Gamble of Our Lives*, and co-director of the upcoming film *Secret Ingredients*, about families and individuals who recover from significant health conditions after switching to non-GMO organic food.

**See associated Editorial
and Comments attached
with this article**

Footnotes and Citations:

- ¹ Grossbard, E. and Atkinson, D. (1985). *The Herbicide Glyphosate*. Butterworths, London; Harris, W.R., Sammons, R.D., Grabiak, R.C., Mehrsheikh, A., and Bleeke, M.S. Computer simulation of the interactions of glyphosate with metal ions in phloem. *Journal of Agricultural and Food Chemistry*. (2012). Vol 60, 6077-6087.
- ² Bohn and Fagan (2014). <http://www.sciencedirect.com/science/article/pii/S0308814613019201>; <https://www.independentsciencenews.org/news/how-extreme-levels-of-roundup-in-food-became-the-industry-norm/>; and <https://www.testbiotech.org/en/node/926>
- ³ USDA Economic Research Service, Adoption of Genetically Engineered Crops in the U.S. (2016). <https://www.ers.usda.gov/data-products/adoption-of-genetically-engineered-crops-in-the-us.aspx>
- ⁴ USDA Economic Research Service, Adoption of Genetically Engineered Crops in the U.S. <https://www.ers.usda.gov/data-products/adoption-of-genetically-engineered-crops-in-the-us.aspx>
- ⁵ Ridler, K. (2015). USDA to approve Simplot's genetically engineered potato that resists late blight. Associated Press. Sourced in US News and World Report, Aug 28, 2015. <https://www.usnews.com/news/business/articles/2015/08/28/usda-to-approve-simplots-genetically-engineered-potato>
- ⁶ Gurian-Sherman, D. (2003). Holes in the Biotech Safety Net: FDA Policy Does Not Assure the Safety of Genetically Engineered Foods, Center For Science in the Public Interest, January 1, 2003 https://cspinet.org/sites/default/files/attachment/fda_report_final.pdf
- ⁷ See for example, Alan M. Rulis Ph.D., Director, Office of Premarket Approval, Center for Food Safety and Applied Nutrition, U.S. Food and Drug Administration, to Dr. Kent Croon, Regulatory Affairs Manager, Monsanto Company, September 25, 1996. <http://www.fda.gov/Food/IngredientsPackagingLabeling/GEPlants/Submissions/ucm161107.htm>
- ⁸ "Statement of Policy: Foods Derived from New Plant Varieties," Federal Register 57, no. 104 (May 29, 1992): 22991
- ⁹ To view 24 memos by various FDA employees, visit <http://responsibletechnology.org/fraud/quotes-from-fda-scientists/>.
- ¹⁰ Memorandum from David Kessler, Commissioner of Food & Drugs. Subject: "FDA Proposed Statement of Policy Clarifying the Regulation of Food Derived from Genetically Modified Plants—DECISION." Dated March 20, 1992. (4 pages) <http://responsibletechnology.org/fraud/quotes-from-fda-scientists/>

- ¹¹ Robin, M. (2013). *The World According to Monsanto: Pollution, Corruption, and the Control of the World's Food Supply*. The New Press, Dec 13, 2013, Pp 234-235
- ¹² A full discussion of the limitations of industry safety studies is found in Jeffrey M. Smith, *Genetic Roulette: The Documented Health Risks Of Genetically Engineered Foods*, Yes! Books, Fairfield, IA, USA 2007, Part 3. Samples of other sources include: Ian F. Pryme and Rolf Lembcke, "In Vivo Studies on Possible Health Consequences of Genetically Modified Food and Feed—with Particular Regard to Ingredients Consisting of Genetically Modified Plant Materials," *Nutrition and Health* 17 (2003): 1–8; A. Pusztai and S. Bardocz, "GMO in animal nutrition: potential benefits and risks," Chapter 17, *Biology of Nutrition in Growing Animals*, R. Mosenthin, J. Zentek and T. Zebrowska (Eds.) (Elsevier, October 2005); Terje Traavik and Jack Heinemann, *Genetic Engineering and Omitted Health Research: Still No Answers to Ageing Questions*, TWN Biotechnology & Biosafety Series 7, 2007; William Freese, "Genetically Engineered Crop Health Impacts Evaluation: A Critique of U.S. Regulation of Genetically Engineered Crops and Corporate Testing Practices, with a Case Study of *Bt* Corn," Friends of the Earth U.S., <http://www.foe.org/camps/comm/safefood/gefood/index.html>; Marc Lappé and Britt Bailey, "American Soybean Association Response," Center for Toxics and Ethics, June 25, 1999, <http://www.cetos.org/articles/asaresponse.html>; Jeffrey M. Smith, "GM Peas," *Ecologist*, March 2006; and Jeffrey M. Smith, "Science or Nonsense? Jeffrey Smith responds to Hans Lombard," *Noseweekonline*, January 2006.; Jeffrey M. Smith, *Seeds of Deception* (Fairfield, IA: Yes! Books, 2003).; Doug Gurian-Sherman, "Holes in the Biotech Safety Net, FDA Policy Does Not Assure the Safety of Genetically Engineered Foods," Center for Science in the Public Interest, http://www.cspinet.org/new/pdf/fda_report_final.pdf; "Elements of Precaution: Recommendations for the Regulation of Food Biotechnology in Canada; An Expert Panel Report on the Future of Food Biotechnology prepared by The Royal Society of Canada at the request of Health Canada Canadian Food Inspection Agency and Environment Canada" The Royal Society of Canada, January 2001.; Food Legislation and Regulation Advisory Group (FLRAG) of the Public Health Association of Australia (PHAA) on behalf of the PHAA, "Comments to ANZFA about Applications A372, A375, A378 and A379," <http://www.iher.org.au/>; Mae-Wan Ho, "Exposed: More shoddy science in GM maize approval," *Science in Society* 22 (Summer 2004).; Judy Carman, "The Problem with the Safety of Round-Up Ready Soybeans," from *Diverse Women for Diversity*, 8th August 1999, <http://members.inet.net.au/~rabbit/noticsoy.htm>; William Freese, "A Critique of the EPA's Decision to Reregister *Bt* Crops and an Examination of the Potential Allergenicity of *Bt* Proteins," adapted from comments of Friends of the Earth to the EPA, Dec. 9, 2001, <http://www.foe.org/safefood/comments.pdf>; William Freese and David Schubert, "Safety Testing and Regulation of Genetically Engineered Foods," *Biotechnology and Genetic Engineering Reviews* 21 (November 2004); David Schubert, "A different perspective on GM food," *Nature Biotechnology* 20(2002): 969.; A. Pusztai, "Can science give us the tools for recognizing possible health risks of GM food?" *Nutr. Health* 16(2002): 73–84; Jeffrey M. Smith, "Genetically Modified Corn Study Reveals Health Damage and Cover-up," *Ecologist*, October 2005; Arpad Pusztai, "Why the Mon 863 Study Should Have Been Rejected," press backgrounder, June 22, 2005; Judy Carman, GE Foods and Human Health Safety Assessments, www.truefood.org.au/downloads/gefandhumanhealthsafetyassessments.pdf; Marc Lappé and Britt Bailey, "ASA Response," June 25, 1999, <http://cetos.org/articles/asaresponse.html>; Jack Heinemann et al., "Submission on Application A549 Food Derived from High Lysine Corn LY038: to permit the use in food of high lysine corn," Submitted to Food Standards Australia/New Zealand (FSANZ) by New Zealand Institute of Gene Ecology, January 22, 2005; William Freese, "The StarLink Affair, Submission by Friends of the Earth to the FIFRA Scientific Advisory Panel considering Assessment of Additional Scientific Information Concerning StarLink Corn," July 17–19, 2001; Masaharu Kawata, "Dr. M. Kawata: Questions CDC and EPA testing of Starlink and allergies," *Renu Namjoshi*, June 28, 2001; Barbara Keeler, "News Column," *Whole Life Times*, August 2000; Mae-Wan Ho, "Unstable Transgenic Lines Illegal," Institute for Science in Society, Press Release, March 12, 2003, <http://www.i-sis.org.uk/UTLI.php>;
- ¹³ "Genetically modified foods, who knows how safe they are?" CBC News and Current Affairs, September 25, 2006.
- ¹⁴ Friends of the Earth Press release, "Supermarket Loyalty Cards to Track GM Food Threat," January 25, 1999 https://www.foe.co.uk/resource/press_releases/0125card
- ¹⁵ American Academy of Environmental Medicine, (1999). Genetically Modified Foods, May 1999 <https://www.aaemonline.org/gmo.php>
- ¹⁶ "The American Academy Of Environmental Medicine Calls For Immediate Moratorium On Genetically Modified Foods," Press Advisory May 19, 2009 <http://aaemonline.org/aaemonline/oldsite/gmopressrelease.html>
- ¹⁷ Schubert D, (2002). Personal communication to H. Penfound, Greenpeace Canada, October 25, 2002
- ¹⁸ Fagan, J., Antoniou, M., & Robinson, C. (2014). GMO Myths and Truths. Earth Open Source. Pp 13-14
- ¹⁹ Bao, PH, Granata,, S., Castiglione, S., Wang, G., Giordani, C., Cuzzoni, E., (1996). Evidence for genomic changes in transgenic rice (*Oryza sativa* L.) recovered from protoplasts. *Transgenic Res* 1996 Mar;5(2):97-103; Labra, M., Savini, C., Bracale, M., Pelucchi, N., Colombo, L., Bardini, M., & Sala, F. (2001). "Genomic changes in transgenic rice (*Oryza sativa* L.) plants produced by infecting calli with *Agrobacterium tumefaciens*," *Plant Cell Rep* 20 (2001): 325-330.
- ²⁰ Srivastava, M., Eidelman, O., Pollard, HB. (1999). Pharmacogenomics of the cystic fibrosis transmembrane conductance regulator (CFTR) and the cystic fibrosis drug CPX using genome microarray analysis. *Mol Med*. 1999 Nov;5(11):753-67.
- ²¹ The original study: Stephen R. Padgett et al., "The Composition of Glyphosate-Tolerant Soybean Seeds Is Equivalent to That of Conventional Soybeans," *The Journal of Nutrition* 126, no. 4, (April 1996), left out the data from the cooked soybeans, which was recovered from the Journal by Barbara Keeler and later referenced in A. Pusztai and S. Bardocz, "GMO in animal nutrition: potential benefits and risks," Chapter 17, *Biology of Nutrition in Growing Animals*. Elsevier, 2005.
- ²² Zolla, L., Rinalducci, S., Antonioli, P., Righetti, PG (2008). Proteomics as a complementary tool for identifying unintended side effects occurring in transgenic maize seeds as a result of genetic modifications. *J Proteome Res*. 2008 May;7(5):1850-61. doi: 10.1021/pr0705082. Epub 2008 Apr 5.
- ²³ Betarbet, R. et al (2000). Chronic systemic pesticide exposure reproduces features of Parkinson's disease. *Nature Neurosci*. 3, 1301-1306 (2000); cited in cited in David Schubert, Regulatory regimes for transgenic crops. *Nature Biotechnology* 23, 785 - 787 (2005) doi:10.1038/nbt0705-785b.
- ²⁴ GMWatch, (2016). GM maize NK603 is not substantially equivalent to its non-GMO counterpart, <http://gmwatch.org/news/latest-news/17378-gmo-maize-nk603-is-not-substantially-equivalent-to-non-gmo-counterpart>
- ²⁵ citing Mesnage, R., Agapito-Tenfen, S., Vilperte, V., Renney, G., Ward, M., Séralini, GE, Nodari, N., Antoniou, MN. An integrated multi-omics analysis of the NK603 Roundup-tolerant GM maize reveals metabolism disturbances caused by the transformation process. *Scientific Reports*, 2016; 6:37855. <http://www.nature.com/articles/srep37855>
- ²⁶ Prescott, V. E. et al (2005). Transgenic Expression of Bean α -Amylase Inhibitor in Peas Results in Altered Structure and Immunogenicity, *J. Agric. Food Chem*. 2005, 53
- ²⁷ Zolla, L., Rinalducci, S., Antonioli, P., Righetti, PG (2008). Proteomics as a complementary tool for identifying unintended side effects occurring in transgenic maize seeds as a result of genetic modifications. *J Proteome Res*. 2008 May;7(5):1850-61. doi: 10.1021/pr0705082. Epub 2008 Apr 5.
- ²⁸ Heinemann, J., Zanon Agapito-Tenfen, S., & Carman, J. (2013). A comparative evaluation of the regulation of GM crops or products containing dsRNA and suggested improvements to risk assessments. *Environment International*, vol. 55, May 2013. Pp 43-55
- ²⁹ Netherwood et al., (2004). "Assessing the survival of transgenic plant DNA in the human gastrointestinal tract," *Nature Biotechnology* 22 (2004): 2.
- ³⁰ See for example, Jeffrey M. Smith, *Seeds of Deception*, (Iowa: Yes! Books, 2003) Chapter 1.
- ³¹ Ewen, Stanley W. B., and Pusztai, A. (1999). "Effect of diets containing genetically modified potatoes expressing *Galanthus nivalis* lectin on rat small intestine," *Lancet*, 1999 Oct 16; 354 (9187): 1353-4.
- ³² Arpad Pusztai, "Can science give us the tools for recognizing possible health risks of GM food," *Nutrition and Health*, 2002, Vol 16 Pp 73-84;

- Stanley W. B. Ewen and Arpad Pusztai, "Effect of diets containing genetically modified potatoes expressing *Galanthus nivalis* lectin on rat small intestine," *Lancet*, 1999 Oct 16; 354 (9187): 1353-4; A. Pusztai and S. Bardocz, "GMO in animal nutrition: potential benefits and risks," Chapter 17, *Biology of Nutrition in Growing Animals*, R. Mosenthin, J. Zentek and T. Zebrowska (Eds.) Elsevier, October 2005; and A. Pusztai, et al., "Genetically Modified Foods: Potential Human Health Effects," in: *Food Safety: Contaminants and Toxins* (ed. JPF D'Mello) (Wallingford Oxon, UK: CAB International), 347–372.
- ³² See for example, Silvia Caccia, (2016). Midgut microbiota and host immunocompetence underlie *Bacillus thuringiensis* killing mechanism, Proceedings of the National Academy of Sciences, August 23, 2016 vol. 113 no. 34, <http://www.pnas.org/content/113/34/9486.full>
- ³³ Fagan, J., Antoniou, M., & Robinson, C. (2014). GMO Myths and Truths. Earth Open Source. Pp 172-173
- ³⁴ Vazquez, A. et al. (2000). "Intragastric and intraperitoneal administration of Cry1Ac protoxin from *Bacillus thuringiensis* induces systemic and mucosal antibody responses in mice," 1897–1912; Vazquez, A. et al., "Characterization of the mucosal and systemic immune response induced by Cry1Ac protein from *Bacillus thuringiensis* HD 73 in mice," *Brazilian Journal of Medical and Biological Research* 33 (2000): 147–155; Vazquez, A. et al., "Bacillus thuringiensis Cry1Ac protoxin is a potent systemic and mucosal adjuvant," *Scandinavian Journal of Immunology* 49 (1999): 578–584; Vazquez, A. et al., "Cry1Ac protoxin from *Bacillus thuringiensis* ssp. *kurstaki* HD73 binds to surface proteins in the mouse small intestine," 54–58.
- ³⁵ Finamore, A., Roselli, M., Britti, S., Monastra, G., Ambra, R., Turrini, A., Mengheri, E (2008). Intestinal and Peripheral Immune Response to MON810 Maize Ingestion in Weaning and Old Mice. *J Agric Food Chem*. 2008; 56 (23): 11533–11539.
- ³⁶ de Vendômois, JS, Roullier, F., Cellier, D., Séralini, GE (2009). A Comparison of the Effects of Three GM Corn Varieties on Mammalian Health. *Int J Biol Sci*. 2009; 5:706-726. Available from <http://www.biolsci.org/v05p0706.htm>; and John M. Burns, "13-Week Dietary Subchronic Comparison Study with MON 863 Corn in Rats Preceded by a 1-Week Baseline Food Consumption Determination with PMI Certified Rodent Diet #5002," December 17, 2002. See also Stéphane Foucart, "Controversy Surrounds a GMO," *Le Monde*, 14 December 2004.
- ³⁷ Mezzomo, et al., Hematotoxicity of *Bacillus thuringiensis* as Spore-crystal Strains Cry1Aa, Cry1Ab, Cry1Ac or Cry2Aa in Swiss Albino Mice, *Journal of Hematology & Thromboembolic Diseases*, 2013, 1:1 <http://gmoevidence.com/dr-mezzomo-bt-toxins-toxic-to-blood-of-mice/>
- ³⁸ Mesnage R, Clair E, Gress S, Then C, Székács A, Séralini GE (2013). Cytotoxicity on human cells of Cry1Ab and Cry1Ac Bt insecticidal toxins alone or with a glyphosate-based herbicide. *J App. Toxicol*. 2013 Jul;33(7):695-9. doi: 10.1002/jat.2712. Epub 2012 Feb 15.
- ³⁹ Washington State Department of Health (1993). "Report of health surveillance activities: Asian gypsy moth control program," (Olympia, WA: Washington State Dept. of Health, 1993); Green, M. "Public health implications of the microbial pesticide *Bacillus thuringiensis*: An epidemiological study, Oregon, 1985-86," *Amer. J. Public Health* 80, no. 7(1990): 848–852; M.A. Noble, P.D. Riben, and G. J. Cook, "Microbiological and epidemiological surveillance program to monitor the health effects of Foray 48B BTK spray" (Vancouver, B.C.: Ministry of Forests, Province of British Columbia, Sep. 30, 1992).
- ⁴⁰ Gupta, A. et. al. (2005). "Impact of Bt Cotton on Farmers' Health (in Barwani and Dhar District of Madhya Pradesh)," *Investigation Report*, Oct–Dec 2005.
- ⁴¹ *Sunday Indian*, according to hospital records: "Victims of itching have increased massively this year . . . related to BT cotton farming." October 26, 2008
- ⁴² Aris, A., Leblanc, S. (2011). Maternal and fetal exposure to pesticides associated to genetically modified foods in Eastern Townships of Quebec, Canada. *Reprod Toxicol*. 2011 May;31(4):528-33. doi: 10.1016/j.reprotox.2011.02.004. Epub 2011 Feb 18..
- ⁴³ Mezzomo, et al., Hematotoxicity of *Bacillus thuringiensis* as Spore-crystal Strains Cry1Aa, Cry1Ab, Cry1Ac or Cry2Aa in Swiss Albino Mice, *Journal of Hematology & Thromboembolic Diseases*, 2013, 1:1 <http://gmoevidence.com/dr-mezzomo-bt-toxins-toxic-to-blood-of-mice/>
- ⁴⁴ *Sets of Scientific Issues Being Considered by the Environmental Protection Agency Regarding: Bt Plant-Pesticides Risk and Benefit Assessments*, SAP Report No. 2000-07 March 12, 2001, FIFRA Scientific Advisory Panel Meeting, October 18-20, 2000
- ⁴⁵ Ho MW, Unstable Transgenic Lines Illegal, I-SIS report, 23 October 2003. <http://www.i-sis.org.uk/CAGMMAD.php>
- ⁴⁶ FAO-WHO, "Evaluation of Allergenicity of Genetically Modified Foods. Report of a Joint FAO/WHO Expert Consultation on Allergenicity of Foods Derived from Biotechnology," Jan. 22–25, 2001; <http://www.fao.org/es/ESN/food/pdf/allergvym.pdf>; D.D.Metcalf et al., "Assessment of the Allergenic Potential of Foods Derived from Genetically Engineered Crop Plants," *Critical Reviews in Food Science and Nutrition* 36(S) (1996): S165–186; and Gendel, "The use of amino acid sequence alignments to assess potential allergenicity of proteins used in genetically modified foods," *Advances in Food and Nutrition Research* 42 (1998), 45–62; H. P. J. M. Noteborn, "Assessment of the Stability to Digestion and Bioavailability of the LYS Mutant Cry9C Protein from *Bacillus thuringiensis* serovar *tolworthi*," Unpublished study submitted to the EPA by AgrEvo, EPA MRID No. 447343-05 (1998); and H. P. J. M. Noteborn et al., "Safety Assessment of the *Bacillus thuringiensis* Insecticidal Crystal Protein CRYIA(b) Expressed in Transgenic Tomatoes," in *Genetically modified foods: safety issues*, American Chemical Society Symposium Series 605, eds. K.H. Engel. (Washington, DC, 1995): 134–47.
- ⁴⁷ Benbrook, C. (2012). Impacts of genetically engineered crops on pesticide use in the U.S.—the first sixteen years, *Environmental Sciences Europe* 2012, 24:24 <http://www.enveurope.com/content/24/1/24>
- ⁴⁸ Peterson Myers, J. (2016). Concerns over use of glyphosate-based herbicides and risks associated with exposures: a consensus statement, *Environmental Health* 2016 15:19 DOI: 10.1186/s12940-016-0117-0, <http://ehjournal.biomedcentral.com/articles/10.1186/s12940-016-0117-0>
- ⁴⁹ Benbrook, C. (2016). Trends in glyphosate herbicide use in the United States and globally, *Environmental Sciences Europe Bridging Science and Regulation at the Regional and European Level* 2016 28:3, 2 February 2016
- ⁵⁰ See for example: <https://www.regulations.gov/document?D=EPA-HQ-OPP-2012-0132-0009>; and Mary Ellen Kustin, americans At Greater Risk Of Glyphosate Exposure than Europeans, *Ag Mag*, EWG, FEB 3, 2016
- ⁵¹ Samsel, A. & Seneff, S. (2013). Glyphosate's Suppression of Cytochrome P450 Enzymes and Amino Acid Biosynthesis by the Gut Microbiome: Pathways to Modern Diseases, *Entropy*, 18 April 2013, <http://www.mdpi.com/1099-4300/15/4/1416>
- ⁵² See list at <http://responsibletechnology.org/list-of-references-on-the-health-effects-of-glyphosate/>; Antoniou, M. et al., GM SOY, Sustainable? Responsible? Earth Open Source, September 2010 <http://earthopensource.org/earth-open-source-reports/gm-soy-sustainable-responsible/>; and Antoniou, M. et al., Roundup and birth defects: Is the public being kept in the dark? Earth Open Source, 2011 <http://www.scribd.com/doc/57277946/RoundupandBirthDefectsv5>; and Watts, M., Glyphosate, Monograph prepared for Pesticide Action Network, International, October 2016
- ⁵³ International Agency for Research on Cancer, World Health Organization (2015). *IARC Monographs Volume 112: evaluation of five organophosphate insecticides and herbicides*, March 20, 2015 <http://www.iarc.fr/en/media-centre/iarcnews/pdf/MonographVolume112.pdf>; and WHO IARC Q & A on Glyphosate, 1 March 1, 2016 https://www.iarc.fr/en/media-centre/iarcnews/pdf/Q&A_Glyphosate.pdf
- ⁵⁴ See for example, Ganson, R.J. and Jensen, R.A. (1988). The essential role of cobalt in the inhibition of the cytosolic isozyme of 3-deoxy-D-arabino-heptulosonate-7-phosphate synthase from *Nicotiana glauca* by glyphosate. *Arch Biochem. Biophys.* 260(1):85-93; Lee, T.T. 1982. Mode of action of glyphosate in relation to metabolism of indole-3-acetic acid. *Physiol. Plantar.* 54:289; and Liu, L., Punja, Z.K. and Rahe, J.E. 1997. Altered root exudation and suppression of induced lignification as mechanisms of predisposition by glyphosate of bean roots (*Phaseolus vulgaris* L.) to colonization by *Pythium* spp. *Physiol. Mol. Plant Pathol.* 51:111-127.
- ⁵⁵ See for example: Ganson, R.J. and Jensen, R.A. (1988). The essential role of cobalt in the inhibition of the cytosolic isozyme of 3-deoxy-D-arabino-heptulosonate-7-phosphate synthase from *Nicotiana glauca* by glyphosate. *Arch Biochem. Biophys.* 260(1):85-93. Johal, G.R. and Rahe, J.E. 1984. Effect of soilborne plant-pathogenic fungi on the herbicidal action of glyphosate on bean seedlings. *Phytopathology* 74:950-955. Johal, G.R. and Huber, D.M. 2009. Glyphosate effects on diseases of plants. *Eur. J. Agron.* 31:144-152. Zobiolo, L.H.S., Oliveira Jr., R.S.,

- Constantin, J., Kremer, R.J., Biffe, D.F. 2010. Amino acid application can be an alternative to prevent glyphosate injury. *J. Plant Nutr.* Zobiole, L.H.S.; Oliveira Jr., R.S.; Constantin, J. and Biffe, D.F. 2011. Prevention of RR soybean injuries caused by exogenous supply of amino acids. *Planta daninha* [online]. 2011, vol.29, n.1, pp. 195-205. ISSN 0100-8358. Zobiole, L.H.S. 2010. Use of exogenous amino acid to prevent glyphosate injury in glyphosate-resistant soybean. *Planta daninha* [online]. 2010, vol.28, n.3, pp. 643-653. ISSN 0100-8358. doi: 10.1590/S0100-83582010000300022.
- ⁵⁶ Yano JM et al. (2015) Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. *Cell*, 161: 264-276.
- ⁵⁷ See for example, Katholm, C.L. (2016). Effects of Roundup (glyphosate) on gut microorganisms of farm animals. http://library.au.dk/fileadmin/www.bibliotek.au.dk/faqsider/jordbrug/Specialer/Master_Thesis_final.pdf; Krüeger, M., Shehata, A., Neuhaus, J., Muller, T., Kotsch, M., and Schrod, W. 2011. Chronischer Botulismus des Rindes eine Faktorenkrankheit Welche Rolle spielt das Totalherbizid Glyphosat? Institut für Bakteriologie und Mykologie, Universität Leipzig.
- Krüeger, M., Shehata, A., Neuhaus, J., Muller, T., Kotsch, M., and Schrod, W. 2012. Chronic botulism of cattle, a multifactorial disease. What is the role of the herbicide glyphosate? *Inst. Fur Bakteriol. Und Mykol., Univ. Leipzig, Germany*; Krüeger, M., et al., 2013. Glyphosate suppresses the antagonistic effect of *Enterococcus* spp. on *Clostridium botulinum*. *Anaerobe* 20:74-78; Lu, W., Li, L., Cheng, M., Zhou, Z., Zhang, W., Ping, S., Yan, Y., Wang, J. and Lin, M. 2013. Genome-wide transcriptional responses of *Escherichia coli* to glyphosate, a potent inhibitor of the shikimate pathway enzyme 5-enolpyruvylshikimate-3-phosphate synthase. *Mol. Biosyst.* 9:522-530. Doi: 10.1039/C2MB25374G; Shehata, A., Schrod, W., Aldin, A.A., Hafez, H.M., Krueger, M. 2012. The effect of glyphosate on potential pathogens and beneficial members of poultry microbiota in vitro. *Current Microbiology* DOI 10.1007/s00284-012-0277-2; Shehata, A., Schrod, W., Neuhaus, J., and Krueger, M. 2012. Antagonistic effect of different bacteria on *Clostridium botulinum* types A, B, C, D and E in vitro. *Veterinary Rec.* 1-2. Doi: 10.1136/vr.101184; Shehata, A.A., Schrödl, W., Aldin, A.A., Hafez, H.M., Krüger, M. 2012. The Effect of Glyphosate on Potential Pathogens and Beneficial Members of Poultry Microbiota in Vitro. *Curr Microbiol.* Doi: 10.1007/s00284-012-0277-2; Shehata, A.A., Kuhnert, M., Haufe, S., and Kruger, M. Neutralization of the antimicrobial effect of glyphosate by humic acid in vitro. *Chemosphere* (2013). <http://dx.doi.org/10.1016/j.chemosphere.2013.10.064>.
- ⁵⁸ Kim, YH (2013). Mixtures of glyphosate and surfactant TN20 accelerate cell death via mitochondrial damage-induced apoptosis and necrosis. *Toxicol In Vitro.* 2013 Feb; 27(1):191-7. doi: 10.1016/j.tiv.2012.09.021. Epub 2012 Oct 23; and Heu, , A step further toward glyphosate-induced epidermal cell death: involvement of mitochondrial and oxidative mechanisms. *Environ Toxicol Pharmacol.* 2012 Sep;34(2):144-53. doi: 10.1016/j.etap.2012.02.010. Epub 2012 Mar 15; and Peixoto F., Comparative effects of the Roundup and glyphosate on mitochondrial oxidative phosphorylation. *Chemosphere.* 2005 Dec;61(8):1115-22. Epub 2005 Apr 26.
- ⁵⁹ Paganelli, A. et al. (2010). Glyphosate-Based Herbicides Produce Teratogenic Effects on Vertebrates by Impairing Retinoic Acid Signaling. *Chem. Res. Toxicol.*, 2010, 23 (10), pp 1586–1595; Dallegrave, E., Mantese, F. D. 2007. Pre- and postnatal toxicity of the commercial glyphosate formulation in Wistar rats. *Arch Toxicol* 81: 665–673; Dallegrave, E., Mantese, F. D. 2003. e teratogenic potential of the herbicide glyphosate-Roundup in Wistar rats. *Toxicol Lett* 142(1–2): 45–52; and Lajmanovich, R. C., Sandoval, M. T., Peltzer, P. M. 2003. Induction of mortality and malformation in *Scinax nasicus* tadpoles exposed to glyphosate formulations. *Bull. Environ. Contam. Toxicol.* 70, 612–618.
- ⁶⁰ See for example, Cassault-Meyer E, et al., [An acute exposure to glyphosate-based herbicide alters aromatase levels in testis and sperm nuclear quality](#), *Environ Toxicol Pharmacol.* 2014 Jul;38(1):131-40; and Kwiatkowska, et al., [Glyphosate and its formulations--toxicity, occupational and environmental exposure](#), *Med Pr.* 2013;64(5):717-29.
- ⁶¹ Séralini, GE (2014). Republished study: long-term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize, *Environmental Sciences Europe*, 2014, 26:14
- ⁶² Varayoud, J. (2016). Effects of a glyphosate-based herbicide on the uterus of adult ovariectomized rats, *Environmental Toxicology*, 27 July 2016
- ⁶³ Thongprakaisang, S. et al. (2013). Glyphosate induces human breast cancer cells growth via estrogen receptors. *Food Chem Toxicol.* 2013 Sep;59:129-36.
- ⁶⁴ Mesnage R, Phedonos A, Biserni M, Arno M, Balu S, Corton JC, Ugarte R, Antoniou MN, Evaluation of estrogen receptor alpha activation by glyphosate-based herbicide constituents, *Food Chem Toxicol.* 2017 Jul 12;108(Pt A):30-42. doi: 10.1016/j.fct.2017.07.025. [Epub ahead of print]
- ⁶⁵ Séralini, GE, Clair, E., Mesnage, R., Gress, S., Defarge, N., Malatesta, M. (2012). Long term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize. *Food Chem Toxicol.*, 2012; 50(11):4221-4231; NS Gilles-Eric Séralini, Republished study: long-term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize, *Environmental Sciences Europe*, 2014, 26:14
- ⁶⁶ Mesnage, R., Arno, M., Costanzo, M., Malatesta, M., Séralini, GE, & Antoniou, M. (2015). Transcriptome profile analysis reflects rat liver and kidney damage following chronic ultra-low dose Roundup exposure, *Environmental Health*, 2015; 14:70
- ⁶⁷ Mesnage, R., Renney, G., Séralini, GE, Ward, M. & Antoniou, M. (2017). Multiomics reveal non-alcoholic fatty liver disease in rats following chronic exposure to an ultra-low dose of Roundup herbicide. *Scientific Reports*, 2017; 7:39328. <http://www.nature.com/srep/2017/170104/srep39328/full/srep39328.html>
- ⁶⁸ Fagan, J., Antoniou, M., & Robinson, C. (2014). GMO Myths and Truths. Earth Open Source. Chapter 3, <http://earthopenresource.org/gmomysandtruths/sample-page/3-health-hazards-gm-foods/3-1-myth-gm-foods-safe-eat/>
- ⁶⁹ Séralini GE, Clair E, Mesnage R, et al., Long term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize, *Environmental Sciences Europe*, 26:14 June 24, 2014 <https://doi.org/10.1186/s12302-014-0014-5>
- ⁷⁰ Gab-Alla AA, El-Shamei ZS, Shatta AA, Moussa EA, Rayan AM. Morphological and biochemical changes in male rats fed on genetically modified corn (Ajeeb YG). *J Am Sci.* 2012;8(9):1117–1123; and El-Shamei ZS, Gab-Alla AA, Shatta AA, Moussa EA, Rayan AM. Histopathological changes in some organs of male rats fed on genetically modified corn (Ajeeb YG). *J Am Sci.* 2012;8(10):684–696.
- ⁷¹ Hines FA. Memorandum to Linda Kahl on the Flavr Savr tomato (Pathology Review PR–152; FDA Number FMF–000526): Pathology Branch's evaluation of rats with stomach lesions from three four-week oral (gavage) toxicity studies (IRDC Study Nos. 677–002, 677–004, and 677–005) and an Expert Panel's report. US Department of Health & Human Services; 1993. Available at: <http://www.biointegrity.org/FDAdocs/17/view1.html>; Pusztai A. Witness Brief – Flavr Savr tomato study in Final Report (IIT Research Institute, Chicago, IL 60616 USA) cited by Dr Arpad Pusztai before the New Zealand Royal Commission on Genetic Modification. 2000. Available at: <http://www.gmcommission.govt.nz/>; Pusztai A. Can science give us the tools for recognizing possible health risks of GM food? *Nutr Health.* 2002;16:73-84; Scheuplein RJ. Memorandum: Response to Calgene amended petition. 1993. Available at: <http://www.biointegrity.org/FDAdocs/11/view1.html>; Pusztai A, Bardocz S, Ewen SWB. Genetically modified foods: Potential human health effects. In: D'Mello JPF, ed. *Food Safety: Contaminants and Toxins*. Wallingford, Oxon: CABI Publishing; 2003:347–372. Available at: <http://www.leopold.iastate.edu/sites/default/files/events/Chapter16.pdf>.
- ⁷² Prescott VE, Campbell PM, Moore A, et al. Transgenic expression of bean alpha-amylase inhibitor in peas results in altered structure and immunogenicity. *J Agric Food Chem.* 2005;53:9023–30. doi:10.1021/jf050594v.
- ⁷³ Finamore A, Roselli M, Britti S, et al. Intestinal and peripheral immune response to MON810 maize ingestion in weaning and old mice. *J Agric Food Chem.* 2008;56:11533–39. doi:10.1021/jf802059w; and Kroghsbo S, Madsen C, Poulsen M, et al. Immunotoxicological studies of genetically modified rice expressing PHA-E lectin or Bt toxin in Wistar rats. *Toxicology.* 2008;245:24-34. doi:10.1016/j.tox.2007.12.005.
- ⁷⁴ Krzyzowska M, Wincenciak M, Winnicka A, et al. The effect of multigenerational diet containing genetically modified triticale on immune system in mice. *Pol J Vet Sci.* 2010;13:423-30.
- ⁷⁵ Malatesta M, Caporaloni C, Gavaudan S, et al. Ultrastructural morphometrical and immunocytochemical analyses of hepatocyte nuclei from mice fed on genetically modified soybean. *Cell Struct Funct.* 2002;27:173–80; Vecchio L, Cisterna B, Malatesta M, Martin TE, Biggiogera M. Ultrastructural analysis of testes from mice fed on genetically modified soybean. *Eur J Histochem.* 2004;48:448-54; and Malatesta M, Boraldi F,

- Annovi G, et al. A long-term study on female mice fed on a genetically modified soybean: effects on liver ageing. *Histochem Cell Biol.* 2008;130:967–977.
- ⁷⁶ Malatesta M, Boraldi F, Annovi G, et al. A long-term study on female mice fed on a genetically modified soybean: effects on liver ageing. *Histochem Cell Biol.* 2008;130:967–977.
- ⁷⁷ Tudisco R, Lombardi P, Bovera F, et al. Genetically modified soya bean in rabbit feeding: Detection of DNA fragments and evaluation of metabolic effects by enzymatic analysis. *Anim Sci.* 2006;82:193–199. doi:10.1079/ASC200530.
- ⁷⁸ Brasil FB, Soares LL, Faria TS, Boaventura GT, Sampaio FJ, Ramos CF. The impact of dietary organic and transgenic soy on the reproductive system of female adult rat. *Anat Rec Hoboken.* 2009;292:587–94. doi:10.1002/ar.20878.
- ⁷⁹ Carman JA, Vlieger HR, Ver Steeg LJ, et al. A long-term toxicology study on pigs fed a combined genetically modified (GM) soy and GM maize diet. *J Org Syst.* 2013;8:38–54.
- ⁸⁰ Séralini GE, Mesnage R, Clair E, Gress S, de Vendômois JS, Cellier D. Genetically modified crops safety assessments: Present limits and possible improvements. *Environ Sci Eur.* 2011;23. doi:10.1186/2190-4715-23-10; De Vendômois JS, Roullier F, Cellier D, Séralini GE. A comparison of the effects of three GM corn varieties on mammalian health. *Int J Biol Sci.* 2009;5:706–26; and Séralini GE, Clair E, Mesnage R, et al., Long term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize, *Environmental Sciences Europe*, 26:14 June 24, 2014 <https://doi.org/10.1186/s12302-014-0014-5>
- ⁸¹ Séralini GE, Cellier D, Spiroux de Vendômois J. New analysis of a rat feeding study with a genetically modified maize reveals signs of hepatorenal toxicity. *Arch Environ Contam Toxicol.* 2007;52:596–602.
- ⁸² Kilic A, Akay MT. A three generation study with genetically modified Bt corn in rats: Biochemical and histopathological investigation. *Food Chem Toxicol.* 2008;46:1164–70. doi:10.1016/j.fct.2007.11.016.
- ⁸³ US Food and Drug Administration (FDA). Biotechnology consultation note to the file BNF No 00077. Office of Food Additive Safety, Center for Food Safety and Applied Nutrition; 2002. Available at: <http://bit.ly/ZUMiAF>.
- ⁸⁴ Trabalza-Marinucci M, Brandi G, Rondini C, et al. A three-year longitudinal study on the effects of a diet containing genetically modified Bt176 maize on the health status and performance of sheep. *Livest Sci.* 2008;113:178–190. doi:10.1016/j.livsci.2007.03.009.
- ⁸⁵ Ewen SW, Pusztai A. Effect of diets containing genetically modified potatoes expressing Galanthus nivalis lectin on rat small intestine. *Lancet.* 1999;354:1353–4. doi:10.1016/S0140-6736(98)05860-7; and Pusztai A, Bardocz S. GMO in animal nutrition: Potential benefits and risks. In: Mosenthin R, Zentek J, Zebrowska T, eds. *Biology of Nutrition in Growing Animals*. Vol 4. Elsevier Limited; 2006:513–540. Available at: <http://www.sciencedirect.com/science/article/pii/S1877182309701043>.
- ⁸⁶ Fares NH, El-Sayed AK. Fine structural changes in the ileum of mice fed on delta-endotoxin-treated potatoes and transgenic potatoes. *Nat Toxins.* 1998;6(6):219–33.
- ⁸⁷ Poulsen M, Kroghsbo S, Schroder M, et al. A 90-day safety study in Wistar rats fed genetically modified rice expressing snowdrop lectin Galanthus nivalis (GNA). *Food Chem Toxicol.* 2007;45:350–63. doi:10.1016/j.fct.2006.09.002.
- ⁸⁸ Schrøder M, Poulsen M, Wilcks A, et al. A 90-day safety study of genetically modified rice expressing Cry1Ab protein (Bacillus thuringiensis toxin) in Wistar rats. *Food Chem Toxicol.* 2007;45:339–49. doi:10.1016/j.fct.2006.09.001.
- ⁸⁹ Gu J, Krogdahl A, Sissener NH, et al. Effects of oral Bt-maize (MON810) exposure on growth and health parameters in normal and sensitised Atlantic salmon, *Salmo salar* L. *Br J Nutr.* 2013;109:1408–23. doi:10.1017/S000711451200325X.
- ⁹⁰ Agodi A, Barchitta M, Grillo A, Sciacca S. Detection of genetically modified DNA sequences in milk from the Italian market. *Int J Hyg Env Health.* 2006;209:81–8. doi:10.1016/j.ijheh.2005.08.005.
- ⁹¹ Aris, A., Leblanc, S. (2011). Maternal and fetal exposure to pesticides associated to genetically modified foods in Eastern Townships of Quebec, Canada. *Reprod Toxicol.* 2011;31. <http://www.ncbi.nlm.nih.gov/pubmed/21338670>
- ⁹² Chowdhury, EH, Shimada, N., Murata, H., Mikami, O., Sultana, P., Miyazaki, S. (2003). Detection of Cry1Ab protein in gastrointestinal contents but not visceral organs of genetically modified Bt11-fed calves. *Vet Hum Toxicol* 2003;45:72–5. [39]; Chowdhury, EH, Kuribara, H., Hino, A., Sultana, P., Mikami, O., Shimada, N (2003). Detection of corn intrinsic and recombinant DNA fragments and Cry1Ab protein in the gastrointestinal contents of pigs fed genetically modified corn Bt11. *J Anim Sci* 2003;81:2546–51. [40] Lutz, B., Wiedemann, S., Einspanier, R., Mayer, J., Albrecht, C (2005). Degradation of Cry1Ab protein from genetically modified maize in the bovine gastrointestinal tract. *J Agric Food Chem* 2005;53:1453–6
- ⁹³ Monsanto Company. Residues in Roundup Ready soya lower than conventional soy. http://www.monsanto.co.uk/news/99/june99/220699_residue.html . 1999.
- ⁹⁴ Bøhn, T., Cuhra, M., Traavik, T., Sanden, M., Fagan, J. and Primicerio, R. (2014). Compositional differences in soybeans on the market: Glyphosate accumulates in Roundup Ready GM soybeans. – *Food Chemistry* 153: 207–215. See also, Thomas Bøhn and Marek Cuhra, How “Extreme Levels” of Roundup in Food Became the Industry Norm, *Independent Science News*, March 24, 2014 <https://www.independentsciencenews.org/news/how-extreme-levels-of-roundup-in-food-became-the-industry-norm/>
- ⁹⁵ Chinese Report Shows High Levels of Glyphosate in Imported GM Soybeans, Posted by Sustainable Pulse, May 19, 2014, <http://sustainablepulse.com/2014/05/19/chinese-report-shows-high-levels-glyphosate-imported-gm-soybeans/#>
- ⁹⁶ Toxicological Summary for: Aminomethylphosphonic acid, Minnesota Department of Health, February 2017 CAS: 1066-51-9
- ⁹⁷ Bøhn, T., Cuhra, M., Traavik, T., Sanden, M., Fagan, J. and Primicerio, R. (2014). Compositional differences in soybeans on the market: Glyphosate accumulates in Roundup Ready GM soybeans. – *Food Chemistry* 153: 207–215.
- ⁹⁸ Samsel, A. & Seneff, S. Glyphosate, pathway to modern diseases IV: cancer and related pathologies. *J. Biol. Phys. Chem.* 15 (2015) 121–159.
- ⁹⁹ Gillam, C. (2015). Fears over Roundup herbicide residues prompt private testing, *Reuters*, April 10, 2015; and Zen Honeycutt, More Roundup Found in USA Mom's Breast Milk, August 04, 2015
- ¹⁰⁰ McGuire, et al., Glyphosate and aminomethylphosphonic acid are not detectable in human milk, *Am J Clin Nutr.* 2016 May;103(5):1285–90. doi: 10.3945/ajcn.115.126854. Epub 2016 Mar 30; and Steinbor, et al., Determination of Glyphosate Levels in Breast Milk Samples from Germany by LC-MS/MS and GC-MS/MS, *J Agric Food Chem.* 2016 Feb 17;64(6):1414–21. doi: 10.1021/acs.jafc.5b05852. Epub 2016 Feb 8.
- ¹⁰¹ Heinemann, J. (2009). Report on animals exposed to GM ingredients in animal feed. Prepared for the Commerce Commission of New Zealand, 24 July 2009
- ¹⁰² Glyphosate; Pesticide Tolerances, A Rule by the Environmental Protection Agency May 1, 2013, <https://www.federalregister.gov/documents/2013/05/01/2013-10316/glyphosate-pesticide-tolerances>
- ¹⁰³ Glyphosate: Unsafe on Any Plate, Food Testing Results and Scientific Reasons for Concern, Report by Food Democracy now and The Detox Project, November 2016 <http://bit.ly/glyphosateFood>
- ¹⁰⁴ Robinson, C. (2017). Roundup causes non-alcoholic fatty liver disease at very low doses. *GMWatch*. Referencing Mesnage R, Renney G, Séralini GE, Ward M, Antoniou MN. Multiomics reveal non-alcoholic fatty liver disease in rats following chronic exposure to an ultra-low dose of Roundup herbicide. *Scientific Reports*, 2016; 6:39328. <http://www.nature.com/articles/srep39328>
- ¹⁰⁵ Seidler, R. (2017). The time for glyphosate-based herbicides is over. *GMWatch*. 03 February 3, 2017 <http://gmwatch.org/news/latest-news/17456-the-time-for-glyphosate-based-herbicides-is-over>
- ¹⁰⁶ Battaglin, W.A. (2014). Glyphosate and Its Degradation Product AMPA Occur Frequently and Widely in U.S. Soils, Surface Water, Groundwater, and Precipitation. *Journal of the American Water Resources Association*, April 1, 2014, Vol 50, Issue 2, Pp 275–290

- ¹⁰⁷ Genetically Modified Organisms and the deterioration of health in the United States N.L. Swanson, 4/24/2013; <https://people.csail.mit.edu/seneff/glyphosate/NancySwanson.pdf>. This document was first published as a series of articles on Seattle examiner.com
- ¹⁰⁸ Genetically Modified Organisms and the deterioration of health in the United States N.L. Swanson, 4/24/2013; <https://people.csail.mit.edu/seneff/glyphosate/NancySwanson.pdf>. This document was first published as a series of articles on Seattle examiner.com
- ¹⁰⁹ Swanson, N. (2014). Genetically engineered crops, glyphosate and the deterioration of health in the United States of America, *Journal of Organic Systems*, 9(2), 2014
- ¹¹⁰ Swanson, N. (2014). Genetically engineered crops, glyphosate and the deterioration of health in the United States of America, *Journal of Organic Systems*, 9(2), 2014
- ¹¹¹ Pusztai, A. (2002). "Can science give us the tools for recognizing possible health risks of GM food," *Nutrition and Health*, 2002, Vol 16 Pp 73-84; and Stanley W. B. Ewen and Arpad Pusztai, "Effect of diets containing genetically modified potatoes expressing *Galanthus nivalis* lectin on rat small intestine," *Lancet*, 1999 Oct 16; 354 (9187): 1353-4.
- ¹¹² Pusztai, A. (2005). "Facts Behind the GM Pea Controversy: Epigenetics, Transgenic Plants & Risk Assessment," Proceedings of the Conference, December 1st 2005 (Frankfurt/Main, Germany: Literaturhaus, 2005). <http://www.oeko.de/oekodoc/277/2006-002-en.pdf>
- ¹¹³ Pusztai, A. (2001). "Genetically Modified Foods: Are They a Risk to Human/Animal Health?" *Action Bioscience*. June 2001
- ¹¹⁴ Carman, J. (2013). A long-term toxicology study on pigs fed a combined genetically modified (GM) soy and GM maize diet, *Journal of Organic Systems*, Volume 8 Number 1, June 2013
- ¹¹⁵ Stephen R. Padgett et al, Table 2 in "e Composition of Glyphosate-Tolerant Soybean Seeds Is Equivalent to that of Conventional Soybeans," *e Journal of Nutrition* 126, no. 4 (April 1996).
- ¹¹⁶ Pusztai, A. & Bardocz, S. (2005). "GMO in animal nutrition: potential benefits and risks," Chapter 17, *Biology of Nutrition in Growing Animals*, R. Mosenthin, J. Zentek and T. Zebrowska (Eds.) (Elsevier, October 2005)
- ¹¹⁷ Pusztai, A. & Bardocz, S. (2005). "GMO in animal nutrition: potential benefits and risks," Chapter 17, *Biology of Nutrition in Growing Animals*, R. Mosenthin, J. Zentek and T. Zebrowska (Eds.) (Elsevier, October 2005)
- ¹¹⁸ Malatesta, et al, "Ultrastructural Analysis of Pancreatic Acinar Cells from Mice Fed on Genetically Modified Soybean, *Journal of Anatomy*, 2002; Nov: 201(5) 409-415
- ¹¹⁹ Mesnage, R. (2012). Cytotoxicity on human cells of Cry1Ab and Cry1Ac Bt insecticidal toxins alone or with a glyphosate-based herbicide. *J Appl Toxicol*. 2013 Jul;33(7):695-9. doi: 10.1002/jat.2712. Epub 2012 Feb 15.
- ¹²⁰ Traavik, T. (2004). Conference presentation to delegates of the Cartagena Protocol for Biosafety, Kuala Lumpur, sponsored by the Third World Network, February 22, 2004
- ¹²¹ See for example, Barocelli E and Ballabeni V, Histamine in the control of gastric acid secretion: a topic review; *Pharmacol Res*. 2003 Apr;47(4):299-304. <https://www.ncbi.nlm.nih.gov/pubmed/12644386>
- ¹²² Fares, N., El-Sayed, A. (1998). "Fine Structural Changes in the Ileum of Mice Fed on Endotoxin Treated Potatoes and Transgenic Potatoes," *Natural Toxins* 6, no. 6 (1998): 219-233.
- ¹²³ Senapati, T., Mukerjee, AK, & Ghosh, AR (2009). Observations on the effect of glyphosate based herbicide on ultra structure (SEM) and enzymatic activity in different regions of alimentary canal and gill of *Channa punctatus* (Bloch). *Journal of Crop and Weed* 5(1): 236-245, 2009.
- ¹²⁴ Senapati, T., Mukerjee, AK, Ghosh, AR (2009). Observations on the effect of glyphosate based herbicide on ultra structure (SEM) and enzymatic activity in different regions of alimentary canal and gill of *Channa punctatus* (Bloch) *Journal of Crop and Weed*. 2009;5(1):236-245.
- ¹²⁵ Shehata, A.A., Schrödl, W., Aldin, A.A., Hafez, H.M., Krüger, M. (2013). The effect of glyphosate on potential pathogens and beneficial members of poultry microbiota in vitro. *Curr. Microbiol*. 2013 66, 350-358.
- ¹²⁶ Monika Krüger, Awad Ali Shehata, Wieland Schrödl, Arne Rodloff. Glyphosate suppresses the antagonistic effect of *Enterococcus* spp. on *Clostridium botulinum*. *Anaerobe*. 2013 Feb 6. Epub 2013 Feb 6. PMID: 23396248; see also Monika Krüger, Wieland Schrödl, Jürgen Neuhaus and Awad Ali Shehata. Field Investigations of Glyphosate in Urine of Danish Dairy Cows, *Journal of Environmental & Analytical Toxicology*, <http://dx.doi.org/10.4172/2161-0525.1000186>. R. Rulff, W. Schrödl, S. Basiouni, J. Neuhaus, M. Krüger. Is downer cow syndrome related to chronic botulism? *Polish Journal of Veterinary Sciences* Vol. 18, No. 4 (2015), 759-765. <http://doi10.1515/pjvs-2015-0098>; Ramón Rulff, Shereen Basiouni, Monika Krüger. Biochemical Investigations and Glyphosate Detection in Downer Cow Syndrome. *International Journal of Advancements in Research & Technology*, Volume 5, Issue 1, January -2016 ISSN 2278-7763; Monika Krüger, Awad Ali Shehata, Wieland Schrödl Arne Rodloff. Glyphosate suppresses the antagonistic effect of *Enterococcus* spp. on *Clostridium botulinum*. *Anaerobe* 20 (2013) 74-78; Gerlach H, Gerlach A, Schrödl W, Haufe S, Schottdorf B, (2014) Oral Application of Charcoal and Humic Acids Influence Selected Gastrointestinal Microbiota, Enzymes, Electrolytes, and Substrates in the Blood of Dairy Cows Challenged with Glyphosate in GMO Feeds. *J Environ Anal Toxicol* 5:256. doi: 10.4172/2161-0525.1000256; Gerlach H, Gerlach A, Schrödl W, Schottdorf B, Haufe S et al., (2014) Oral Application of Charcoal and Humic acids to Dairy Cows Influences *Clostridium botulinum* Blood Serum Antibody Level and Glyphosate Excretion in Urine. *J Clin Toxicol* 4:186. doi: 10.4172/2161-0495.1000186
- ¹²⁷ Shehata, A.A., Schrödl, W., Aldin, A.A., Hafez, H.M., Krüger, M. (2012). The Effect of Glyphosate on Potential Pathogens and Beneficial Members of Poultry Microbiota In Vitro. *Curr Microbiol*. 2012 Dec 9. Epub 2012 Dec 9. PMID: 23224412
- ¹²⁸ Krüger, M., Große-Herrenthey, A., Schrödl, W., Gerlach, A., Rodloff, A. (n.d.). Visceral botulism at dairy farms in Schleswig Holstein, Germany e Prevalence of *Clostridium botulinum* in feces of cows, in animal feeds, in feces of the farmers, and in house dust. *Anaerobe*, in press; and Böhnel H, Behrens S, Loch P, Lube K, Gessler F. Is there a link between infant botulism and sudden infant death? Bacteriological results obtained in central Germany. *Eur J Pediatr*. 2001 Oct;160(10):623-8.
- ¹²⁹ Arnon, SS, Midura, TF, Damus, K., Wood, RM, Chin, J. (1978). Intestinal infection and toxin production by *Clostridium botulinum* as one cause of sudden infant death syndrome. *Lancet*. 1978 Jun 17;1(8077):1273-7.
- ¹³⁰ Fasano, A. (2011). Zonulin and Its Regulation of Intestinal Barrier Function: The Biological Door to Inflammation, Autoimmunity, and Cancer. *Physiol Rev* 91: 151-175, 2011. <http://www.ncbi.nlm.nih.gov/pubmed/21248165>
- ¹³¹ Shin-Ichiro Miyashita, Yoshimasa Sagane, Ken Inui, Shintaro Hayashi, Keita Miyata, Tomonori Suzuki, Tohru Ohyama, Toshihiro Watanabe, Koichi Niwa (2013). Botulinum Toxin Complex Increases Paracellular Permeability in Intestinal Epithelial Cells via Activation of p38 Mitogen-Activated Protein Kinase. *J Vet Med Sci*. 2013 Jul 25. Epub 2013 Jul 25. PMID: 23884081
- ¹³² Goldstein, J., Morris, W., Loidl, CF, Tironi-Farinati, C., McClane, B., Uzal, F., Fernandez Miyakawa, M. (2009). *Clostridium perfringens* epsilon toxin increases the small intestinal permeability in mice and rats. *PLoS One*. 2009 ;4(9):e7065. Epub 2009 Sep 18. PMID: 19763257
- ¹³³ Yong-guo Zhang, Shaoping Wu, Yinglin Xia, Jun Sun (2013). *Salmonella* infection upregulates the leaky protein claudin-2 in intestinal epithelial cells. *PLoS One*. 2013 ;8(3):e58606. Epub 2013 Mar 11. PMID: 23505542
- ¹³⁴ Vasiluk, L., Pinto, LJ, Moore, MM (2005). Oral bioavailability of glyphosate: studies using two intestinal cell lines. *Environ Toxicol Chem*. 2005 Jan;24(1):153-60.
- ¹³⁵ Sikander, A. (2009). Role of serotonin in gastrointestinal motility and irritable bowel syndrome. *Clin Chim Acta*. 2009 May;403(1-2):47-55. doi: 10.1016/j.cca.2009.01.028. Epub 2009 Feb 4.